of site and area in a cardiac *bruit*, the nature of the means by which we seek for evidence that may turn the scale is momentous in the extreme. It is strange that such issues have been left so long to hang upon the clumsy devices of our forefathers.

This description of the compound stethoscope, with regard to the prime conditions it fulfils, will not fail to suggest abundant illustrations of its advantages as an instrument for clinical use. I will conclude by a reference to one or two points, as I think, of much practical consequence. It is no small advantage that both ears are completely closed to extraneous sounds when the stethoscope is in use; the ears are engaged only by the sounds which it is desired to hear. Moreover, the attention is not diverted during the shifting of the collectors from point to point. Ease of position to the auscultator and patient are secured; it is possible to auscultate patients whose decubitus would preclude the use of the uniaural stethoscope, and the comfort of the patient who is in pain is eminently consulted. Apart from the fact that attention in hearing is liable to distraction when, at the same time, the attention is solicited in other directions through the eye, the sense of hearing is much aided and the evidence furnished to the ear is much corroborated by coincident inspection of the chest by the eye. I have frequently derived assistance, in determining the relations of a cardiac murmur to the sounds it has wholly or partially replaced, from having the cardiac impulse before the eyes. It is often essential to diagnosis that the point of maximum intensity of sounds should be accurately localised. To this end, means of making rapid exploration of the areas over which the sounds are heard, and means of marking the results of comparisons as they are successively made, are the desiderata. It is needless to point out how these are supplied by the two component stethoscopes of my instrument, either of which can be brought into use or rendered impervious instantly, and without disturbing the adjustment to the ears. Finally—and it is the student that this illustration chiefly concerns—the first recognition of a morbid sound depends on being able to compare it with its healthy correlative, and the more rapidly the one can be exchanged for the other in the ears the better will be the appreciation of the difference. Or, take the case of a sound which is morbid, not from its special character, but because it is heard in an abnormal position, a similar rapidity of exchange and comparison is required. In most patients, standard healthy sounds coexist in the same chest with sounds that are morbid both as to character and site. One collector being applied over the site of a healthy standard and the other over the site of a morbid sound, the sounds can be admitted to, or shut off from, the ears alternately, and thus compared under conditions which could hardly fail to produce accurate impressions. During an extended use of the instrument in the wards and out-patient rooms of a large hospital, its ready response to the endless and varied calls made upon it by the exigencies of clinical practice justify the expectation that it will stand the test of experience.

ON HYPEREXCITABILITY AND PARESIS.

By C. HANDFIELD JONES, M.B., F.R.C.P., F.R.S.,
Physician to St. Mary's Hospital, etc.

[Concluded from p. 372 of last number.]

THE next point on which we shall remark is the affinity between hyperexcitability and paresis. This has already been illustrated, to some extent, in the preceding remarks, but it deserves to be noticed more particularly.

1. Commencing with the faculty of sensation, we find, first, that the most marked hyperæsthesia is produced by conditions which also give rise to anæsthesia. Thus, in Dr. Waller's experiments on refrigeration of the ulnar nerve, the ultimate effect of the chilling was anæsthesia of the district supplied by the nerve; but this was preceded by hyperæsthesia. A case is recorded in the *Medico-Chirurgical Transactions*, vol. vii, p. 177, where, during the gradual recovery of the brachial plexus from injury produced by a fracture of the clavicle, the skin of the upper part of the arm possessed natural sensibility, another (probably the mid-region) was morbidly sensible, while that immediately beyond was quite insensible. Trousseau (*Clinique Médicale*, vol. ii, p. 312) states that cutaneous anæsthesia is often seen to succeed hyperæsthesia, especially in cases of herpes zoster. Secondly, inflammation of a nervous centre, as the spinal cord, giving rise to motor paralysis, will also give rise to hyperæsthesia, the two conditions existing together. In a case of alcoholic paralysis which I have recorded, complete motor palsy of the lower limbs coexisted with extreme hyperæsthesia and pain. Her powerless hands were numb, but not in nearly so much pain as the feet. Here the three phenomena were manifestly co-products of the same cause, and at the same time. Anæsthesia, however, is the more usual accompaniment

of palsy. Trousseau (*Clinique Médicale*, vol. i, p. 287) mentions, speaking of a case of complete paraplegia, that the anæsthesia, which had been considerable up to a certain date, was replaced the next day by hyperæsthesia.

2. *Hyperæsthesia is often associated with pain*, as in abdominal neuralgia, the hysterical breast, and irritable testis. In Dr. Fuller's case of obstruction of the arteries of the lower limbs with fibrinous deposit, the first symptom was acute pain, associated with exquisite tenderness of the whole limb, so great that the slightest touch caused intolerable suffering. Now, pain is regarded by so good an authority as Dr. Anstie as ranking on the same level as paralysis.

3. The remedies for hyperæsthesia are mainly of a restorative and recreative kind—not quite so absolutely, indeed, as those of neuralgia, but still so much so as to indicate not obscurely the nature of the process. Those who admit the truth of the adage, "Naturam morborum remedia ostendunt," will not think lightly of this argument.

Motor hyperexcitability exhibits a relation to paralysis in the following particulars. 1. The same conditions which usually produce paralysis of a motor centre, will also produce hyperexcitability. Thus, in a case of myelitis under my care, the left leg was simply paralysed, as well as the sphincters; but the right leg started and jumped, and was affected with twitching sensations. In another case, where there was very considerable paralysis of both motor and sensory power in the lower limbs, and the power of the sphincters was impaired, strong tremors were produced by passive movements of the legs, or by attempts to move them. As improvement took place, the tremor ceased. In a third case, where there had been complete paralysis of motion and sensation, in connexion with diseased vertebrae and abscess, the left thigh, at the time when I saw the patient, was affected with considerable tremor when he attempted to use the limb. In a fourth case, where motor power, which at first was quite lost, had been very much regained, there was great tremor of the lower limbs. In all these instances, the damaged nerve-tissue was hyperexcitable. A case of lead-poisoning was once under my care, where the symptoms were not the ordinary paralytic, but were decidedly indicative of hyperexcitability. The patient's arms were agitated by clonic convulsions when he attempted to use them. He could scarcely carry a glass to his lips without spilling the liquid, or hold it so as to drink. At one time, his arm used to be fixed to his side by tonic spasm for several hours every day. If he put himself out in any way, he shook violently, "like a leaf on a tree," and his gait became unsteady and jerky. He improved very much with strychnia, cod-liver oil, and faradisation. Here the motor centres were certainly affected by the poison; but the result was hyper- instead of non-excitability. The remedies which availed were those which serve us when paralysis exists. Surely all this plainly points to a close connection between the two states just mentioned. Regarding embolism as the cause of chorea, which it probably is in some instances, it will not be forgotten that the same cause gives rise to paralysis when it obstructs the blood-flow to the corpus striatum. Dr. Lockhart Clarke has recorded a case (*British and Foreign Medico-Chirurgical Review*, July 1864) in which there was partial paralysis of one arm; and so long as the arm was enveloped and kept warm in flannel, it remained perfectly still; but whenever it was exposed to the impression of cold air, it began to move in a spasmodic and jerking manner. Here the lesions and disintegrations found in the cord bore a very striking resemblance to those occurring in tetanus. 2. When hyperexcitability exists in one part, there is often also failure of nervous power in another. In a case where the sural muscles of the left leg contracted spasmodically, so that the patient walked on the point of his toes, and there was some permanent rigidity of both the flexor and extensor muscles of the ankle, the mental faculties were impaired; he was mopy, dull, listless, and more or less drowsy. These cerebral pareses had obliged him to quit his occupation. 3. In chorea, Trousseau states it as the general rule, that the intellectual faculties are more or less weakened as long as the motor disorder continues. He also affirms that anæsthesia or other derangements of the sensory faculty are present in the majority of instances. A medical friend related to me a case of tænia under his care, where there existed paralysis of one leg and spasmodic contraction of the opposite arm. Mr. Moodeen Scherriff has recorded some instances of paralysis not depending on organic lesion, in which there was rigidity of one limb and simple immobility of the other on the same side, the same cause having given rise to both conditions, and the same treatment removing them. Dr. Wilks mentions a case (*Guy's Hospital Reports*, 1866, p. 261) where, after a mental shock, a girl aged 15 passed into a semi-maniacal condition. Her face and hands were constantly affected by choreal movements, which at times became almost tetanic. After her menstrual period passed by, she became calmer and more like an ordinary choreic. She appeared to understand what was said to her, but was perfectly anæsthetic. Dr.

Wilks tested her condition in many ways, and was convinced that sensation was altogether absent. In another of his cases, the patient was cataleptic, and had lost all sense of feeling, either from true anaesthesia or from unconsciousness. In all these instances, hyperexcitability of some motor nerve-apparatus coincides with paresis of other centres, either motor, sensory, or intellectual. The latter defect, being more easy to interpret, throws light on the nature of the former. 4. There is little doubt, I think, that motor hyperexcitability may sometimes be very beneficially treated by tonic and recreative remedies—by means, as Dr. Radcliffe states, which are calculated to exalt vital energy in general, and nervous energy in particular. He has seen two cases in which strong and general tetanic spasms relaxed very rapidly under a treatment of which the essential part was to give wine to a point just short of inebriation. He knows of four cases where attacks of laryngismus stridulus, which were frequently recurring under a treatment in which it was thought essential to regulate the secretions by grey powder and so on, were put a stop to once and for all by a tonic and restorative plan of treatment—wine, beef-tea, steel, chloric ether, etc.; and he can speak with confidence of the beneficial results of a similar plan of treatment in a considerable number of cases of whooping-cough. Though the above experience applies to a good many cases of this kind, it certainly does not to all; and this argument must, therefore, not be overstrained.

In the intellectual organs, hyperexcitability is evidently not remote from paresis in the downward descent. The two conditions, we may say, overlap one another; so that, while at one period hyperexcitability may predominate, in another it is more or less completely replaced by paresis. Delirium is sometimes of such a quality as implies the extremest activity of the cerebral nerve-cells; at others, it is a mere mauling, but little removed from complete stupor; while in both states it is to be observed that the real functional power of the organ is totally lost. The same poison—say that of typhus—at the same period of its action, may produce either active and maniacal delirium, or low and muttering. In severer grades of the disease, the muttering delirium passes into a heavy stupor, or into coma vigil. It does not seem possible to account for the objective difference in the symptoms by any demonstrable state of the brain, for Dr. Buchanan writes (Reynolds's *System of Surgery*) that the nervous system commonly shows, after death, no lesion whatever to account for even intense head-symptoms. Not only does the hyperexcitability in this disease culminate in stupor, but, as in other fevers, mental languor and dulling of the intellect actually precede the occurrence of delirium. The two phases of nerve-disorder are more or less blended, in fact, throughout. In common drunkenness, and also in chloroform and belladonna intoxication, we have the same sequence of disorder as in typhus. Noisy more or less excited inebriety precedes sopor and unconsciousness. There seems to be good reason to believe that, in these and in analogous cases, hyperexcitability is the earlier and milder effect of the toxic agent, and paresis the later and severer. If the dose of alcohol be small, no effect may ensue except a moderate amount of excitement, attended with increase rather than with diminution of power. If it be very large, coma is rapidly produced, without previous excitement. Epilepsy affords some notable evidence of the close connection between hyperexcitability and paresis. The usual sequel of the "grand mal" is coma of longer or shorter duration; but in some instances this is replaced by fierce delirium, which is hard to tame even by chloroform. There can be no question that the circulation is free in both morbid states. We can only refer the difference in result to some peculiarity in cerebral constitution. This shows how closely allied the two derangements really are. In the paroxysm itself, we cannot help noticing the coincidence of extreme hyperexcitability in the motor centres, with total paresis in the intellectual. Again, in chorea, outbreaks of maniacal delirium are not very infrequent, though the usual condition of the brain approximates much more to torpor. In a case of diphtherial poisoning I saw lately, after the false membrane had disappeared, the patient became semicomatose, with dilated pupils, and marked but incomplete ptosis of the right eye. He was well fed and stimulated, and became rational, but soon after was violently delirious in the night, and continued so next day. Subcutaneous morphia quieted him, the delirium disappeared, and for some days he seemed pretty well; then coma suddenly came on again, and proved fatal. Here the toxæmia caused two attacks of paresis, with an intervening paroxysm of hyperexcitability. Dr. Cheadle records (BRITISH MEDICAL JOURNAL, 1871, vol. ii, p. 545) a case of starvation from extreme cirrhosis of the liver. The later symptoms were great dulness of intellect, with jerking of left arm and head, succeeded by wild delirium with shrieking, and, lastly, right hemiplegia, coma, widely dilated and insensible pupils, and continued jerking of the left arm and leg. Nothing was found in the brain at the autopsy to account for the cerebral symptoms. Here signs of excitement and of

paresis are evidently due to the same cause, and occur in close connection.

But, closely allied as these derangements are, it is quite certain that *they are not inseparable*. Hyperexcitability may occur in its most marked form, and not be associated with, or followed by, any paresis. Of this we have frequent instances in ordinary tetanus. Cardiac palpitation, asthma, photophobia, are others. The same may be affirmed of paresis: that it may be quite unattended with any signs of hyperexcitability. Of this, diphtherial paralysis affords a good example, as also do common instances of coma.

The *diverseness of hyperexcitability from paresis* is strongly shown in the cases where the former is successfully treated by large doses of depressants, as antimony, tobacco, aconite. In these, it is of course essential that the heart should be able to withstand the depressing influence; that it should not be paretic, nor its ganglia. If this be the case, the hyperexcitability of the brain or the cord may be very beneficially reduced by depressing agents. Of course, if the heart be feeble, they are not to be thought of. Sometimes the hyperexcitability of the nervous centres seems to confer an increase of power on the heart for the time, rendering it capable of withstanding agents which, when the nervous centres have regained their normal state, would dangerously depress it. This diverseness is also shown by the common experience that in many of the neurotic disorders, especially in females of the middle and upper classes, where there is also great debility, no drug of a tonic kind is tolerated, it aggravates the hyperexcitability so much. These facts certainly show that the morbid state conditioning hyperexcitability is very diverse from that belonging to paresis. One consists in hurried imperfect vito-chemical change; the other, in cessation of normal change.

A few remarks on sedatives may appropriately follow this discussion of hyperexcitability. These remedies differ much more from one another than the several tonics do. In fact, some of the sedatives approximate closely to the stimulants or tonics. Digitalis, which is a tonic to the heart, may be an efficient sedative to the brain, quieting delirium and promoting sleep. Opium, famous for its *vis dormitiva*, is scarcely less so for its sustaining and invigorating effect, as well as its tonic action on nasal nerves. Indian hemp, while producing sedative effects, is also stimulating and exciting. The same may be said of alcohol. That this should be so, appears no mean argument in favour of the view that most derangements of nerve-force partake very much of the nature of paralyses. I should distribute the so-called sedatives into three groups. The first, comprising opium, alcohol, camphor, and Indian hemp, might be termed that of the stimulating sedatives; the second, comprising belladonna, henbane, conium, bromide of potassium, chloral, and chloroform, might be denominated that of the simple sedatives; the third, comprising physostigma, tobacco, aconite, and antimony, that of the depressing sedatives. The second group show a manifest approximation to the third, in that they are apt to produce actual paralysis when given in large doses. It is clear that, in the choice of a sedative to combat hyperexcitability, one must be guided very much by the amount of coexisting paresis manifested either in the nerve-centres or the heart. Doseage is also to be carefully regarded, a large dose often acting as an effective sedative, when small ones prove exciting or inoperative.

Finally, it may be remarked that the existence of these two great forms of nerve-disorder, which, though closely allied, as we have seen, and often more or less blended together, yet are often met with quite distinct and separate, goes far to show that nerve-apparatuses possess two faculties, either of which may be impaired without the other suffering. One of these is the faculty of *evolving force* on the solicitation of the will or a physical excitant; the other that of *regulating the evolution of the force*, so that it shall not take place, so to speak, in an explosive manner, but in due order, according to the requirements of the function. In delirium, mania, and convulsions, the second of these faculties is lost; in dementia, coma, and paralysis, the first. Both hyperexcitability and paresis, therefore, imply failure and loss of vital power, though of a different kind. It is, therefore, quite what we should expect that both are often produced by the same causes, and that they should often exist together. On this view it is quite intelligible why "excitement without power" is of so frequent occurrence. If we apply this view to therapeutics, we foresee that there will be (1) Cases of pure paresis, mental, motor, or sensory, to which the tonic and stimulant regimen in its entirety is quite appropriate; (2) Cases of pure hyperexcitability where different remedies will be requisite, viz., the depressing sedatives; (3) A large number of mixed cases, where the treatment must also be of a mixed kind, according as the symptoms indicate one or other defect to predominate. All this has been arrived at by empiricism, but I think we see now that it is a rational procedure.

From what has been advanced, I think it may be concluded—I. That hyperexcitability and paresis do not depend principally on altera-

tions in the amount of blood-flow; 2. That they do depend on alterations in the nerve-tissue effected, as all neuroses are, either by exhaustion, toxæmia, or remote irritation; 3. That the morbid processes giving rise to these states are diverse, and yet have much affinity, and may exist together; 4. That hyperexcitability, as a necessary result, tends to produce more or less paresis by exhaustion; 5. That paresis probably consists in slackening and cessation of normal vital chemical change, hyperexcitability in setting up of abnormal change, the former being comparable to an atrophy, the latter to a morbid growth; 6. That the therapy of either morbid condition needs to be modified by regard to the amount of the other coexistent with it.

THE DENTITION OF THE KOSTROMA PEOPLE.

By CHARLES S. TOMES, M.A.; and
OAKLEY COLES.

THE Kostroma people now being exhibited in London, independently of their hirsute appendages, are of great interest on account of the remarkable condition of their jaws.

The man, Adrian, has in the upper jaw only one tooth, apparently a lateral incisor, though Virchow describes it as a canine; whilst in the lower jaw he has four incisors, or probably three incisors and a canine; and, from their texture, these teeth are doubtless of the second dentition. The man gives no account of any milk-teeth; but it is only rea-

Fig. 1.—Adrian.

Fig. 2.—Feodor.

sonable to suppose that those at present existing were preceded in the usual way by the deciduous teeth. The boy, Feodor, has an edentulous upper jaw, and but four (temporary) incisors in the lower jaw. These

Fig. 3.—The Man's Upper Jaw.

Fig. 4.—The Man's Lower Jaw.

are fairly developed, and well placed as to position. The accompanying engravings, taken from photographs, give an excellent idea of the

personal appearance of the "hairy men"; whilst the woodcuts below show the jaws of Adrian and Feodor. The alveolar ridges of both jaws in the two subjects of this notice are not at all fully developed, and do not convey an impression of the teeth being retained within their osseous crypts, but rather of an absence of the dental organs altogether. Taking the face in its entirety, we observe that the jaws are the only parts that have apparently been dwarfed in the process of development, the malar region being normal in character. The angle of the lower jaw has, so far as can be ascertained, retained its natural

Fig. 5.—The Child's Upper Jaw.

Fig. 6.—The Child's Lower Jaw.

form, so that, when the mouth is closed, the lower teeth, lip, and chin project in much the same way that they would do in the case of a patient who had been deprived suddenly of all the teeth with the exception of the lower incisors.

A comparison of the upper jaws of the man and the boy with one another, and with normal jaws of various known ages, in which the Museum of the Odontological Society is very rich, reveals some relations of interest. The upper jaws were selected for this purpose, as there is, unfortunately, some doubt as to the accuracy of the model of the lower jaw of the man; that of the lower jaw of the boy is certainly accurate. The alveolar arch in Feodor's (the boy's) mouth corresponds very closely with that of a well grown child four years of age; that is to say, it is of just the same size and shape as an alveolar arch, which contains all the deciduous teeth in their places. Its posterior cornua fall short of those of children of five years, inasmuch as these latter are prolonged behind the last deciduous molar, to give space for the forming first permanent, or six-year-old molar. The alveolar arch in the man differs but slightly either in shape or size from that of the boy. Owing to its being rather less defined than that of the boy, it is difficult to measure it with the same accuracy; but it appears to exceed it very slightly in antero-posterior length, and only a little more in width. As has been already noted, this applies only to the alveolar border, and not to the whole upper maxilla. An abrupt bulging is noticeable in the upper part of the model, in the sulcus between the cheek and the alveolar border; and the malar processes are of large size, so that the face is broad and heavy.

It is, of course, unsafe to draw general conclusions from such scanty data; yet the relations existing between these two jaws seem to give some confirmation to the views as to the growth of the jaws set forth by Mr. Tomes, sen., and subsequently confirmed by Professor Humphry of Cambridge; for they indicate that the sole difference between the child's jaw and that of an adult is (disregarding a slight increase of width due to the greater breadth of the alveolar processes) a backward elongation of the cornua, strictly dependent on the successive addition of the posterior molars. For here is the alveolar portion of the jaw of a bulky man, the rest of whose upper maxilla is in no degree stunted, so far as we can judge, barely exceeding in size that of his child; and the explanation seems obviously to be that, as no posterior molars have been added, so no posterior elongation of the jaw has been effected. And the fact that the jaw of the boy has attained to its present size and form would seem further to indicate that the growth of the jaw up to a certain point is independent of the teeth, but that, after that point, its enlargement, which is actually little else than an elongation of its cornua backwards, is strictly and absolutely dependent on the successive additions of teeth posteriorly.

It should be added, to save misunderstanding, that the foregoing remarks apply solely to the form and size of the arch. The alveolar process, which is strictly dependent on the teeth which it carries for its depth, is in this respect necessarily deficient.