If, in reference to the possibility of overlooking the nervous phenomena, or, in order to make assurance doubly sure, you require collateral evidences of approaching danger, I will mention a few symptoms which are held to be of peculiar significance. A dry unspiring skin; a profuse and intensely red miliary rash; an abatement of the pain and swelling in the joints, or their total disappearance; diarrhoea, and a copious flow of pale alkaline or neutral urine, have all been observed to accompany or to precede the paroxysms: they may even precede the nervous prodromata, specially so called. In our case, a free flow of urine occurred before delirium, and diarrhoea occurred with it. The articular pains, for many days before the outbreak of delirium, appeared in all the joints, and disappeared from all in the most desultory way. Again, during the four days prior to that outbreak, there was a slow rise in the temperature, which on the fourth evening culminated at 104.2 deg.; the pulse-rate, also, during the same period, ruled higher than before; but the complexity and continued severity of the case seemed sufficient to explain these occurrences. An unexplained rise in either direction would certainly be a suspicious symptom: perhaps even a rise of any kind should put us on our guard. Unfortunately, the collateral evidences are all inconstant, but they are useful auxiliaries, and when they appear they ought to awaken attention. Let me remark, by way of parenthesis, that, as I should be vigilant myself, so I should inculcate the extremest vigilance on every one entrusted with the care of the patient; and I should direct the attendants, on the appearance of anything strange in his aspect or demeanour, to use the thermometer assiduously, and, if ever it rose to 105 deg., to send at once for the physician.

Gentlemen, I have laid down the law that, for all practical purposes, as far as the bath is concerned, we must take the thermometer for our guide. I wish, in some measure, to modify this hard and fast rule. So singularly calming is the bath, even when there is no great exaltation of the body-heat, that I should be disposed to administer it as a simple palliative in all exacerbations of delirium and restlessness, although the heat may have fallen far below its point of maximum intensity, and may register at the time no more than from 102.5 to 104 deg. in the axilla. At a lower level than the above it would not be desirable to use the bath, for such temperature would not allow a sufficient range to cover both the fall within the bath, which we can regulate, and the fall after the bath, which we cannot regulate with exactness; although, from a comparison of Dr. Wilson Fox's cases, I should roughly reckon the average for the after-fall at about four degrees. On the foregoing plan our last bath was administered, when the temperature was no higher than 103.1 deg. in the axilla, and it accomplished its purpose. Perhaps I might have adopted the same plan with advantage during the prolonged agitation which preceded the climax. In this connexion I would remark that, with a temperature descending gradually from 95 or 90 to 70 deg. or thereabouts, the bath is at least as much a warm bath as a cold bath; and warm baths have for ages been recognised as sedatives to the nervous system.

Before dismissing the subject of the bath, it is important to observe that the chest-complications, severe and extensive as they were, including pneumonia, pleurisy, bronchitis, and pericarditis, underwent no perceptible change for the worse in consequence of its use.

With regard to medicinal remedies, opium and hydrate of chloral are the only soporifics and sedatives of any real value. They may be given freely enough during the premonitory period, or stage of suspicion, to control mere delirium or restlessness; but if I saw the danger-signal in the thermometer, I should put on the break at once and stop the opium for the time being. I might resort to it again if, after the subsidence of the hyperpyrexia, the agitation persisted, and chloral failed to soothe it. Under these circumstances, we cannot afford to dispense with opium altogether; but let it never be forgotten that there is no sharp and unvarying line of distinction between the comatose and the delirious class of symptoms; that stupor may alternate with excitement, and that death by coma is the natural termination of the attack.

In cases characterised by stupor from the first, with or without alternations of excitement, you may apply ice to the forehead or scalp, or the so-called revulsives to the trunk and extremities. I confess I should be afraid of blisters, wherever applied. As the symptoms progressed, I should trust principally to the early employment of the bath, as before stated. In all cases, of whatever type, I should feed and stimulate the patient largely, and I should take especial care not to weaken him in any way.

You are aware that modern researches have established for quinine a high reputation as a refrigerant; but in all the cases that I have perused, I fail to find any illustration of this reputed power. In my own case, as the boy happened to be doing well, on the principle of letting well alone I was unwilling to alter one iota of the treatment, and so he remained for days and weeks under the influence of quinine,

whatever that might be. I cannot determine its value; and the same thing is true of the digitalis associated with it. The possible production of cinchonism is a great drawback to the use of quinine, and might embarrass the prognosis in a class of cases where deafness may be a spontaneous symptom, and a dangerous one.

Gentlemen, in this lecture I have enlarged in great measure on the premonitory symptoms, and passed lightly over the features of the developed attack. With the addition of hyperpyrexia, the last stage is, in the main, the climax and complement of the first. The symptoms may be more conspicuously typhoid in character; there may be convulsions or cyanosis; and there is sure to be coma. I have preferred to dwell upon the premonitions, because the great practical point is to discern the nascent mischief at the moment of its possible origin, and to arrest its ulterior evolution.

I have not ventured formally to affix the name hyperpyrexia to my case: indeed, I know not how to define hyperpyrexia exactly; but this I know, that no one who saw the patient on the night of his first bath would hesitate to say that, without the bath, he must have died in the course of a few hours, with hyperpyrexia on any definition.

In conclusion, I beg to acknowledge the great services rendered by the resident medical officer and all the clinical assistants, without whose aid the patient would never have lived.

CLINICAL LECTURE ON INJURIES FROM LIGHTNING.

BY JAMES R. LANE, F.R.C.S.,
Surgeon to St. Mary's Hospital.

GENTLEMEN,—Opportunities for witnessing the effects of lightning on the human body are not in ordinary seasons very common in this country; and the cases of which I propose to speak to you to-day are well worth your attention, not only from their rarity, but also on account of the many curious and interesting circumstances connected with them.

On the afternoon of Tuesday, June 18th, two men, having with them a horse and cart, were at work, road-making, in a field near Willesden. A severe thunder-storm coming on, they took shelter under a large oak-tree growing in an earthen bank, with no hedge or other tree within a considerable distance. The horse and cart stood near, but not under, the tree. The men were close together, but not touching the trunk of the tree. Suddenly they were both struck down; one was rendered completely insensible; the other did not lose his senses, but was unable to rise from loss of power in his lower limbs. Some men who saw what had occurred went to their assistance, and, as soon as a conveyance could be procured, they were brought to the hospital. When the men were struck, the horse took fright and bolted with the cart, but neither horse nor cart sustained any injury. A large branch was torn off from the trunk of the tree.

One of the men was but slightly injured. When admitted, he appeared to be suffering from shock; his pupils were dilated, and he was unable to stand. The loss of power, however, was confined to the lower extremities, on both of which an erythematous blush was seen extending from the knee downwards. There was nowhere any breach of surface or ecchymosis, and his clothing was uninjured. This man's symptoms soon subsided; by the next morning the redness was gone, and the paralysis had completely passed off. He said he felt quite well, and was allowed to leave the hospital.

With the other man the case was very different. He is reported to have been quite unconscious for about ten minutes after the occurrence; then he recovered himself for awhile, but again lost his senses, probably from faintness. He says that he remembers feeling a violent blow on the head, which made "his eyes strike fire". When admitted, he was conscious, but in a state of collapse, and his pupils were widely dilated. He had bleeding from the right ear, the external part of which was of a dark purple colour, and seemed both excoriated and bruised. Some of the hair above the ear, and also the right whisker, had been burnt away. From the whisker there was an ecchymosis extending down the neck, and ceasing by forming a narrow circular line round the back of the neck, corresponding to the position of a silver watch-chain which he was wearing when he was struck. From this point, down to the right groin, there was no sign of injury; but from the groin downwards there was a purple discoloration of the skin, about two inches wide, passing round the outside of the thigh to

the back of the knee and thence to the heel. Along this track the hair had disappeared, having apparently been singed off.

The condition of his clothing, which you see on the table, and which is to be preserved in the Museum, is very remarkable. His cloth cap has a large ragged hole in it on the right side, as if a piece had been torn out; but some padding of cotton-wool, and also the inner lining of the cap, are uninjured. The side of the cap was, no doubt, the point first struck, and thence the lightning passed over the ear and down the neck till it met with the silver watch-chain, which, from its greater conducting power, carried it onwards till it reached his watch, which was in the right pocket of his trousers. From this point the current travelled down his right thigh and leg to the ground, leaving the marks on the skin already referred to.

His calico shirt is blackened and burnt into several small holes round the back of the neck. It is blackened also in wide patches down the front, where the silver watch-chain rested; and here at one point it is burnt into a large hole. The front flap of the skirt is torn to shreds, but neither burnt nor blackened.

A red woollen jersey, which he wore under the shirt, is blackened, and evidently singed at a point corresponding to the hole in the front of the shirt, but not burnt completely through. The lower part is torn into shreds, both in front and behind. Both shirt and jersey were smouldering when he was picked up.

The silver watch-chain is broken into fragments. One piece about six inches long was found, and also several smaller pieces; the rest is missing. The large piece shows signs of partial fusion in places; the smaller ones are more completely melted. The relation of the fused watch-chain to the black patches on the shirt is interesting. These patches at first were thought to have been caused by scorching; but it seems more probable that they result from an impregnation of the texture by metallic silver in a state of minute subdivision, for the analysis by Dr. Wright of a piece about half an inch square indicated the presence of silver in considerable quantity.

As to the watch, the glass was broken into small pieces. The face is partially driven in, and is stripped of its white enamel. The hands are twisted and broken. The silver case shows signs of fusion in places, but the works do not seem to be injured. The watch was forced through a hole, partly burnt and partly torn, in the pocket, and through the torn trousers; for it was found detached from the chain and lying on the ground at some distance from the man's side. Some money, which was in a leather purse in the left trousers pocket, does not appear to have been touched.

The right leg of his fustian trousers is ripped open along the seams from the top to the bottom, but not burnt. The calico lining is torn to shreds.

The back of his right stocking is singed above, but torn open below, whence a large piece appears to have been blown away. The back and outer side of the right boot have also been blown away; there are holes in the heel, from which several large nails have evidently been forcibly driven out; but the majority of the nails remain undisturbed.

Such was the condition of the man and his clothing when brought to the hospital. His subsequent progress has been, on the whole, favourable. He soon recovered from the collapse; but the shock had evidently been severe, for he remained for several days in a very weak and languid condition. It was found that he was deaf in the right ear; an examination of which, by Dr. Peter Allen, showed a laceration of the membrana tympani; and when he makes an effort at expiration with the nose and mouth closed, you may hear the air whistling through the auditory meatus. On July 1st, he complained of severe pain in the ear and over the mastoid region, and this was accompanied by quick pulse, hot skin, and other feverish indications, so that we feared that some meningeal complication, arising out of the injury to the tympanum, might be setting in; but on the following day there was a free purulent discharge from the ear, and these unpleasant symptoms subsided. Complete deafness in this ear, however, still remains, which renders it probable that the nervous textures as well as the membrana tympani have suffered. The dark marks on the neck and right thigh gradually passed away. They were evidently ecchymoses, and not burns, as was at first supposed. The only sign of burning, with the exception of the scorched hair and some vesications on the ear, was found on the right thigh, in which a small piece of fused watch-chain was imbedded.

The operation of the lightning on the man's body in this case seems to have been in the nature of a dull heavy blow. In its passage it produced, for the instant, sufficient heat to scorch the hair, but not to burn the skin. It must have exercised at the same time great rending power in the outward direction, as indicated by the condition of the torn clothing. From the neck to the groin it left no mark, this interval having been protected to a great extent by the conducting power

of the watch-chain; silver being, in fact, the best of metallic conductors. The chain, however, had not an area sufficient to permit so intense a current to pass without injury to its own texture by fusion and pulverisation, and consequent scorching of the shirt and jersey on which it rested.

Another curious phenomenon presented by this patient, to which I have not yet alluded, is a partial paralysis, together with a complete, or almost complete, absence of pulse in the left upper extremity, although this part shows no outward sign of injury. Soon after his admission, he complained that his left arm felt weak and numb, and he was unable to grasp anything firmly in his left hand. No pulse could be felt at the wrist, or in the brachial or subclavian arteries; and, in probable relation with this latter circumstance, there is a loud *bruit* to be heard below the sterno-clavicular articulation, over a limited area corresponding in position to the first part of the left subclavian artery. It was at first thought that this might depend on an aneurism involving the subclavian; but it is not probable that this is the case, for there is no dulness on percussion over the spot where the *bruit* is most distinctly heard, and none of the pressure-effects which an aneurism would be likely to produce have been observed. The *bruit* varies in intensity in different positions of the body; but change of position makes no difference in the pulse. The paralysis has gradually passed off, and now (a month after the occurrence) he is hardly conscious of any weakness remaining. There is very little difference, however, in the pulse; the slightest possible thrill being all that can be perceived.

With respect to the origin of the symptoms affecting the left arm, the evidence of the patient leaves no doubt that the paralysis was the result of the accident; but, the abnormal condition of pulse may of course possibly have existed previously without his knowledge. Looking, however, to the very curious and bizarre effects of lightning on the human body, the serious internal lesions which it may produce without leaving any trace of its action on the surface, I am of opinion that both the stoppage of the pulse and the paralysis were caused by the stroke. We know the tendency of electricity operating within a blood-vessel to cause coagulation; and it is not difficult to conceive that a current of extreme intensity, conducted as this was, from link to link of a silver chain of insufficient bulk for its free transmission, and which lay across the subclavian artery and brachial plexus of nerves, may have produced plugging of the vessel by coagulation, and have affected the nerves so as to cause loss of muscular power and numbness of sensation in the limb.

Time will not permit me to enter upon the question of the very curious and varied forms of injury which may be caused by lightning; but I would refer those who wish to inquire further into the subject to a very elaborate and exhaustive treatise, in two large volumes, by Dr. F. Sestier, published in Paris in 1866 (*De la Foudre, de ses Formes et de ses Effets sur l'Homme, les Animaux, les Végétaux et les Corps Bruts: Des Moyens de s'en préserver, et des Paratonnerres*). You will there find statistics and records of cases in abundance, showing at length and systematically the effects of lightning on each organ of the body; showing that persons may be struck dead without any trace of injury being discoverable either internally or externally; that there may be fatal internal injuries without lesion of the surface; that the surface may be extensively and deeply burnt without injury to the clothes; and that every article of clothing may be stripped off and scattered in all directions, leaving the wearer naked and unhurt.

There is, however, one practical consideration indirectly growing out of these cases, and in the nature of what may be styled *preventive surgery*, which is not without interest in this season of frequent storms, and on which I will say a few words, in order that you may be able to give some sort of answer to the question not infrequently asked by timid persons, as to the safest course to pursue when overtaken by a storm of thunder and lightning.

There is no doubt that the safest place for shelter is in the interior of a dwelling-house or other enclosed building, at a distance from windows and street-doors; and in a cellar, perhaps, for choice. Not only is the chance of being struck infinitely less, but the risk of serious injury is also much diminished. According to Dr. Sestier, the proportion of *fatal* cases occurring within houses is not more than one-fifth of the persons struck, and of these the greater number were standing near a window or on a door-step; whereas in exposed situations in the open country, three-fourths of the cases are fatal.

There is no doubt, also, that the popular notion, that it is imprudent to take shelter under a tree, is well founded, especially if, as in the cases before us, the tree be an isolated one. A low tree, or a hedge with several high trees in its proximity, is less objectionable, as the lightning will generally be attracted by preference to the most prominent objects. Trees standing together in a wood are seldom struck; the electric cloud, coming within the attraction of a mass of trees, pro-

bably discharges itself insensibly through the innumerable points of foliage. A wood, therefore, is not an unsafe place, though even there it may be well to keep away from a tree which is higher than its neighbours. Many persons have been killed while standing under a hay or corn rick; these, therefore, should be avoided. From their dryness they are worse conductors than the human body, so that the current passes from them to the latter, as the readiest channel by which it can reach the ground.

But if it is unsafe to stand under a tree or a haystack, is it safer to remain in the middle of a large open space? This is a doubtful question; for a man in the erect position, though less prominent than a tree, still offers a dangerous point of attraction when no other object is near, and if struck, the whole force of the stroke will pass through his body, entering probably by his head; whereas, under the tree, the current is likely to be divided and split up, so that though the chance of being struck may perhaps be greater, the risk of fatal injury is considerably less. According to M. Sestier's statistics, of those struck in exposed situations in the open country, three-fourths are killed; under a stack of hay or corn, two-thirds; under a tree, one-half; under a hedge, one-third; within houses, only one-fifth.

It appears to be pretty generally agreed that the safest plan, supposing shelter within a house to be unattainable, is to remain in the vicinity of some prominent object, such as a tree, but on the side opposite to that from which the storm is proceeding, and at a distance sufficient (say twenty or thirty yards) to avoid the risk of the electricity being attracted from the tree to the person. In the example before us, the horse and cart, which were standing near, but not under, the tree, escaped without injury.

Under any circumstances the recumbent is undoubtedly safer than the erect position; elevated and prominent situations being of course carefully avoided. It would be better, for instance, to lie down in a furrow than on a ridge. Additional security may also be obtained by depositing watch and chain, money, or other metallic substances which attract electricity, at a safe distance. Probably you will say that, by pursuing the course suggested, you are certain to get wet through, while under a tree you only run a remote chance of being struck by lightning. That is true, and the alternative selected will, I fancy, depend pretty much on individual temperament. Wet clothes, however, are not without a compensating advantage: they are all the better conductors of electricity; and, if they do not convey safely to the ground the whole of the current, they will transmit a much larger proportion of it, so that there will be all the less risk of serious personal injury. The men of whom I have been speaking went under the tree with their clothes only slightly sprinkled by the rain.

A gun is a very unsafe companion: nothing could well be worse than to walk about in a thunderstorm with a gun over your shoulder. To do so, is to convert yourself into the upper or attractive part of a lightning-conductor, without the accompanying provision for conveying the current safely away into the earth. Neither is it wise to walk along an exposed road under an umbrella, especially one with metallic stem and framework.

CLINICAL MEMORANDA.

HYSTERIA.

WHEN summoned to a female patient in hysterical convulsions, we are apt to ignore danger, and to speak of the attack as "only hysteria." The cases related by Dr. Foss in the last number of the JOURNAL prove that death may happen in one of the paroxysms; and I have lately met with a similar instance.

A married lady, of middle age and hysterical temperament, in whom faintness was readily induced by nervous excitement, was suddenly called to the bedside of her mother, who was supposed to be dying. The moment she saw her, a paroxysm of hysterical convulsion came on, followed by failure of pulse, cold perspiration, and unconsciousness. Although she partially recovered from this state of syncope, under the care of my friends Mr. Fryer and Mr. Hallett, death took place within twelve hours. A *post mortem* investigation by Mr. Hallett revealed a degenerated and softened condition of the heart. She did not come under my own notice until some hours had elapsed; and then I found her so much better that I unfortunately gave a very favourable prognosis, and was astounded when, next morning, I received a message announcing her death. As an old patient of my own, I had frequently seen her in similar attacks of more or less severity, and was not at all prepared for a fatal result on the present occasion.

Bristol, July 11th.

W. F. MORGAN.

ABSTRACT OF LECTURES ON UTERINE HÆMORRHAGE.

Delivered at the Hospital for Women, Soho Square.

BY ARTHUR EDIS, M.D.,

Physician-Accoucheur to the British Lying-in Hospital, etc.

LECTURE I.

GENTLEMEN,—The subject for our consideration this evening is one that cannot fail to interest every one engaged either in the busy routine of general practice or in the study of diseases incidental to women. I intend using the term uterine hæmorrhage in its strictly correct, but, at the same time, not in its ordinarily accepted signification; this being applied almost exclusively to floodings connected with pregnancy and parturition. The term *menorrhagia*—literally, excessive menstruation—is far too limited to include all that I wish to say on the subject; and the term *metrorrhagia* is only applied to a steady flow of blood from the uterus, whether profuse or not, during the intervals of menstruation. Hence I employ the more comprehensive term uterine hæmorrhage, and shall limit my remarks principally to cases occurring in the unimpregnated uterus.

GENERAL CAUSES OF UTERINE HÆMORRHAGE.

I enumerate here, for your guidance, the most frequent general causes of this affection. They are as follows: Mental worry, distress, shock, etc.; emotional causes; ovarian irritation; heart-disease; tubercle, bronchitis, and pneumonia; emphysema; residence in tropical climates; congestion of liver; general plethora and luxurious living; sedentary and unhealthy occupations; abdominal tumours; constipation; hæmorrhoids; residence in marshy districts—ague; albuminuria; hæmophilia; exanthemata, cholera, fevers, etc.; anæmia; scurvy; lead-poisoning; period of sexual involution at change of life. A mere glance will suggest to you that, though it may be prudent to devote time and attention to a *speciality*, it is unwise to become a pure *specialist*; for uterine hæmorrhage cannot be regarded otherwise than as a symptom of the most varied affections, both general and local; and the sooner you get out of the way of regarding it as an entity or a disease *per se*, the sooner will you be in a position to treat rationally and successfully the cases with which you will meet, and will not lose time and credit in giving the numerous vaunted specifics for menorrhagia.

The consequences of uterine hæmorrhage, if allowed to go on unchecked, are sterility, hydræmia, hysteria, dyspepsia, extreme emaciation, death. The woman who repeatedly suffers from severe menorrhagia, without adopting means to restrain it, will become prematurely old; will, at a comparatively early period, present evidences of approaching decrepitude and of that re-arrangement of nutrition which ushers in old age. Who amongst you has not noticed in society individuals whose blanched appearance, dark areolæ surrounding the eye, excessive languor and extreme irritability, whose very existence seems a burden too great for them to bear, and to whom every exertion is an effort? All this is due to some form of uterine hæmorrhage, which, either from neglect on the part of the patient, or inexperience on that of the practitioner, leads to much useless suffering and discomfort.

Before, however, we are in a position to appreciate morbid alterations in the discharge of blood from the uterus, we must first, very briefly, consider the normal condition of menstruation, or that periodic discharge of a sanguineous fluid from the uterus, which is attended by an engorged or congested state of that organ, as well as the ovaries and adjacent tissues. A greater part of the fluid under ordinary circumstances, probably the whole, comes from the uterus itself, and appears to be poured out from the open mouth of the uterine glands, which run from the free to the attached surface of the mucous membrane in a tortuous manner, this latter presenting a deeply red, velvety, soft structure. The cavity of the cervix does not ordinarily pour out blood. The tissue of the uterus itself contains large sinuses, receptacles for venous blood, which during menstruation become filled; and the veins being unprovided with valves, congestion of the uterine plexuses readily occurs, and the uterine tissue becomes engorged and expanded. It would appear that this condition, during the sexual life at least, is liable to occur also during intercourse. It is important to remember that ovulation and menstruation bear a peculiar relation to each other, the latter process not occurring in the absence of the former, although the contrary does not hold true. Ovulation, if it do not actually produce menstruation, is first in the chain of events; the ovary is, as it were,