probably only one stone being present, there may, indeed there will, soon be many stones, for each fragment becomes a nucleus for fresh deposit, and this hastens on with greatly increased rapidity. Even the nucleus, always a centre of interest, may be as appreciable in the fragments from lithotripsy as when displayed by the saw. It has happened to me in a case of crushing, in a female, to be struck with the appearance of redness in many of the fragments; and on investigation the mystery has been revealed on confession, that the patient had been in the habit of tickling herself with a stick of red sealing-wax, a portion of which had broken off in the urethra and remained in the bladder. The fragments will be seen in the collection.

Again, I once was aware, in using a lithotrite in a male, that I had clutched something peculiar. On withdrawing the instrument, there was a black substance about an inch long between the blades. A surgeon present, who had been in charge of the case for years, immediately exclaimed, "Egad! this is the end of my gutta percha catheter." A terrible revelation, for in the *interim* the patient had undergone prolonged treatment for chronic inflammation of the bladder, and had actually gone a voyage to Madeira in search of health.

I put as much faith as any man does in the chemical treatment, if I may so call it, of the diathesis of stone; but when once a stone has formed (and in most instances it is so without marked premonitory warning), the "fact" of stone is established, and there may be room for doubt whether chemical treatment does not then make matters worse; for, whatever the quality of urine, the chances are that a nucleus being present, deposition of stone will go on with increased rapidity, equivalent to the increase of size of the stone. That there may be exceptions to this rule, I admit; and there are two stones in my collection from one bladder, which are so smoothly polished by attrition that the formation of more stone had probably ceased for many months, if not years, before they were removed.

I have referred, gentlemen, to the chemical treatment of stone in the bladder, chiefly for the purpose of ventilating a sort of heresy of my own—viz., that in our treatment of stone, and in our estimate of specimens of stone in our museums, the chemical composition has been improperly the feature most referred to as the one of the greatest importance. Stone in the bladder is essentially a surgical disease; it can be treated effectually only by the surgeon; and to him the size, or, I may call it, the circumference of the substance to be removed, possesses the most engrossing interest, whether he looks to his own mechanical work or the safety of his patient; for I hold it as a maxim, particularly in lithotomy, that the bigger the passage required for egress, the more difficult and the more dangerous is the operation. The *accoucheur* considers the size of the head, but does not trouble about its chemical qualities or composition. So should the surgeon the stone, both in regard to lithotomy and lithotripsy.

But time will not permit me to say more on such subjects; and I shall conclude by a partial quotation from one of my lectures, delivered seven years ago at the College of Surgeons;—some of my language holding as strong an impression on my mind as ever. In the collection which I have the honour to exhibit, "there are examples of comminution from lithotripsy, enough to satisfy those not much acquainted with the operation, and entire stones, the result of lithotomy; exhibiting specimens from a few grains in weight to nine ounces—from one solitary stone in the bladder to forty-two. There is the first, taken out in 1832; and the last, removed in 1872. There is a single, solitary, grim, savage-looking mulberry calculus which held possession of its vesical home for forty years; there are numbers, smooth and less formidable in aspect, yet equally painful and dangerous to the sufferers. A momentary glance can sweep over the range; but only those who, like myself, have been engaged in such work, can estimate the days, the nights, the months, the years, of anxious thought and labour, associated with the results which I now put before you."

THE HOSPITAL FOR SICK CHILDREN.—The Princess of Wales, since the laying of the foundation-stone of the new buildings of the Hospital for Sick Children, has sent the patients a quantity of fruit and toys.

POST PARTUM EXPLOITS.—At a recent meeting of the Obstetrical Society of Edinburgh, Dr. Keiller observed that he had recently witnessed several instances of extraordinary exploits on the part of women immediately after their confinement without subsequent suffering. As showing what some women can do, without any bad consequences, he stated that the other day a woman was delivered in the Old Assembly Close, no one being with her at the time. After the placenta came away, she put her room in order, and then, placing the child and the placenta in a basket, she walked coolly to the Maternity Hospital, within an hour of her confinement, to see if everything was right with her. Dr. Keiller exhibited the fetus, which was apparently about the seventh month.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF MIDWIFERY.

At the Annual Meeting of the British Medical Association,
in Birmingham, August 1872.

By EVORY KENNEDY, M.D.,

Late Master of the Dublin Lying-in Hospital; President of the Section.

As I understand, it is the duty of your Presidents to open each Section of the British Medical Association with something of an address. It occurs to me that, in place of occupying the valuable time of so many of my *confrères* with an historical analysis of the progress of our branch of medicine—the usual stock subject of all introductory lectures, and one on which you are all very likely more competent to descant than myself—I might possibly occupy your time more profitably and pleasantly by giving you a few cases of not very ordinary occurrence which I have selected from my notebooks. Be not alarmed. I shall not weary you with detail in these brief *excerpts*, but merely touch the salient points of interest in each case.

CASE I. Fatal Femur embedded in Uterus.—A lady was brought to me from the country, anæmic, wasted, and with a countenance expressive of pain and suffering. She had a constant foetid vaginal discharge, purulent, and occasionally sanious. She was said to have miscarried a year before. She complained of pelvic distress, with lumbar pains and frequent micturition. On vaginal examination, the uterus was found about double its natural size. The os was slightly patulous, and a solid hard resistance was experienced on introducing the finger. The os was gradually dilated with a two-bladed uterine dilator constructed for the purpose, and a bony substance was distinctly felt, traversing the neck of the uterus at the junction with the body of the organ. This was so deeply embedded in the walls of the uterus, that neither extremity could be dislodged. I introduced a pair of small beaked bone-forceps, constructed for the purpose, broke the bone across, and extracted piecemeal the femur of a fetus of about four months' development, which had been all the time embedded in the walls of the uterus, which was converted into an abscess by the presence of what should have been its natural occupant, thus forming an unnatural nidus in its walls in place of its cavity. The discharge subsided; and the lady recovered her usual health; but the look of health was never quite restored. She, however, bore children after an interval of some years, and the uterus appeared to have quite regained its normal condition.

CASE II. Excision of Part of Neck of Uterus.—Dr. Kitson of — brought a patient from the country, suffering from ulceration of the os uteri. The neck was enlarged considerably, and elongated; the ulcer, which impressed us both as presenting all the characters of malignancy, occupied about one-third of the neck. It had taken a rapid course, bled at intervals freely, and upon the slightest touch, and was attended with pain, sleeplessness, and marked constitutional disturbance. It was, however, circumscribed and limited to the part ulcerated; the remainder of the neck and os being healthy to the appearance and touch, although larger than natural. The lady had borne children. The part of the neck engaged extended from the posterior along the left side of the os, and the diseased structure appeared to occupy the entire substance of the wall. Under these circumstances, the case promised little or nothing from the application of the ordinary caustics, and the choice appeared to lie between the free application of potassa fusa and excision. The latter was determined on; first, because of the limited extent of the part engaged; secondly, because of the apparent malignancy; thirdly, from the difficulty of destroying by the potass the whole diseased structure, without extending its action to the adjoining vital parts. On the other hand, the diseased structure came well within our view; the neck was long, affording facilities for the use of the knife. The patient was placed on her back. The vaginal wall and labia were distended by my four brass tractors, firmly held by Dr. Hans Irvine and Dr. Kitson. An ebony spatula, nine inches long and half an inch broad, was introduced and placed within the os. This I held firmly in my left hand, whilst I introduced the scalpel which I now exhibit, which, you perceive, has a handle seven inches long, while the blade is scimitar-shaped. Cutting from without inwards towards the resisting spatula, commencing near the point of junction with the neck and body of the uterus, above the central part of the diseased structure, by two divaricating incisions *A*, a triangular section was removed. I was prepared to draw the uterus down with the double tenaculum; but this was unnecessary, from the perfect manner in which my assistants used their tractors. This allowed me the

assistance and security of the spatula to cut upon. It has occurred to me that, in a case where excision is preferred, and where the facilities I describe do not exist, the spatulum might be armed on the reverse side with two hooks, when it would perform the double office of uterine tractor and spatula, as necessary. The vagina was simply plugged with Ruspini's styptic. There was scarcely any hæmorrhage. The patient recovered speedily and perfectly, and in about two years afterwards conceived and carried a living child to the full period. Her labour was easy and natural; and I had an opportunity of examining her at an interval of several years afterwards, when she was quite well, and the uterus, with the exception of the loss of a portion of the neck, was perfectly sound.

CASE III. Portion of Placenta thrown off in Pregnancy.—A lady, in the seventh month of her fifth pregnancy, was seized with hæmorrhage, ascribed to over-exertion. There were no labour-pains. On examination, a portion of the placenta was found protruding through the os uteri. The hæmorrhage continued for several days, but to no serious extent; and still there was no labour. At length, foetid grumous discharges, mixed with a little blood, occurred, attended with sense of downward pressure. The portion of placenta descended lower in the vagina; its connection with the interior of the os separated; and I removed it with very little assistance. As no increase of hæmorrhage occurred from this, I thought it unnecessary to plug the vagina. The hæmorrhage and discharge ceased, and the patient went on without any inconvenience, except the precaution of keeping the horizontal position for six weeks longer, when she was delivered of a living boy apparently at or near the full time. The edge of the placenta that remained could not be felt near the os, and the portion that came away consisted of the vascular structure without the reflected membranes. There was no discharge of liquor amnii until the labour set in.

I have already had the honour of calling your attention to some of the more rapidly destructive of puerperal diseases in a paper read for me, in my absence, by your Secretary, at your Dublin meeting, under the head of purpuric puerperal fever. It is now my intention to allude briefly to other forms of blood-poisoning, but more especially to puerperal arthritis and puerperal gangrene; premising that, when this disease shows itself, it is usually most rapid and unsparing in its onslaught, and no tissue in the body escapes its ravages.

CASE IV. Destructive Inflammation of the Eye, combined with Arthritis.—Bennet was delivered of her first child after a labour of nine hours, on June 14th, 1838. On the 16th and 17th, there was abdominal tenderness, with a quick small pulse. She was leeches. On the 18th, her countenance was anxious and shrunken; pulse 140, small. On the 19th, sudden inflammation of the right eye set in; the conjunctiva and iris being engaged. Leeches were applied and blisters to the temples. Mercury was given. On the 21st, she was salivated. The whole globe of the eye appeared prominent with serous infiltration beneath the conjunctiva. All the tissues of the eye were engaged; the sclerotic was very vascular, the cornea prominent, the aqueous humour turbid, the iris hardly discernible; there was total loss of vision. The conjunctival vessels were freely scarified. This evening, inflammation of the wrist set in, with erythema of the back of the hand. Excitement and delirium were present. On the 22nd, the other wrist became engaged; her countenance was shrunken and excited. Pulse 125, small and thready. The wrists were treated with solid nitrate of silver. She sank, and died on the 24th. At the *post mortem* examination, effusion of blood was found on the right side and posterior lobe of the cerebrum, and a small quantity between the arachnoid membrane and the pia mater on the left side. The brain in other respects was healthy; there was no fluid in the ventricles. The right eye was much suffused; the conjunctiva very vascular, the cornea perfectly opaque. On extracting the eye, some pus was found round the lacrymal sac. On cutting down on the carpal bones, a quantity of pus was found in the sheaths of the tendons, unconnected with the joints. Pus was found in the vessels of the broad ligaments of the uterus; but there was no fluid in the cavity of the abdomen.

I would now call your especial attention to the drawing which I exhibit to you as a type of disease known as sloughing of the vagina, but which would be more correctly designated as lying-in hospital gangrene; and also to a group of cases occurring in puerperal women as the result of their state and condition, but which are now clearly established as prevailing so commonly in great lying-in hospitals, and so rarely out of them, as to stand in relation of effect and causation with the locality of their delivery. As in surgical hospitals, so in lying-in hospitals, these cases are ascribed to hospital or poisoned atmosphere.

These cases generally occurred in groups. For instance, in April 1837, four occurred in the Dublin Lying-in Hospital at the same time, not one of which recovered. One occurred in June of the same year, complicated with arthritis of the elbow-joint; one in October, and one

in November—all being fatal. Another at the same time was complicated with erysipelatous inflammation of the buttock, followed by abscess and inflammation of the elbow, sterno-clavicular and shoulder-joints; and yet this woman (Keating), the drawing of whose disease I exhibit to you, struggled through all this and recovered, after two months of as acute suffering, from the number of joints consecutively engaged, as ever I witnessed. Free incisions were made into the buttock, with escape of pus. The sterno-clavicular joint suppurated, and an incision was made; but the inflammation of the elbow- and shoulder-joints yielded to the free application of nitrate of silver and potassa fusa. On the 8th of December, another fatal case occurred. This case was complicated with slight peritonitis; and the *post mortem* examination disclosed a peculiar lesion of the peritoneal surface of the uterus at the junction of the neck and body anteriorly, which I have rarely observed, but which of itself I esteem a very fatal occurrence.

CASE V. Puerperal Arthritis: Erosion of Cartilages of Elbow, Hip, and Ankle Joints.—Kenny, three weeks delivered after a difficult and protracted labour, was awakened from sleep in the night by an acute pain in the left groin. In the morning, she observed a swelling in the middle of the thigh, which at the end of two days had completely engaged the entire limb. The pain became less acute as the swelling increased, but never entirely subsided. Some days subsequently to the swelling of the thigh, she was seized with violent pain in the elbow, but did not perceive any swelling. All these symptoms progressively increased, notwithstanding frequent leeching, stuping, poulticing, opiates, and mercury. She was admitted into hospital on January 28th, 1829; and, on the 30th, there was an obscure sense of fluctuation over the outer third of the thigh. An incision was made into it, but no pus followed. On February 3rd, she had a severe rigor; and on the 4th she died comatose. A *post mortem* examination was made twelve hours afterwards. The cellular tissue throughout the entire thigh was filled with gelatinous lymph. An extensive abscess extended from nearly one extremity of the thigh to the other, between the periosteum and muscles. The muscles were pale and flabby, and appeared much softer than natural. About one inch of the upper part of the femoral vein contained pus; its inner tissue was vascular, but did not appear to have lymph upon its surface. The synovial membranes of the hip, knee, and ankle joints were filled with puriform matter. The cartilage covering the bones of the hip appeared healthy; whilst that covering those of the knee and ankle was in part removed by absorption, particularly in the ankle, where scarcely a trace of cartilage could be detected. The uterus was vascular, and inclined towards the left side. The lymphatic glands along the iliac vein were enlarged and vascular. The cartilage was removed altogether from the extremities of the bones forming the right elbow-joint. The viscera appeared healthy.

CASE VI.—Puerperal Arthritis: Sterno-clavicular Articulation and Buttock engaged: Recovery.—Keating was delivered in November 1837 of a still-born child, after a labour of thirty-five hours. There was some sloughing of the vagina, but not considerable, with a good deal of tympanitis of the abdomen and irritation, which were relieved by turpentine followed by opium. On the eleventh day, there was an erysipelatous state of the right elbow, which had been, she stated, painful three or four days previously. She was treated with nitrate of silver locally, and sulphate of quinine with hyoscyamus constitutionally. On the twelfth day, she was attacked with erysipelatous inflammation of the left buttock (the elbow having become improved), which was also treated with the caustic and with quinine. On the sixteenth day, there was an appearance of pointing in the buttock, and an eschar was made with potassa fusa. There was this day noticed some tenderness in the right iliac region. On the seventeenth day, she complained of the left sterno-clavicular articulation, which was swollen, red, and painful; it was scored with caustic, the other treatment was continued, and wine was given. On the twentieth day, a fresh eschar was made on the buttock, through the slough consequent on which a lancet was passed on the twenty-ninth day, giving exit to matter of a healthy character. The swelling of the sterno-clavicular articulation continuing, with decided fluctuation and uterine pain, there being present also great constitutional disturbance, an opening was made on January 11th, 1838, and exit given to healthy pus. She gradually improved after this, and left the hospital. There still remained some swelling, but the opening had healed.

I can conceive what I term lying-in hospital gangrene to occur out of hospital; yet, after a protracted and sufficiently extensive opportunity in my own practice and in consultation, I cannot tax my memory with ever having met with a case of it out of hospital. Would that I could make the same statement of my experience in hospital! It occurs in isolated cases, but more frequently prevails in groups of cases, when a tendency to its congeners—puerperal fever, erysipelas, arthritis, and

peritonitis—also prevails. It is usually the effect of long continued pressure or lesion, or of the force used by the application of instruments in forced deliveries. Indeed, so constantly is the disease observed to occur as the result of these agencies, that they may be fairly looked upon as the exciting or determining cause, and the disease classed as a traumatic affection, and treated accordingly. This is so well understood by the observant hospital physician, that he looks with dread to the occurrence of any lesion when the hospital is in an unhealthy state; and, for the same reason, hesitates to perform the simplest operations, lest they be followed by this affection—operations that, in a healthy state of the hospital, would not cost him a thought.

I hold that the congregation of a number of patients in common chambers generates what we term a hospital atmosphere; that this hospital atmosphere is, or more properly becomes, a poison; that, by a persistence in the causes of its production (with which we shall presently deal), it undergoes a process of what may be termed cumulation, pervading every part of an enclosed building, until it eventually arrives at a stage which we may term saturation, when the whole hospital is charged with a poison capable of seizing upon those who are susceptible to its influence, or who are in what we term a state of receptivity.

Now, from this it will be seen that, by crowding patients in a hospital, we are actually exposing them to a new disease generated by the very means we adopt to cure them of the disease under which they chance to labour. But, unfortunately, the new disease is generally one most fatal in its character, as few there be who survive it.

The characters or phases of the hospital disease vary under the different circumstances of the victims susceptible to it. Thus one may be attacked with blood-poisoning or empyema; another with erysipelas; a third with hospital gangrene; and a fourth with metria or puerperal fever. The laws which regulate the habits of this family of zymotic disease are perhaps best arrived at by a study of the last named poison, as, in its occurrence in our great lying-in hospitals, where it principally commits its ravages, it is less exposed to disturbing influences, and consequently pursues its own natural course free from interruptions and complications.

Out of a hundred and eleven years, for which the great Dublin Lying-in Hospital has been established, it has been haunted by puerperal fever ninety-three years. For twelve years it has been comparatively, and only for eight years has it been totally, free from this fell disease. The deaths of those admitted for the last years amount to 1 in 33. Let us remember that in three small cottage hospitals in Ireland, in which accurate tables have been kept (Kilkenny, Newry, and Waterford), we find that the mortality has been 1 in 282. On the comparison of these two proportions, the conclusion is inevitable that eight out of nine patients have died in the Dublin Lying-in Hospital who would not, in all likelihood, if they had taken refuge in the cottage hospitals, which were comparatively free from the hospital miasm or poison that prevailed or lurked in the great unhealthy hospital.

A fatal error, into which we are prone to fall, is the confounding epidemic and endemic disease; and the amount of loss of life that has occurred from this error I believe to be incalculable. This will be easily appreciated when we state that what, in strictness, we should call true epidemic diseases, are unavoidable; whilst the latter, or endemic diseases, are, with rare exceptions, preventable and perfectly within our control. True, they may be convertible; but this makes the distinction the more important, in order to prevent their extension or fixture.

The contagious nature of hospital miasm is now beyond question; as also that of most of those modifications of hospital zymotocene with which we are familiar, especially metria and erysipelas.

Having satisfied ourselves that a poison is generated by the mere crowding of numbers of patients into a common atmosphere, and also that this poison spreads by contagion, the next principle or law we require to establish is that this poison is cumulative, or developing in its quantity, commencing with a single poisonous emanation, and increasing in its quantity until the atmosphere, walls, floors, and furniture become imbued or charged with it. Unfortunately, as yet we are unable to detect this miasm or poison and display it by its sentient properties; but of its existence we cannot have the slightest doubt, from its effects, from its laws, from analogy, and especially the spread of diseases by inoculation and contact.

But why should the hospital wards ever be free from hospital miasm and these fatal results, if all this be true? This is a question to be answered by the law upon which we now dwell—cumulation. The ordinary epidemic disappears; having, as we say, worn itself out. This it does in hospital as out of it. Precautions are taken; patients zymotically affected are separated; ablution and ventilation are carried out; admissions are refused; the wards are emptied; and the hospital becomes healthy. Weeks, months, and longer, pass over before the poison

again shews itself. The cumulative process, however, is in steady operation; and at an uncertain period the poison again shews itself, and snatches up its victims. The same measures of precaution are taken to banish it, and with the same results. But it does not rest here, as has been abundantly proved by the history of our great hospitals. When the cumulation has gone on repeating itself again and again, a further stage, or that which I shall denominate the state of complete saturation is arrived at, and then the hospital becomes the fixed *habitat* of the poison.

The death of the consumptive costermonger is not to be placed in the category of the victims to hospitalism. It has its analogue in the dog slain in the Grotto del Cane for the instruction of his slayers, by holding him so close to the surface of the deadly cavern that he can only inhale mephitic gas. Nor, indeed, is it only in St. Pancras that we have this going on. Have we not our No. 11 wards and grottoes in nightly operation for the benefit of the upper ten thousand in the West-end *réunions*? Witness the pallid and poisoned state to which our belles are reduced, at the end of the season, from breathing mephitic and animal-poisoned air for several hours nightly, with their lungs at nearly as great a disadvantage as the costermonger's, by tight lacing and waltzing. The Black Hole of Calcutta and the middle passage rendered our fathers familiar with the more immediate effects produced from a number of people breathing and rebreathing the same atmosphere; and its more gradual effect in the production of jail-fever, grappled with by Howard in his mission of mercy, has reached us as a matter of history. We were in hopes that the labours of Boswell Reid and Arnott had exposed the defects of ventilation and directed a remedy. But the enormities occasionally cropping up from neglect in this respect shews that much remains to be done. Dr. Farr, in a paper lately read at the Leeds Conference, pointed out the want that exists of an authoritative organisation charged with the ministry of public health. As it is, what is every man's business is left undone. When will our legislators waken up from their piecemeal attempts at sanitary improvement, and grapple with this great subject as it demands, and as the well-being of the community requires?

The fact is that, when the poison is generated, crowding or the cohabitation of patients in a state of receptivity secures its spread. A process of cumulation of the contagium or poisonous element occurs; and when this arrives at the stage of saturation, outbursts of erysipelas, phagedæna, gangrene, pyæmia, and metria occur; the patients are attacked wholesale; and the diseases become, as we before stated, endemic or fixed, and continue to haunt the hospitals, be they surgical or obstetrical.

Now that the blot has been hit, it requires no great philosophy to meet the difficulty. Simply cease to crowd such patients into common buildings under the same roof. All the advantages of an asylum can be afforded whilst segregation is secured, and facilities for medical instruction preserved to the schools. The substitution of cottage-hospitals for these great hotbeds of contagion should be insisted upon in all future arrangements for housing cases liable to endemic poisons. The term epidemic should be applied in its true meaning. These large surgical and maternity hospitals, in which metria, erysipelas, and pyæmia have committed such havoc, should be either applied to purposes compatible with the safety to human life, or so altered in their construction as to isolate each ward from the common atmosphere now pervading the whole hospital. This may be very simply and inexpensively accomplished by opening separate entrances into each ward direct from the open air. In the existing hospitals, the communications with the common hall should be built up, and the same system carried out in the upper storeys by opening the ward-doors on flying galleries. The lifts and stairs should be placed outside of the building; the existing halls and passages being still retained for the staff, and affording an approach to the respective storeys, but no covered approach or communication of any kind allowed to remain between these and the wards. By these simple expedients, and the limiting the number of beds to three in each ward, it is probable a large saving of human life would result, and hospitalism become, like jail-fever and small-pox, a thing of the past. Pity the founders of the great hospitals so recently erected had not applied our dear-bought experience in their magnificent constructions. The sooner these altered arrangements are made in such great palatial institutions as St. Thomas's and the Leeds Hospitals, as well as the Maternities and our other great charities subject to erysipelas, pyæmia, and phagedæna, the sooner will the insidious approaches and the fell ravages of our common enemy, hospitalism, be subdued.

Stow, in his *Survey*, mentions, in speaking of the King's Bench Prison, that the great mortality that occurred there in the six years preceding 1579 was produced through a certain contagion called "The Sickness of the House." Those learned physicians who uphold the

crowding in large hospitals, and imagine they can meet the difficulty by disinfectants, I would refer to the simple experience of that benefactor of our species, John Howard, as conveyed in his preface a century since. He says: "I guarded myself by smelling to vinegar while I was in those places. This I did constantly and carefully when I began; but, by degrees, I became less attentive to these precautions, and have long since entirely omitted them." John Howard discovered and carried out the best means, on true scientific principles, of disposing, once and for ever, of "the sickness of the house." If the obstinate hospital crowdors would take example by him, and meet, as he did, crowding, the cause of "the diseases of the house," and, as he did, remove them, they would speedily produce an equally satisfactory result, and, like Howard, give up their futile attempt to meet the case of dealing with emergency by "smelling to vinegar."

CASE VII. *Lying-in Hospital: Gangrene.*—Catherine Barnes was delivered by crotchet, after a tedious labour, of her first child. She had one fit of convulsions on entering the ward, and was treated with tartar emetic. There was no return of convulsions. Extensive sloughing of the vagina showed itself the day after delivery, with great debility and exhaustion. Tympanitis and delirium occurred, and she gradually sank, and expired on the fourth day after her delivery. The appearances exhibited in the plate presented themselves. A small quantity of limpid serum was observed in the peritoneal cavity. The whole vagina and the lining of the uterus were in a state of slough. The os uteri was jagged and disorganised; and there was serous infiltration underneath the pelvic cellular membrane.

These cases are sometimes attended with hæmorrhage. For instance, in Murphy (October 1837) there was lingering labour with slow dilatation. She was treated with tartar emetic and ipecacuanha injections. She was delivered naturally on the 17th, but with laceration of the perineum, of a still-born male child. Sloughing was perceptible, with rigors and prostration. Hæmorrhage from the vagina set in as the sloughs were separating, on the 24th; and she sank on the 25th.

In Black's case, the drawing of which I here exhibit, she had tedious labour, followed by sloughing inflammation of the vagina and external parts, of erysipelatous character, spreading from the vulva to the buttocks and back. She sank from exhaustion after a long struggle. In addition to the appearances here represented, the omentum was found adherent to the pelvis and uterus; and an abscess in the walls of the latter organ opened into its cavity.

[To be continued.]

COLD FOOD FOR INFANTS.—Under this heading, Surgeon King, U.S.A., writes to the *Philadelphia Medical Times* as follows:—"Our best authorities direct that the [cow's] milk should be given to the child at the same temperature as that of the mother's milk—from 90 deg. to 95 deg. Fahrenheit—and, when great accuracy is required, a thermometer employed. On reflection, it is obvious that these instructions can never be carried out so that the little one will take all its food at the same temperature; for during a meal the bottle becomes cold, and there may frequently be considerable difference of temperature between the first and the last milk imbibed by the infant. It is unnecessary to state that very little will upset the feeble powers of the digestive organs in the early days of infantile life; and this difference in the temperature of the food, I am disposed to believe, is one of the causes of gastric and intestinal disorder which we so often have to deal with among infants brought up by hand. Instead of giving warm milk, I have adopted the plan of giving cold milk entirely—ordering the babe's bottle to be kept standing in iced water in the summer and in a cold place in winter. This method I have found, from practical experience, to answer remarkably well. If there is any tendency to diarrhoea, I recommend the milk to be heated to 212 deg. Fahr., and afterwards allowed to get quite cold before being used. In private practice, I am of opinion that bottle-fed infants generally have their food given them too warm. . . . They soon like it even better than warm food, and during the teething period cold milk seems especially agreeable to the inflamed gums of the little sufferers." The above plan for the preparation of food for infants is, the *Boston Medical Journal* thinks, worthy of trial. We all know the difficulty of having artificial food properly prepared, notwithstanding minute directions are given concerning it and the importance of keeping the nurse-bottle scrupulously clean. Many devices are resorted to by those who have the care of infants to avoid the trouble of freshly preparing the food every time it is wanted, and the temptation is great, particularly at night, to have in readiness a quantity sufficient for several meals. As a consequence come the various disturbances to the system resulting from the ingestion of food which is often sour before it is taken. If it can be given cold without detriment to the child, there seems to be no good reason why the diet should not always be fresh.

AN ADDRESS ON MEDICAL PSYCHOLOGY,

DELIVERED AT

The Opening of the Psychological Section at the Annual Meeting of the British Medical Association, in Birmingham, August 1872.

By HENRY MAUDSLEY, M.D., F.R.C.P.,
Professor of Medical Jurisprudence in University College; President
of the Section.

IN beginning the work of this Section, over which I have the honour to preside, I shall confine myself to a few introductory remarks of a general character, leaving to those who will come after me the more exact scientific work of which we have fair promise in the papers that are to be read. The occasion seems fitting to take a short survey of the position of medical psychology in relation to certain important questions of the day, and to consider the bearing which its progress must eventually have upon them. Permit me, then, to ask you, first, to look back a little way at what medical psychology was, in order the better to realise what it is, and, if possible, to forecast something of the character of its future work. A glance at the past will show how great a step forward has been made, and may yield some reason for congratulation; a glance at the present, showing, as it cannot fail to do, how small a proportion the gains bear to what remains to be acquired, will prove that as yet we have rather discovered the right path than made much way on it—that we are, in truth, only on the threshold of the history of medical psychology as a science.

One of the saddest chapters in human history is that which describes the cruel manner in which the insane were treated in times past. Notwithstanding that it is happily a thing of the past, it will not be without profit to inquire from what causes the barbarous usage sprang: for it was not common to all nations and all times; on the contrary, it had its birth in the ignorance and superstition of the dark ages of Christian Europe. Whatever may have been thought of madness among the peoples who preceded the ancient Greeks—and there is evidence that the Egyptians adopted a singularly enlightened and humane treatment—it is certain that the Greeks had comparatively sound theories of the nature of insanity as a disease to be cured by medical and moral means, and adopted principles of treatment in conformity with those theories. Their dramatic poets, it is true, present terrible pictures of madmen pursued by the anger of the gods; but these were poetical representations, which must not be taken as a measure of the best knowledge of the time. Then, as now, and indeed as ever in the history of mankind, the true thinkers were emancipated from the fables and superstitions of the vulgar: the just measure of Greek intellect must be sought in the psychology of Plato, in the science of Aristotle, and in the medical doctrines of Hippocrates. This eminent physician and philosopher expressly repudiates the notion that one disease is of more divine origin than another. After saying that the Scythians ascribe the cause of certain disorders to God, he goes on to give his opinion that these and all other disorders are neither more nor less of divine origin, and no one of them more divine or more human than another; that each has its own physical nature, and that none is produced without or apart from its nature. In what he says of the psychological symptoms of various diseases of the body, he evinces such enlarged views of the scope of medical observation and practice as are not often evinced at the present day; and the few observations in his works respecting the symptoms of delirium—"evidence that clear and correct view of disease which has made this first observer a model to all succeeding times". He directs attention to such facts of observation as the physical insensibility of the insane, the appearance of mental diseases in the spring, the occurrence of disorder of the intellect after a continuance of fear and grief, the union of melancholy and epilepsy, the critical importance of hæmorrhoidal discharges in mania, the difficulty of curing madness which commences after the age of forty, and the like. And as there was no superstition in these doctrines, so there was no barbarism in his treatment, which was medical, and consisted principally in evacuation by the use of hellebore. But moral treatment was not unknown among the Greeks; for Asclepiades, who seems to have been the real founder of a psychological mode of cure, made use of love, wine, music, employment, and special means to attract the attention and exercise the memory. He recommended that bodily restraint should be avoided as much as possible, and that none but the most dangerous should be confined by bonds. Without going further into particulars, enough has been said to show that the Greeks had acquired