CLINICAL REMARKS

ON A

CASE OF CONCUSSION FOLLOWED BY CHOREA AND DROWSINESS.

By C. HANDFIELD JONES, M.B. Cantab., F.R.S.,
Physician to St. Mary's Hospital.

J. S., aged 26, painter, was admitted December 9th, 1871. He stated that he had colic first six years ago, and again five weeks ago. He had no agitation then. He was always very excitable and quick tempered. In May 1869, he fell off a scaffold, and suffered concussion and injury of the spine, besides a large bruise on his thigh. He was in a hospital in London three weeks after this accident. On leaving, he walked, he said, three times round the hospital before he could make a start in the right direction for home, and then he was two and a half hours getting home to Park Lane, Upper Baker Street. He lay down on the bed, but the children distracted him, and he became worse. One day, while walking in Baker Street, he saw a child run over by a cab. A crowd collected, and he stood among them until he was desired by the police to move on. He found then that he could not move his legs, and had to get the aid of two persons to enable him to return home. The next day he applied to a public dispensary, where he was advised to go into a hospital, as he "could not walk without assistance; his head, arms, body, and legs, twisting and writhing like a toy made of India-rubber, causing his stomach great pain". The motive power seemed to come from his back, which very soon began to ache on exertion, and then he was glad to go home and lie down. When he was in bed his muscles would quiver; sometimes his arms, legs, and often his back, would spring up in bed almost upright; then it would gradually decrease, so that he could lie quiet. On getting up, his joints would feel very loose; his knees would give way under him, throwing him forward. He had never suffered in this way before. Some time after this he got a letter of admission to a convalescent hospital near Windsor (St. Andrew's). The shaking of the railway-carriage made him worse than ever; his difficulty of articulation was increased, and his unsteadiness. In fact, he could not put one foot before the other for a little time, and was unable, on his arrival at the station, to walk to the hospital. He was kept very quiet after he was admitted into the hospital, and otherwise judiciously treated, and gradually got better. In the course of a month every bad symptom was gone, and he was able to walk miles without the slightest inconvenience. He remained another fortnight, and when he came to work the only indication of his illness was a slight headache, an inclination to drowsiness, which he could shake off unless he had drank beer, and an inability to carry anything on his head. In the course of 1870, the sleepiness increased, and on ascending a ladder the back of his legs would quiver. At that time he began to be morose and melancholy; he would drop off to sleep as soon as he came home in the evening, sometimes before he had his tea, and never went out afterwards, and very rarely spoke to any one. In 1871, he had the gout in his right foot; his headache increased; and he would often go to sleep on a scaffold. He has known himself drop off to sleep on a standard ladder sixty feet from the ground. In October he began to feel very ill, had a spurious sort of hæmoptysis, soreness of his mouth and tongue, costiveness, inability to keep anything on his stomach, and aching of his joints like gout. He gradually became worse, and then came to St. Mary's.*

When I saw him, his condition was very much that of a choreic patient. His speech was much interfered with; it was not always easy to comprehend what he wanted to say. There was a good deal of jactitation. The registrar's note states that he could not keep a limb quiet. His grasping power was rather defective, but he could move both arms well. Sometimes the muscles of his arms and legs became rigid. He slept a good deal, and quietly. The heart's sounds were normal. His habits were temperate. He smoked little. The bowels were costive; face pallid. The agitation was induced by exertion. He could walk sometimes two hundred or three hundred yards before it came on, then his legs jumped and started about. He was ordered two drachms of succus conii in water three times a day; milk, beef-tea, and a warm bath.—December 11th. The bath produced an immediate effect in calming and steadying him. He was quite still now if allowed to be quiet, but if spoken to he began to twitch and yawn a good deal. Yesterday

there was a notable amount of bloody sputa; to-day he had very little. He said that he slept day and night. On the 13th, he was ordered eight grains of hypophosphite of soda three times a day, and two drachms of cod-liver oil once daily. After a week's stay in the hospital, he reported himself much better, was able to walk about briskly, and did not feel sleepy in the day-time. On the 18th, as he had still much pallor, I gave him five grains of hypophosphite of soda and eight grains of citrate of iron and quinine in water three times a day. On the 20th December, he had a good deal of choreic agitation or unsteadiness; he had much difficulty in making the house-surgeon understand that he wanted a certificate. On the 21st he was ordered a drachm of succus conii in water three times a day, and 20 grains of saccharated carbonate of iron three times a day. On December 28th he was better; the palpitation of heart, with which he used to wake up, had ceased. He was still rather unsteady, but walked about well and went into the open air. On January 2nd, 1872, I examined the eyes with the ophthalmoscope, and detected nothing more abnormal than some white streaks on the left disc, and along some of the retinal vessels. On the 4th, he went out. He was seen again on the 19th, when he stated that he had been going on very well up to that morning; but, his child having been run over, he brought it to the hospital, and while he was there the choreic agitation returned in a most marked manner. He was ordered to take two-drachm doses of the succus conii again.

My notes do not state whether there was a blue line on the gums in this case or not; but his occupation, and the fact of his having had two attacks of colic, make it certain that his system was under the toxic influence of lead. This is further substantiated by the occurrence of gout, which Dr. Garrod has shown to affect with increased frequency those who work with this metal. The question then arises whether the choreic disorder can be referred to this cause. Mercurial tremor is of course a well known affection; but few writers, as Dr. Sanders states, admit that lead can act in the same way. It appears that artisans using lead only suffer from the ordinary disorders, but the case is different with regard to lead-miners. "Brockmann, from his experience in the Harz mountains, has described a species of lead tremor which affects the miners there, and which is almost identical with the mercurial shaking palsy, consisting, like it, of oscillating spasmodic contractions of the muscles, and consequent tremulous motion in various parts of the body. The tremor may be local, usually affecting only the upper extremities, or general. The latter form never results except from deeply rooted lead-poisoning" (Reynolds's *System of Medicine*, vol. ii, p. 215). It is apt to be the precursor of paralysis or cerebral disease. In J. S., however, you will observe that the irregular movements were clonic spasms, and not tremors; there was no continuous muscular twitching, but frequent movements of considerable extent. The character, then, of the muscular disorder, the circumstance that the patient was not a lead-miner, and that the toxic action of the metal was not carried to any great degree, incline me to believe that his choreic agitation was dependent rather on the spinal injury than on lead-poisoning. This view is confirmed by his having never suffered from any thing like chorea until shortly after his accident, and by the fact that, even before the chorea appeared, his locomotor power was markedly impaired, as was evidenced by the difficulty he had in going home from the hospital.

The remarkable drowsiness from which J. S. suffered after his accident, as well as his moroseness and melancholy, afford proof that his encephalon—probably his hemispheres—had suffered damage, and, owing to impaired nutrition, performed their functions badly. As concussion often produces impairment of the intellectual powers, I can see no improbability in the view that it may have in this instance impaired motor power also, having injured the corpus striatum and other associated centres, and rendered their tissue unstable. The coexistence of symptoms of defective power and of increased excitability, frequently observed, was well marked in this case. It is most probable that the damage occasioned by the injury was not of a kind readily demonstrable; for, if such had been the case, it would be difficult to account for the great improvement produced by the residence at the convalescent asylum, and subsequently by the conium.

The cessation of another form of muscular disorder—viz., the palpitation, while taking the conium, deserves notice. The remedies for this neurosis are by no means so satisfactory, that we can regard with indifference the offer of another which may prove more effective. The effect of emotional excitement on our patient was remarkable, and tends materially to assimilate his malady to ordinary chorea. The action of conium in this and similar instances seems evidently not to be merely sedative, but, as Dr. J. Harley describes it, recreative and restorative. Not only did it tranquillise the irritable motor centres, but it rendered them more capable of exertion, and restored the hemispheres from their parietic and languid state. My recent experience of conium inclines me

* The foregoing statements are taken from the patient's own account, which he wrote out at my request.

to hope that we possess in it a remedy of great value in many states of nervous system characterised by the blending of hyperexcitability and debility. I have said nothing of the gout, which he seems to have had, as a *causa mali*, for it seems to me quite improbable that it had anything to do with the production of the motor disorder or the drowsiness: they did not alternate with the articular affection.

Shortly, we may regard the case as one where the nervous centres were originally of rather infirm, unstable constitution, and subsequently had their nutrition impaired by injury received in a fall, so that they performed their functions feebly and unsteadily. The lead-poisoning may at the same time have contributed something to the deterioration of the nerve-force by acting injuriously on the nerve-cells, but it was by no means the principal cause of disorder.

It may be well to consider how this poor fellow would have fared if his concussion had been received in a railway collision, and he had sought compensation in a court of law. Would not suspicions have been expressed rather freely that he was malingering, or at any rate exaggerating considerably? and might he not have had difficulty in convincing the jury that his malady was genuine? It is quite right that all claims for compensation should be closely scrutinised; but mistrust may be carried too far, and I call your attention to this case as a proof that concussion may produce symptoms which, though very real, might be regarded with much suspicion by those who have chiefly given their attention to cases of severe injury.

REMARKS ON A CASE OF HÆMATURIA FROM THE PRESENCE OF THE BILHARZIA HÆMATOBIA.

By HENRY SIMPSON, M.D.,

Physician to the Manchester Royal Infirmary.

THE publication of Dr. Cobbold's interesting paper in the JOURNAL of July 27th reminds me of a patient, suffering from the presence of the Bilharzia hæmatobia, who was under my care last year in the Manchester Royal Infirmary. He was seen by many of my medical friends, and I read a paper before the Medical Society of this city, on the History of the Bilharzia and the Phenomena due to its presence as a Parasite. This I had intended to send to the JOURNAL; but, as Dr. Cobbold's paper goes over much of the ground I traversed, and in a manner I cannot attempt to rival, I shall not do so. As, however, but few of these cases have been reported, and they still possess some of the interest of novelty, and of difficulty as regards efficient treatment, I may perhaps be allowed, as briefly as possible, to touch on some of the points of interest in the case.

The patient, William Ray, aged 20, presented himself as an outpatient about the end of January 1871, suffering from hæmaturia. Mr. Roberts, the physicians' assistant, learning that he had been in Egypt, and recollecting that this disease was endemic there, and connected with the presence of the Bilharzia, brought him to me. I at once examined the urine microscopically, and discovered the ova of the worm. The patient was slight but well built, as befitted his occupation, which was that of a jockey. He looked ill and weak, and was very anæmic, but his chief complaint was of the hæmaturia. He went to Egypt in February 1869, to follow his calling in the service of the Viceroy. The day after his arrival at Cairo, his leg was broken by a donkey he was riding slipping down and falling on him. In three months he left Cairo for Alexandria, where he remained nine months. He then spent six weeks in Cairo, after which he was in Alexandria until his return to England in November 1870.

The history of these cases is chiefly interesting so far as it may throw light on the introduction of the parasite into the body, and suggest preventive treatment.

Griesinger supposed that the young of the Bilharzia exist in the water of the Nile and in the fishes abounding in it, or even in the bread, grain, or fruit. Dr. Cobbold, however, in a paper published early last year, with which I infer that his views are still in accord, states that he thinks it more probable that the larvæ will be found in some of the gasteropod molluscs proper to the localities from whence the adult forms have been obtained. Unfortunately, his recent researches have not as yet proved this supposition to be correct, but it still, I think, suggests the direction for future inquiry. In my paper alluded to, I ventured to criticise the views put forth by Dr. Harley as to the probable mode of its introduction into the human system, as not in unison with those of the best observers. Perhaps I may quote the following sentence. "The view above given" (referring to Dr. Harley's paper) "is certainly opposed to what has been made out as to the deve-

lopment of other trematode worms, and I cannot but believe that it will be found eventually that man becomes infested with the parasite, not when it has just escaped from the egg, but when it has advanced to a stage beyond that of the ciliated embryo as we first see it, and that one or more intermediate hosts will be necessary before he is honoured by its presence."

Now, while my patient was at Alexandria, he drank water from a well; when he was at Cairo, he drank the water of the Nile. In both places, it was brought to the house in skins by Arabs. These men, by the way, used to wash their food, clothes, etc., in the spring at Alexandria. Generally he took the water filtered, but sometimes it was unfiltered. At Cairo, the Nile water was also brought in skins, and generally filtered, but occasionally, when thirsty, he took that also unfiltered. Before filtration, "it was sometimes very dirty-white in colour, and at other times clayey-looking." In Alexandria, he ate a good deal of water-cress, which he said was well washed. His living generally was good. I have no information as to his eating fish or shell-fish.

Now, the above details will not decide the source of infection. I incline to the belief that it was the unfiltered Nile water; but of course the water-cress, though said to be washed, might have conveyed small molluscs, etc., to the stomach.

It was not until January 1870, eleven months after reaching Egypt, that he noticed blood in the urine. It then came in single drops at the end of micturition, but gradually increased in quantity. A week after the appearance of the blood, he felt a burning sensation at the end of the penis after micturition, and a little pain during the act. He said that at one time a piece of clotted blood used to shoot out first, the urine then became clear, but never so clear as healthy urine, and towards the end of the flow it was freely mixed with blood. In August 1870, he had an attack of gonorrhœa, during which time he estimated that the quantity of blood, which was then much increased, amounted at the most to a pint in the twenty-four hours. This is probably an exaggeration, though not, I think, wilful.

During his stay in Egypt, he went under the care of three medical men from this country, none of whom appeared to recognise the cause of his complaint; as one treated him, according to the lad's statement, for inflammation of the kidney and irritation of the bladder, and the other two for stone in the bladder.

He had often great pain round the loins, especially in wet weather, and the testicles often swelled. During the twelve months of his illness he had lost colour, and his pallor was very noticeable when he was admitted, February 4th, 1871. His pulse was 72, small, weak, and irregular in rhythm. Nothing special had to be noticed as to the heart and lungs, except some irregularity in the action of the former. The tongue was slightly furred and moist, but apt to become dry at night. The appetite was generally fair, and the bowels somewhat confined. The abdomen was soft and normal in appearance. There were, however, slight fulness and tenderness on pressure at the epigastrium; and the liver extended a little below the ribs. The spleen was not enlarged. He complained of tenderness on the right side, where the lumbar and hypochondriac regions were grasped by the hand. There was, moreover, some tenderness on deep pressure over the hypogastrium. The desire to pass urine was rather troublesome, and he had frequently to get up three or four times in the night. In the twenty-four hours, he usually passed fifty to sixty ounces of urine, which contained a good deal of albumen, but no sugar. The reaction was acid, and the specific gravity ranged from 1007 to 1015. On standing, there was a dense and copious deposit, consisting chiefly of pus-cells, with blood-corpuscles dispersed among them, and here and there a small clot. Sometimes he passed urine into several vessels, in order that we might notice if the ova were more abundant at one period of micturition than at another. The following instance will sufficiently illustrate this.

Vessel No. 1. Urine of a pale greenish straw-colour. Deposit white and scanty, made up of pus, with numerous blood-corpuscles; but no ova observed.

No. 2. Slightly darker in colour; whitish deposit more abundant. This contained numerous ova and some escaped embryos.

No. 3. Dark brown in colour. Deposit very abundant, and containing numerous small blood-clots. Ova very numerous. In one small shreddy fragment, in which much blood was entangled, I counted twenty-two ova in the field at once, when using the one-inch object-glass. They were, therefore, much more abundant in the latter stage of micturition than the earlier, as, indeed, one would have expected; for the turbid portion of the urine would remain at the bottom of the bladder until the expulsive force, having drawn off the supernatant liquid, began to tell on the dregs.

During his stay in hospital, he never passed much blood at once, but