and ran off. He soon had to let the potato go, as you may suppose, but symptoms of oedema of the glottis set in, which necessitated opening the windpipe. Laryngotomy was performed by Mr. Tufnell, and the case did well. It is not long ago that I had to perform a similar operation on a man who swallowed a quantity of nitric acid, which gave rise to oedema of the glottis. The operation was performed after the admission of the patient into hospital, and in ten days the man went out perfectly well. And in connection with this subject, I may mention an out-of-the-way case, which I had under my care last year. It was that of a lady with rather severe symptoms of oedema of the glottis and swelling about the mouth and throat, which had followed the application of a leech to the gum a short time before I saw her.

The great danger in all cases of oedema of the glottis lies not so much in the oedema, as in the suffocating spasm which accompanies it. You may find a patient still breathing fairly well, but while you are at the bedside there comes a violent spasm, with the greatest difficulty of breathing and impending suffocation. This is the reason why the surgeon is often called upon to perform the operation at a moment's notice; and hence in a case of this kind it is advisable, as far as possible, to have the necessary instruments at hand in the ward in the hospital, and in the patient's house in private practice.

ON SPERMATORRHOEA AND ITS TREATMENT.*

By GEORGE G. GASCOYEN, F.R.C.S.,

Surgeon to the Lock Hospital; Assistant-Surgeon to St. Mary's Hospital.

FROM the brief description already given, it is not difficult to realise how a stimulus applied to or in the neighbourhood of the genital organs, or any sexual emotion, may produce such an effect upon the cerebro-spinal nervous centre as to influence the secretive function of the organs concerned; and we can also account for the instantaneous sexual excitement, almost electrical in its suddenness and rapidity, which the mere contact of these parts occasions. We can, too, thus explain the share the mind has in all sexual matters, as well as the copious secretion poured out by the urethra and its glandular adjuncts under excitement. We can now fully comprehend how healthy sexual excitement, whether from physical or from mental causes, may result in an emission of semen; and how its too frequent repetition may bring about a morbid excitability of the genital organs and of the brain until a state of complete nervous exhaustion is produced, both general and local, and this without any disease whatever in the parts of generation.

It follows, then, that in sleep the slightest contact of the sexual organs, too great warmth, position—anything, in fact, which causes an increased flow of blood to them—may arouse in the brain a lascivious dream, and be followed by genital excitement and an emission more or less pleasurable to the individual; and if this irritable condition of the nerves continue, emissions may become so frequent and scanty as to occur almost without the knowledge of the person until they prove a source of injury to him. When there is much irritability of the genital system, such slight causes as the friction of the clothes in walking, of the perineum against the saddle in riding, licentious talk, indecent pictures, etc., may occasion a slight loss of semen without the usual amount of erection and sensation, often causing the person great alarm and distress of mind, though needlessly. It is not uncommon to be told by a patient of imperfect erection, or even of its complete absence, with a too rapid ejaculation of semen when attempting intercourse, occasioning not only great disappointment and annoyance, but also the most serious apprehension that he is losing, or has lost, his virile power. The knowledge that there is a possibility of failure, together with the shame and vexation of such an occurrence, add much to the probability of non-success when again trial is made; and the loss of confidence thus engendered may really result in a functional impotency, though the structural capacity is adequate for the purpose.

When such a circumstance happens to a person who at some period of his life has indulged in sexual excess or self-abuse—even though many years have elapsed since he practised these habits, and then only for a short period and to a limited extent—he is certain to connect it with his former indiscretions, and to believe that he is permanently injured by them. Or, if it occur to a man who has never in any way abused his sexual powers, he will call to mind that in days gone by he occasionally had seminal emissions, and will attribute his failure to this cause, regarding it as evidence of debility, or of some defect in his reproductive organs. When once such an idea has crept into the mind, every sensation is eagerly watched for; and the slightest nervous

twinge or the least feeling of discomfort in these parts, which formerly would have passed unnoticed, is set down to some abnormal cause.

Constant mental uneasiness soon produces derangement of the digestive organs; heartburn, distension of stomach, flatulence, sense of oppression over the epigastrium, swimming in the head, flushing of the face, palpitation of the heart, shortness of breath, loss of appetite, constipation, occasional muscular pains, feeling of general debility, aching of the limbs, and especially of the back, with a host of other uncomfortable sensations, beset the unhappy man, until he believes that he is suffering from the effects of spermatorrhoea, the result of his former imprudence. The more he torments himself about this, the more urgent becomes his dyspeptic troubles; and, unless treated, he gradually sinks into a state of hypochondriasis, passing his time in noting his symptoms, and brooding over them to the exclusion of almost every other subject. The pain in his loins he attributes to disease of the kidneys or of the spinal marrow; the occasional shooting sensations through the body and limbs, to a grave disease of his nervous system; the feeling of oppression in his head, his forgetfulness and mental inactivity, to approaching loss of intellect. He becomes the prey of the most depressing thoughts; and entertains the firm conviction that his condition is incurable, and that idiocy or insanity is staring him in the face, until at length he falls into a slough of despond, from which it is very difficult to extricate him.

Although a certain amount of genital irritability may be present, there is seldom any positive disease about these patients; their complaint is at first almost entirely mental, until, by dwelling upon their imaginary ailments and morbid fancies, functional disorders are produced, which vary somewhat according to the caprice of each person, but which chiefly affect those organs he believes to be peculiarly influenced by the genital system. The patients who come to us to be treated for what they call nervous debility, impotency, etc., with nocturnal emissions, generally describe such symptoms as are referable to a deranged digestion, and the many abnormal sensations of which they complain arise chiefly from this. The emissions are due to an unnatural sensibility of the genital nerves, especially those of the urethra, resulting from sexual abuse, masturbation, or some other cause which has left them in a state of morbid irritability; and trifling causes, which in their normal condition have passed unperceived by them, act now as stimulants to them, and through them to the brain, causing an emission by reflex action. In these patients, also, the imagination is highly vivid or vitiated, so that a very slight local excitant will suffice to create a strong feeling of sexuality.

In treating these cases, the first thing is to secure the confidence of the patient, and to disabuse his mind of the conviction that he will never recover from his condition, and that his constitution is irretrievably injured; and also to persuade him that nocturnal emissions do not necessarily indicate any serious lesion of the parts of generation, but often the reverse. The next thing is to ascertain the cause of the emissions, which is very variable. If they arise from masturbation or venery, these practices must be discontinued; if from improper thoughts or conversations, all such subjects and objects as excite them must be avoided, as well as everything tending to venereal excitement. If the prepuce be long, so that the secretions collect beneath it and occasion erection from the itching and irritation, circumcision should be performed. Fissures of the anus, hæmorrhoids, ascarides, pruritus ani, etc., have sometimes the same effect, and must be treated. Pruritus scroti may produce erection and emissions in consequence of the scratching and rubbing induced by the intolerable itching; there will sometimes be no obvious cause for the itching, but a lotion of perchloride of mercury—about half a grain to an ounce of water, with a little spirits of wine—will generally stop it; whilst, if it proceed from pediculi, a lotion of double this strength will soon remove them.

Sometimes emissions will result from position, such as sleeping on the face or the back, particularly the latter; or from a custom of sleeping with the hands on the abdomen, or in the neighbourhood of, or in contact with, the genitals. Such habits must be corrected. Emissions will often result from enforced abstinence after a course of sexual indulgence, or after the sudden cessation of marital rights; hard physical exercise, so as to produce fatigue, is the best remedy, for then the brain and the body are too tired to run riot in libidinous dreams and practices.

Little or no treatment, either local or general, is necessary for emissions which arise from the above causes, but merely the application of common-sense measures. Yet, if an irritable condition of the genital organs be established, with frequent emissions, we must then bring to bear such local and constitutional remedies as are at our disposal.

In these cases it must be remembered that the nerves of the genito-urinary passages, and especially those of the urethra, are in a morbidly sensitive condition; that the mucous membrane of the urethra and neck

* Concluded from page 69 of last number.

of the bladder is congested, with often a copious secretion of thin watery mucus; that the prostatic and other venous plexuses within the pelvis are enlarged and engorged in consequence of the frequent afflux of blood to these parts; whilst the peripheral irritation of the genital nerves has caused functional disorders of various kinds.

The occasional introduction of a catheter as large as the urethra will take, is often of the greatest service; it should be passed into the bladder and allowed to remain for five or ten minutes, according to the tolerance of the patient; its mechanical pressure helps to unload the congested capillaries and small vessels of the urethra; its contact deadens and destroys the extreme sensibility of the urethral nerves, and renders them less susceptible to the influence of slight excitants; whilst, by stimulating the muscles, it provokes their contraction, and so renders material assistance in emptying the larger veins. A silver catheter is the best instrument for the purpose, as it exerts firmer pressure than an elastic bougie; and, as the urine can be drawn off through it, the patient will not require to micturate for several hours, which is a point of some importance, as the urethra is often very tender after the passage of an instrument for the first few times. The frequency with which it should be employed depends upon the amount of discomfort its presence occasions; and, if the pain be great, it should not be left in more than a few seconds, lest rigors, swelled testicle, etc., be occasioned. Sometimes the urethra is *extremely* sensitive, and much pain attends the use of the catheter; but this is an additional reason for persisting with it, though a smaller one may be employed at first, so as to cause less pain. I have sometimes found that smearing the catheter with blue or calomel ointment, or with half a grain to a grain of nitrate of silver rubbed down in an ounce of lard, to be of use in obstinate cases; but I prefer the blue ointment to anything I have yet tried. Some camphor, extract of opium, belladonna, etc., may be combined with these ointments if thought desirable. Care should be taken that these applications do not reach much beyond the curve of the instrument, and it should be thoroughly well oiled before using it. The over-secretion of mucus is always checked by the use of the catheter, whether armed with ointment or not.

Cold bathing, cold douches, etc., should not be employed on going to bed. The ordinary bath in the morning does good; but cold applications at night should be forbidden, as the reaction which follows them will increase the local circulation, and so cause congestion and erection of the penis, and increase the probability of emissions.

Not only must the position assumed in sleep be attended to, but undue warmth in bed avoided, whether by using very soft beds or too large an amount of clothing. The bowels should be carefully regulated, to prevent any accumulation within the rectum; and the urine examined from time to time so as to detect an excess of uric acid, the presence of oxalates, etc., which may render its passage irritating to the hypersensitive urethra. Over-distension of the bladder must at all times be guarded against, and the patient warned to pass urine on waking in the morning, lest he doze off again with a full bladder, which is one of the most certain provocations of erection and emission.

Before commencing to treat this affection constitutionally, it is generally necessary to allay the digestive disturbances which are so common and often so severe, by giving such remedies as may be applicable to the condition of the patient, either with or without the more special medicines. By neglecting to do so, we may not only add to the dyspeptic troubles and obtain no benefit from the drugs given, but a valuable medicine may do harm and be brought into disrepute, in consequence of its being administered at a time when the stomach cannot tolerate it.

Internally, I have found astringents of more use in this disorder than tonics; or they may be combined. Gallic acid, the dilute mineral acids, especially the sulphuric, may be given. Tincture of matico will often be of service, and more so, in my experience, than any other plant rich in tannin, as it appears to act upon the genito-urinary track, rather than upon the bowels, as is often the case with the others.

Ergot is one of the most valuable remedies for this affection, and the liquid extract of the *Pharmacopœia* is a very efficient and convenient form for giving it; whilst the dilute sulphuric acid can be added, if thought advisable.

When the urethra is very sensitive, and the passage of urine painful, small doses of copaiba are often most comforting; or the other oleo-resins may be tried if it disagree; but none of them, in my opinion, is equal in value to copaiba, when it can be borne.

I am not disposed to regard strychnine in these cases with very great favour; when there is much irritability of the nerves, I believe it often adds to this; but when this is subsiding, it may be of use as a tonic: so may quinine or iron, but in no other way. I have never given the tincture of iron in the enormous doses (from one to two drachms three

times daily) recommended by some, and so I cannot speak personally of its value in such large quantity.

Cantharides, phosphorus (except the dilute phosphoric acid), and the so-called aphrodisiacs, do harm by acting as stimulants to the nervous system generally, and therefore to the local nerves. Cantharides, also, by its action upon the bladder is, especially when given in large doses, a very injurious drug in these cases. For the same reasons, I disapprove of local blistering; whilst the sore left by the blister acts, moreover, as a source of irritation, and adds to the liability of emissions.

Belladonna, in my hands, has proved to be an uncertain remedy; in some cases it has appeared to do good by allaying irritation, whilst in others there were no beneficial results from it. The dryness of throat, disturbance of vision, and diarrhoea, which are often caused by it, constitute an objection to its employment in full doses, and without them its value is very questionable.

Camphor is a most useful drug; three or four grains made up into two pills, with half a grain or a grain of opium and one or two of aloes, have more frequently allayed irritability and prevented emissions, than anything I have yet tried. Opium alone does not succeed as well, and a larger dose is necessary, so that the untoward symptoms sometimes produced by it are more likely to be incurred.

I have tried chloral in a few cases, and with very great advantage: in doses of fifteen or twenty grains at bedtime, it has answered its purpose admirably.

Bromide of potassium, in thirty or forty-grain doses, will sometimes be of service; but it seems to me a less certain remedy than chloral, which I am disposed to regard as one of the valuable agents we possess for these cases, though as yet my experience of it is limited.

Suppositories vary much in their action whatever drugs they may contain. Occasionally they answer well; but often they do not lessen, and I am not sure that they do not sometimes increase the irritability of the parts.

Galvanism I have not employed myself; but in the few instances where I have known it tried by others, it has seemed to me to do more harm than good, by adding to the nervous irritation.

Lastly, as to cauterisation by the *porte-caustique*, I need scarcely say that I am strongly opposed to this method of treatment; for, if my view of this disorder be correct, this instrument can relieve it in no other way than as the passage of a catheter does. I do not believe that ulceration or other morbid conditions of the ejaculatory ducts are the causes of seminal losses. We have no evidence that these pathological conditions exist except, it may be, in very rare instances; and, if so, the application of nitrate of silver to the prostatic mucous membrane in every case of nocturnal emission must be unnecessary, and, in spite of its alleged harmlessness, I consider it to be a dangerous treatment. I have known two persons die from the effects of the *porte-caustique*, and I have seen others suffer severely from its employment. This may not be the usual result; but I do say that the application of nitrate of silver to the urethra, whether in stick or in strong solution, is, at least, a very sharp remedy, and will often produce violent inflammation, and sometimes lay the foundation of a stricture or of a chronic irritation of the bladder. If, then, caustic be applied on an incorrect surmise as to the condition of, and its effects upon, the prostatic mucous membrane and ejaculatory ducts, it is not only an unnecessary, but, in my opinion, an unsafe method of treatment.

WHAT CASES SHOULD BE SENT TO LUNATIC ASYLUMS? AND WHEN?*

By T. S. CLOUSTON, M.D.

Superintendent of the Cumberland and Westmorland Asylum, Carlisle;
President-elect of the Cumberland and Westmorland Branch.

THE question of sending a patient to a lunatic asylum is one of very great interest from a medical, a legal, and a social point of view; and, unfortunately, its interest to us is by no means that of an abstract question. It must be solved by us frequently and at short notice; and by its wrong solution most grave consequences may, in some cases, happen to ourselves, our patients, and the public. It cannot be unprofitable, therefore, to discuss it shortly; and I propose to confine my remarks chiefly to the medical aspects of the question.

Less than fifty years ago, the question of sending an insane patient to a lunatic asylum was a very simple one. It was determined solely by the fact of the patient's being dangerous to others or to himself, chiefly by the former consideration. After that, there was a decided tendency among the medical profession to simplify it in quite the opposite direc-

* Read at a meeting of the Cumberland and Westmorland Branch.