septic dressing applied. The result, as you see (patient again shown), has been very satisfactory; the wound is soundly healed, and the patient appears to have suffered little either locally or constitutionally from the operation.

REMARKS

ON

HETEROGENESIS IN ITS RELATION TO CERTAIN PARASITIC DISEASES.

By H. CHARLTON BASTIAN, M.A., M.D., F.R.S.,

Professor of Pathological Anatomy in University College, London; Physician to University College Hospital; and Assistant-Physician to the National Hospital for the Epileptic and Paralysed.

II.

THE transformations which I have already described, of the particles in the latex of plants, and the somewhat similar transformation of milk-globules, are most important and typical. Here the change takes place in comparatively large masses of matter, which can be watched with ease; and, moreover, it takes place in masses which, although they are the products of living bodies, can scarcely themselves be said to be "living". There is, perhaps, most room for doubt, in the latter respect, concerning the particles in the latex; and these, being probably poor in nitrogenous materials, evolve into very small and simple organisms. The milk-globules, however, have a highly nitrogenous composition, owing to an admixture of albuminoid constituents with their fatty elements—a combination which seems especially favourable for the occurrence of evolutionary changes. The milk-globules accordingly are seen to produce large specimens of *Penicillium* of a remarkably vigorous growth.*

In the blood we have another highly nutritive fluid, which might be supposed to be capable of giving birth to independent living things under certain conditions. It seems, however, to undergo such changes more frequently in the lower forms of life than in the higher. Amongst insects, several instances may be mentioned in which simple organisms appear to be born from the fluid constituents of the blood, or else produced by modifications of some of its already existing solid elements.

This seems to be the case with 'muscardine', the disease which formerly committed such fearful ravages amongst the silk-worms of France.

It was ascertained by Bassi in 1835, that one of the most prominent features of this disease was the presence of a fungus which at first increased and multiplied within the body of the living animal, and, after the death of the worm or moth, made its appearance externally—coming through the skin in numerous situations as a whitish powdery growth. It was afterwards ascertained by M. Audouin and others, that the disease was not confined to the silk-worm and its moth, since it could be communicated directly by inoculation to many other species of Lepidoptera; and it could also be engendered quite easily in these and in the silk-worm by shutting them up and feeding them for a time in close damp bottles or boxes. The possibility of inducing a "spontaneous" outbreak of this contagious disease was always within reach of the experimenter, even in districts which were, so far as all previous knowledge went, wholly untainted. In this respect, muscardine was found to be similar to typhus fever.† The question arises, however, whether, in such cases of apparently "spontaneous" origin, the unhealthy conditions merely induce a state of the blood and body generally in which omnipresent, though unknown, spores are enabled to develop;‡ or whether the state of the blood thus initiated does (as it

seems) suffice to give rise to the germs and fungus-growths which afterwards constitute such all-important elements of the disease. The former is the view which has been most widely adopted: and yet two of the ablest writers on muscardine, after the most careful investigations with reference to this very point, came to the opposite conclusion. These observers, who adopted in its entirety the belief in the possibility of engendering muscardine *de novo*, were M. Guérin-Ménéville and M. Robinet. The conclusions of the former especially were based upon positive and apparently unambiguous observations.

The blood-corpuscles of the silk-worm during life are elliptic or more or less elongated, but after death they are always found to be spherical. When a little blood is abstracted from a healthy worm, the corpuscles are elongated at first, though they speedily assume the spherical condition; and, when in this latter state, they begin to exhibit amœboid protrusions, although such changes of shape are never seen in healthy blood-corpuscles immediately after they have been drawn from the body. M. Guérin-Ménéville observed that in dead silk-worms,* and also in the cases where blood had been drawn from living animals during the very early stages of the disease, that the spherical amœboid corpuscles contained much larger granulations than usual; and that some of them tended towards the periphery of the corpuscles, from which they ultimately made their exit. These little bodies were ovoid, and from $\frac{25}{1000}$ — $\frac{30}{1000}$ in length. They moved

Fig. 3.—Illustrating the development of *Botrytis Bassiana* in the Blood of Animals suffering from Muscardine (Guérin Ménéville). X400.

a, a. Spherical and amœboid blood-corpuscles containing large particles. b, b. Ovoid particles in the free state. c, c. Germs of *Botrytis* supposed to be derived from such particles, which gradually grow (d, d) into long and simple, and subsequently (e, e) into branched, Fungus-filaments.

about in the serum of the blood more actively than could be accounted for by mere Brownian movements. They gradually increased in size; and, even two or three days before the death of the sick worms or moths, many of them had become so elongated as to be easily recognisable as rudimentary fungus-filaments. Along with this characteristic feature [of muscardine, another very significant peculiarity invariably showed itself—i. e., a well-marked acidity of the blood becomes manifest, instead of the neutral condition which is invariably met with in the healthy worm or moth.†

This fungus, continuing to grow through the tissues of the animal in all directions, soon causes its death; and, after twenty-four hours or

* The conversion of fatty-looking intestinal globules of uncertain origin into Monads and Amœbæ has been fully described by Mr. T. R. Lewis, in his Report on the Micros. Objects found in Cholera Evacuations. Calcutta: 1870. Pp. 26-29.

† Muscardine, however, is undoubtedly associated with the development of a Fungus in the blood; whilst in typhus fever no lower organisms are known to be produced. Their absence in the latter disease is further testified by the large number of recoveries; though, on the other hand, like other diseases in which the blood is affected and numbers of organisms are developed in it, muscardine is invariably fatal. As Audouin says, "dans tous les cas le résultat est le même; aucun de ceux qui sont atteints n'échappe."

‡ It is admitted on all sides that such a predisposing cause is necessary. Thus Robin says: "Il ne suffit pas que les spores arrivent à la surface du corps des larves ou même dans l'intérieur; il faut encore, pour qu'elles puissent agir sur l'animal, que ses humeurs présentent des conditions favorables à leur germination. Si les Vers exposés au contact et à la pénétration des spores sont placés dans des circonstances d'alimentation très favorables, il se pourra probablement que les humeurs dont les principes se renouvellent rapidement, ne présentent pas ces conditions; on dit alors que l'animal résiste par son énergie propre. C'est là certainement, dit Robinet, la ressource la plus sûre et la plus facile à créer par la magnanerie. C'est ainsi que dans des invasions désastreuses de la muscardine une foule d'individus

échappent, non pas à transmission, ainsi qu'on le dit, mais bien au développement des spores. Voilà pourquoi beaucoup d'observateurs ont contesté la transmissibilité de la Muscardine d'un individu à un autre." (*Végétaux Parasites*, 1853, p. 582.)

* In reference to the different conditions under which the same phenomena may be witnessed, it should be remarked that the Fungus can always be developed in dead animals. According to Robin, this was shown by M. Johannys in 1839; for, as he says, "il a pu faire développer ce végétal sur des Vers à soie morts, placés dans des conditions favorables à la fermentation et hors de toute communication avec des lieux infectés de Muscardine. La moisissure se développe aussi bien sur l'animal mort que sur les animaux inoculés de leur vivant, et le végétal obtenu par inoculation est identique avec celui, qui se développe sans cause directement connue." Whilst, according to evidence given by M. Guérin-Ménéville in 1851, "la muscardine sporadique est une terminaison naturelle de l'existence du ver à soie, maladie qu'il est impossible de prévenir d'une manière absolue" (*loc. cit.*, pp. 574 and 603).

† From the evidence now in our possession, it is not perhaps possible to say whether the acidity precedes the appearance of germs, or *vice versa*. We do know, however, from the experiments of Dutrochet, that a mixture of egg-albumen and water will remain unchanged for months, but will always develop fungi within a few days after it has been slightly acidified.—*Ann. des Sc. Nat.*, 1834 (Botan), page 34.

less, it may make its appearance externally through the thinnest portions of the dorsal integument, and through the respiratory spiracles, in the form of delicate snow-white tufts of fungus-mycelium. These continue to grow rapidly, and soon produce multitudes of minute spores,

Fig. 4.—More developed form of *Botrytis Bassiana*, as it grows through the tissues of animals suffering from Muscardine (Audouin). X 400.

by means of which the disease may be disseminated amongst those silk-worms which, owing to bad hygienical conditions or other causes that may have lowered their vitality, are at all predisposed to initiate the morbid changes which are peculiar to the disease.

In its epidemic form, muscardine is most prone to occur at the season when the worms are just about to pass into the chrysalis condition. At this critical period, unknown influences are apt to operate upon them which produce abnormal and specific nutritional changes; and then the disease breaks out like a pestilence, attacking almost all the worms of a given locality.

Another general parasitic disease to which silk-worms are liable has been called 'flacherie'. It is generally acknowledged to be due to a kind of fermentation or unnatural state of the mulberry-leaves on which the worms feed. The affection is characterised by the presence and multiplication of very minute ovoid corpuscles, having a close resemblance to *Torula*. They multiply rapidly by germination, and sometimes are to be found in chaplet-like groups. They are seen in the stomachs of worms which feed upon the decaying leaves, and subsequently may be recognised in myriads throughout the blood and all the tissues of the animal. This affection is, moreover, like 'pebrine' (of which I shall subsequently speak), constantly transmitted to any young that may be produced by diseased moths.

The affected leaves doubtless bring about some important alterations in the fluids and tissues of the animal which feeds upon them, of such a nature as to lead to the evolution of the organisms characterising the disease. An enormous amount of evidence exists tending to show that the consumption of mouldy or fermenting articles of food, whether by higher or by lower animals, is not of itself sufficient to inoculate the system so as to cause general parasitic diseases; and some very strong facts in support of this view will be subsequently cited. Fungi would, indeed, soon destroy all higher forms of life, if the popular belief were true. In the production of 'flacherie', therefore, we are driven to believe that the unhealthy food is the immediately determining cause of some general constitutional condition, which, in its later phases, leads to the evolution of the minute *Torula*-like organisms in the blood and throughout all the organs and tissues of the animal affected. In these respects, the history of this parasitic and highly contagious affection is very similar to that of muscardine, [although 'flacherie' is more distinctly engendered by bad food, whilst muscardine is produced in well-fed worms, so long as they are kept in a close and damp atmosphere.

Again, it is well known that the growth of a mould (*Empusa*) frequently seems to prove fatal to flies in autumn.* This subject has been investigated by Professor Cohn, who came to the opinion that the parasitic growth commenced by an independent development of Fungus-germs in the blood of the sickly animal. He does not believe that the disease is incited by the mysterious introduction of Fungus-germs into their circulating system from without. He sums up by saying (*Hedwigia*, 1855) that "the influence of the spores of *Empusa* in the appearance of this fungus, and of the disease in flies, is by no means evident, since the genesis, the chemical and optical characters of the numberless free cells in the blood, the absence of a special ex-

panded mycelium, and, above all, the whole history of the development, seem to concur in favour of the origination of the cells of the *Empusa* from the diseased blood."†

Although much has been written of late years with reference to the existence of germs of Fungi in the blood of man as producers of many of the most serious diseases to which he is liable, I have elsewhere stated (*BRITISH MEDICAL JOURNAL*, Oct. 7, 1870) how little evidence there is in support of these various statements, and how much other evidence tends to contradict them. As a matter of fact, such organisms are not to be found in the blood of living persons, except in certain rare affections; and doubtless many of the alleged cases are to be explained by the altogether unwarranted assumption of observers that mere particles, which are so commonly met with in the blood, are examples of the so-called "micrococci" of Hallier. But it must always be remembered that there are particles and particles. Particles which are potential fungus-germs cannot be discriminated by the microscope from normal blood-particles; and even if some of these particles were seen to develop into rudimentary fungi, it would by no means prove that they came from fungi, in the face of what we now know concerning the metamorphosis of milk-globules and other organic particles.

There are, indeed, weighty reasons why such a fluid as the blood should not, in a very highly organised creature, easily give birth to heterogenic products during the life of the individual. It is not a mere excretion like the milk, but the most nearly vitalised of all the fluids of the body, and subject to constant changes from moment to moment in all its tissues: so that, as long as the organism lives, the united molecular activities of the various tissues are continually influencing this all-pervading fluid, and tending to maintain its ordinary characters. When death is approaching, however, these united activities become weaker and weaker, and changes may begin to take place in the blood, more closely resembling those which occur in organic fluids out of the body.† It is quite conceivable, also, that the changes which occur in the blood in certain febrile diseases—more especially when these are associated with a very high temperature—may be of such a nature as to make the blood more than usually prone to undergo rapid putrefactive changes after death. And the occurrence of putrefactive changes in animal fluids implies the presence of *Bacteria*.

Facts can, moreover, be cited tending to show that the changes in the blood in these diseases do predispose it to early putrefaction, and that the living organisms which have been observed soon after death must have been produced *de novo* in the organic fluids themselves. One observation of this kind made much impression upon me at the time, and tended strongly to confirm a then growing belief in the truth of doctrines concerning "spontaneous generation". Having made an autopsy (thirty-two hours after death) nearly three years ago, on a man who died in University College Hospital of rheumatic fever, in whom an exceedingly high temperature had existed for a few hours before death, I immediately afterwards proceeded to examine portions of the brain and membranes with the microscope. The skull had been opened and the brain had been removed in my presence but a few minutes before, when I cut through the arachnoid, and carefully separated two convolutions which had previously been in close contact, in order to cut off a portion of the delicate network of vessels lying between them. On submitting this to the microscope, the fluid outside the vessels and also that within, was seen to contain a large number of most actively moving particles. Many of these were mere spherical particles of various sizes, though others were distinct and large *Bacteria*, made up of two almost cellular segments. And every portion of the pia mater examined showed similar moving particles and *Bacteria*. The brain was then covered by a bell-glass, and when portions of pia mater—again taken from between previously unseparated convolutions—were examined after an interval of twenty-four hours, the large *Bacteria* had considerably increased in number, whilst the small spheroidal particles seemed to be as plentiful as before. When portions of brain-substance from some central parts of the organ were also examined at this time, moving particles and *Bacteria* were seen to exist in the greatest abundance amongst the disintegrated nerve-elements.

Now, with regard to the origin of those *Bacteria* which were observed in the vessels a few minutes after the brain had been removed from the body, it is, in the first place, perfectly obvious that they must have existed in the blood of these vessels before the brain had been removed and before the skull was opened. *Bacteria* are not produced in any fluids under two or three hours. Their origin could not, therefore,

* Of course it is not denied that such a disease is capable of being spread by contagion: far from it. We believe that this does occur. The establishment of the mode by which contagion is communicated cannot, however, as some have appeared to think, dispose of the more general question.

† It is quite certain that refuse fluids, such as urine in certain diseased conditions, even in man himself, may contain an abundance of *Bacteria* and *Torula* at the moment when it is passed from the body.

* This mould will always appear on a dead fly which is allowed to float for a few days on the surface of water.

have been due to *Bacteria*-germs derived from the atmosphere, which, on removal of the skull-cap, had in some mysterious way insinuated themselves into the blood-vessels. They must either have existed in the blood during life, or else they must have been produced *de novo* in this fluid after death. There is strong reason for disbelieving that *Bacteria* existed in the blood during the life of the individual. I have several times examined the blood of individuals who were similarly affected with this exaggerated form of rheumatic fever, and have always failed to discover any such organisms.* It is, therefore, by far more likely that they had been newly evolved owing to changes taking place in the blood after death, in or near the situation in which they were found. It would be impossible otherwise to account for their distribution throughout the brain† at a time when the circulation had ceased. This organ is so far, comparatively, from any mucous surface, that even if germ-particles had been able to make their way into the blood-vessels near their surface (which is in itself altogether a gratuitous supposition), it would be impossible for us to imagine that, in such a short space of time, they could have been able to penetrate into the innermost parts of the brain. Their unceasing movements are exceedingly slow in reality; and more than this, they are never continuously progressive. They consist either of slow oscillations, or else of short darting movements hither and thither, in which the same ground is frequently retraced. And again, if *Bacteria* in their adult or in their rudimentary state could make their way from the atmosphere through the superficial layers of the mucous membrane so as to penetrate the vessels—why, if it is to be assumed that they do this after the death of an individual, does not the same thing occur during his life? It cannot be replied that *Bacteria* are unable to live in the blood during life, since the observations of Davaine, Vulpian, and others, clearly show that they can live in the blood of man and also in that of several of the lower animals during their life. There seems, then, to be no reasonable alternative; and we are compelled to fall back upon the assumption, that the *Bacteria* met with in our observations had been evolved out of the blood-plasma and other fluids of the body, just as we have seen that they arise in previously heated organic infusions.‡

During life, and under the influence of all the varied activities of the living body, the plasma of the blood, rich in nutritive materials, is probably constantly giving birth to living particles which speedily develop into leucocytes. These amoeboid corpuscles are the organisms into which such new-born living matter invariably tends to develop in the healthy living body. But when death has supervened, then all is changed; the molecular composition of the fluid may have altered, whilst the activities of the tissues which formerly influenced it have ceased to act. It is, however, a fluid still rich in albuminoid materials; and now that it is released from all the vital influences of the organism of which it formed part, what wonder is it that the new-born particles which are still evolved should assume the familiar shapes of such units (*Bacteria*) as appear in organic solutions outside the body?

Possibly also some of the minute particles which existed in the blood before death was close at hand, may actually alter their destination and develop into *Bacteria*, just as the milk-globules develop into large Fungus-germs.§ And it is also quite possible that a heterogenetic change, similar in nature though different in kind, may overtake red and white blood-corpuscles when they are liberated from the vital influences of the organism in which they have been produced, and also occasionally within the living organism itself. On this subject, however, we have very little evidence at present.||

* Even if they had existed, however, during life, there would still be weighty reasons inclining us to believe that they had been produced *de novo* in the blood, rather than that they had been developed from germs which had gained access to the blood from without.

† In all probability, if examination had been made, they would have been found disseminated throughout all other parts of the body, just as they were actually found in the brain. (See BRIT. MED. JOURNAL, Oct. 7th, 1870.)

‡ See *Modes of Origin of Lowest Organisms*, pp. 18 and 67. The recent investigations of Dr. Sanderson have, moreover, led him to the conclusion that the blood does not naturally contain either visible or invisible *Bacteria* (*Thirteenth Report of Medical Officer of Privy Council*, 1870, p. 65), whilst Professor Lister's antiseptic method of treatment is based upon this belief, as one of its starting points.

§ And such an occurrence would be accounted by Professor Hallier and his school as evidence that Fungus-germs pre-existed in the blood (derived from pre-existing Fungi), and that they were causes rather than consequences of antecedent morbid conditions.

|| White corpuscles are practically young Amœbæ, and there is no saying what changes they may not occasionally be capable of undergoing. *Gregarina*, which are so very abundant in the bodies of lower animals, and which are closely allied to Amœbæ, may perhaps in many cases be derived from transformations of leucocytes. Again, red corpuscles are very similar in many respects to the chlorophyll vesicles of *Alga* and *Characea*, although the latter are probably much less specialised in structure. But I shall show elsewhere that the transformations of such chlorophyll vesicles are often of the most startling description. Quite recently, Mr. Ray Lankester (*Quarterly Journal of Microscopical Science* 1871) has described a peculiar, though small and simple, ciliated Infusorium which he found in the blood of certain frogs; whilst Dr. Boyd Moss (*Monthly Microscopical Journal*, Oct. 1871)

The two modes of origin of organisms to which we have just alluded may, however, lead to the presence of *Bacteria* and larger Fungus-germs within the midst of cell-elements, both in plants and animals.

Myriads of microscopic Fungi belonging to the protean types included under Rust, Smut, Mildew, and Mould,* are habitual dwellers in and upon the surfaces of living plants, especially when they are in a sickly condition—although others are often found in and upon plants which present no other sign of disease. These particular kinds of fungi are encountered only in such situations, and they recur in similar habitats with tolerably constant forms. The ravages of many of them are matters of no small importance to mankind on account of the very serious damage which they help to produce upon our food-supplies. We need only mention the fatal "blights" which they are apt to occasion amongst our cereals, and those devastating diseases of the vine, the hop, and the potato, in which Fungi of this kind appear as the most active agents of destruction. The original mode of origin of these various growths is still involved in doubt and obscurity, though most positive statements have from time to time been made by different observers with reference to their heterogenetic origin,† by changes similar to those which suffice to convert the milk-globule into a *Penicillium* and the globules of many plants into *Amylobacters*. Thus in speaking of the blight produced by *Uredo*, M. Turpin says (*loc. cit.*, p. 346): "Il m'est bien démontré par un grand nombre d'observations, faites sur diverses plantes plus ou moins attaquées de l'Uredinée, que la carie n'est qu'un état morbide, qu'un dégénérescence de la globuline ou féculine du tissu cellulaire du péricarpe du grain de blé." A similar change, however, may also occur in the globules which naturally exist in the cells of the stem or in those of the leaves; so that, according to Turpin, "L'Uredinée est une maladie qui attaque par place, les globules contenus dans les vésicules du tissu cellulaire des plantes, qui leur donne quelquefois plus de volume et toujours les couleurs blanche, jaune, aurore et brune, par lesquelles les mêmes globules passent dans les feuilles qui prennent toutes ces couleurs à l'automne. Ces globules ainsi viciés, peuvent ensuite, par contagion ou par inoculation, altérer de la même manière ceux de la plante nouvelle. . . . Cette affection spéciale est au grain de globuline du tissu cellulaire ce qu'est celle de l'ergot au grain du seigle, du froment et de l'ivraie tout entier." Thus the mode of origin of the blight seemed so indubitable to Turpin, that he was led to suppose the products were mere pathological modifications of pre-existing structures, not possessing an independent life of their own. Several other celebrated botanists, moreover,—amongst whom we may name Fries, Endlicher, and Unger—were equally certain that this and many other Entophytes are derivable from morbid portions of the tissues of plants, although they recognised the fact of their developing into independent living organisms.‡

[To be continued.]

XYLOL IN SMALL-POX.

I have, during the last few weeks of the prevailing epidemic of small-pox here, used xylol with most satisfactory results. I do not, however, agree with giving it in capsules, as recommended, simply on account of the great difficulty generally experienced by patients in swallowing during the severer stages of the disease; nor is it miscible with water alone, as stated. I have given it in doses to adults of 20 minims every three or four hours, either in milk or mixture of almonds; the latter I prefer. I should be glad to hear of its extensive use, as I believe in its efficacy, although I cannot define its specific action.

H. B. HURMAN, Bridgewater.

has described a similarly simple infusorium found in the blood of a Ceylon red deer on several occasions. It seems to me more easy to suppose that such organisms should have arisen by a heterogenetic process, than from 'germs' of delicate, externally existing organisms of this kind, which had not only made their way into the blood, but were capable of flourishing there.

* See a useful little book by M. C. Cooke, entitled *Microscopic Fungi*, 2nd ed., 1870.

† It has been noticed that the leaves of many plants, prior to the appearance of fungi within them, have been remarkable for their "almost unnatural green colour"; and, according to Mr. Cooke (*loc. cit.*, p. 155), "this phenomenon has been noticed in ears of corn, in which every grain was soon afterwards filled with spores of bunt." This fact is one of much interest and importance, since I shall subsequently show (in *The Beginnings of Life*) that the same extremely bright green colour is almost invariably to be observed amongst those portions of *Alga*, or amongst *Euglena* and *Desmidae*, when they are about to undergo a heterogenetic change; and in these latter cases every step of the process of transformation into new organisms may be watched by the microscopist.

‡ We quote the following note from M. Pouchet (*Novelles Expériences*, 1864, p. 117): "Fries, qui classe ces plantes parmi les champignons, les décrit ainsi: *Entophyti vegetatio nulla. Sporidia ex anamorphosi tele cellulosa plantarum vivarum orta; sub epidermide enata et per hanc erumpentia*. Endlicher est encore plus explicite. Voici ce qu'il dit: *Sporidia varia e parenchymate morbo plantarum vivarum sub epidermide orta, hæc rupta erumpentia, et varie sæpè mutata stipata*. Fries, *Syst.*, iii, p. 501; Endlicher, *Genera Plantarum*, p. 16." On this subject, see also Unger in *An. des Sc. Nat.*, vol. ii, n. s., p. 209.