able time in a feverish state. The prognosis formed at the time was most unfavourable, as a lacerated and penetrating wound of the bladder was evident; but no inflammatory symptoms succeeded, and the patient left the hospital in a week well. The bladder having been in a state of fullness, it is almost certain the peritoneal sac had not been opened."

To recur, however, to the cases of rupture occurring in the female: the cases to which I have already alluded were those of Mr. Maurice Collis and Professor Smith. The particulars of the first of these cases are as follows. "The patient, who was a woman of intemperate habits, retired to bed on the night of September 7th without relieving her bladder, and got up in the course of the night for that purpose. Not succeeding, she endeavoured to get into bed again, but fell across the footboard. She fainted at once, and was lifted into bed; and in the morning complained of inability to pass urine, and said that she had felt something give way. On the third day after the accident, she was brought to the Meath Hospital in a state of collapse. A large male catheter was introduced, and about half a pint of urine, mixed with blood and clots, drawn off. About the same quantity trickled away afterwards, and she voluntarily passed water more than once. She died upon the following day, the fourth from the date of the injury. There was an ecchymosis on the front of the abdomen, where the blow had been received. Upon examination *post mortem*, a large transverse rent, which admitted the points of the fingers, and with ragged irregular edges, was found in the anterior wall of the bladder, which was in a contracted state. The large aperture communicated with a gangrenous cavity of considerable size in the neighbouring cellular tissue. There was a communication with the cavity of the peritoneum, which, however, contained a pint of turbid serum, and was inflamed in the vicinity of the bladder."

In the case recorded by my colleague Professor R. W. Smith in the *Dublin Journal of Medical Science* (vol. xxv), a female, aged 50, while in a state of intoxication, fell across the edge of a tub, ruptured the bladder, and died at the end of five days. There was a large transverse rent in the bladder, situated at its posterior and upper part, where it is covered by peritoneum.

PUERPERAL APOPLECTIC CONVULSIONS.

I WAS called at 9 P.M. on Thursday, February 15th, to Mrs. E. S., aged 30, who was in labour with her fifth child. She had been ill since about one in the morning, and the case had been progressing quite favourably until the time when I arrived at her house. I found, upon making an examination, that the presentation was natural and the pains strong. The condition of the patient at the time being good, I left her, and went down stairs; but in less than ten minutes I was called by the nurse in great haste, and, on returning to the bedside of the patient, found that she was quite insensible. The tongue protruded from the mouth, and was most severely bitten; she had loud stertorous breathing and contraction of the pupils; in short, all the symptoms of an apoplectic seizure. At this time the membranes were protruding, but very thick; I therefore at once punctured them, and, the liquor amnii having escaped, the child was born, and the placenta expelled in less than twenty minutes. The patient was all the while in a comatose state, in which condition she remained until 3 o'clock next morning, when she died, twenty-six hours from the commencement of the labour, and five hours and a half after the child was born. Cold water was at first dashed over the face, and afterwards cold was kept to the head by means of cloths dipped in cold water and vinegar. Sinapisms were applied to the legs and to the nape of the neck, and likewise hot water in bottles to the feet. Venesection was not performed on account of the hæmorrhage, which was copious. The child, I may state, was dead, and had, in my opinion, been dead for some weeks, judging from the peculiar discharge with which the patient had been troubled for some time, and which had caused her great anxiety.

I may state in connection with this case, that there had been no premonitory symptoms to warn one of an approaching attack; on the contrary, she had been going on through the day as comfortably as in any of her previous confinements. Mrs. E. S. was naturally of a nervous disposition; and although rather muscular, she was of a flabby fibre: even the thought of having to undergo a vaginal examination would throw her into a state of hysteria, as was shown by her diffidence to be examined three or four weeks before the full time in order to ascertain the source of the vaginal discharge.

I find that Dr. Fleetwood Churchill of Dublin, in his *Theory and Practice of Midwifery*, attributes these cases in some instances to anxiety of mind, and therefore I cannot but refer the above case to the same cause.

JOHN WILLIAM JONES, L.K.Q.C.P., M.R.C.S.E., Coleshill.

REMARKS

ON

HETEROGENESIS IN ITS RELATION TO CERTAIN PARASITIC DISEASES.

By H. CHARLTON BASTIAN, M.A., M.D., F.R.S.,

Professor of Pathological Anatomy in University College, London; Physician University College Hospital; and Assistant-Physician to the National Hospital for the Epileptic and Paralysed.

III.

DURING the examination of some specimens of sugar-cane in a sickly condition which were brought under the notice of the Scientific Committee of the Royal Horticultural Society about three years ago, I first became convinced, from personal observation, that *Bacteria* and larger Fungus-germs may be encountered within the closed cells of living plants. I found, on making thin sections of the central tissue even of young shoots, that many of the cells contained an abundance of *Bacteria*, and others a smaller number of *Torula*-like corpuscles, whilst some of the surrounding cells were quite free from either. I have since found the same kind of thing when examining the central portions of decaying tubers, and other fleshy parts of plants. Actual mycelial growths, are, moreover, to be found in various situations, to which we might pretty confidently suppose that no external germs could ever have penetrated. They have been found, for instance, in the liquid juice taken from freshly broken cocoa-nuts,* from the interior of walnuts and filberts†, and from the central portions of stone-fruits, such as plums and peaches, whilst the surrounding and external fleshy portions are quite uninjured and unaffected. Speaking of *Botrytis infestans*, which he regards as "the proximate cause of the potato murrain", the Rev. M. J. Berkeley says (*Introd. to Cryptogamic Botany*, 1857, p. 65): "The walls of the cavities of the carpels of tomatoes are often covered with the fungus, though there is no communication with the outward air; and a crop of the mould has been seen to grow in a few hours from the cut surface of a diseased potato even though the foliage itself had exhibited no trace of the parasite." Multitudes of such facts might be referred to, but the facts themselves are, I believe, admitted by all. Dr. Lionel Beale, for instance, says: "Lowly vegetable germs appear in closed cavities in the substance of dead animal and vegetable tissues. I have often seen them within vegetable cells in which not a pore could be discovered when the tissue was examined by the highest powers." And again he says: "I have detected them in the interior of the cells of animals, and in the very centre of cells with walls so thick and strong that it seems almost impossible that such soft bodies could have made their way through from the surrounding medium."‡

Nothing is easier for us than to discover such organisms within the very centre of the organs of dead animals whenever the parts begin to exhibit signs of putrefaction. They may easily be met with in the centre of a mass of brain-tissue, for instance; and M. Béchamp (*Compt. Rend.*, t. lxxvi) has also observed that most active *Bacteria* in great abundance are always to be found in the midst of a portion of liver which has been allowed to macerate in water for a day or two. When a section is made through such a mass, the cells in the very central portions are found to be swarming with moving particles and distinct *Bacteria*, though none or very few are to be found in the water in which the portion of liver is immersed. M. Estor has, moreover, found that the cells of the liver in dogs, rabbits, mice, and various kinds of birds, even immediately after death, always contain a number of actively moving particles or mere granules (*microzymæ*), which, according to M.M. Estor and Béchamp, have the power of developing into definitely formed *Bacteria*.

There can be little doubt that the granulations naturally existing in the cells of the liver and of other organs resemble those which increase under conditions of irritation, and which under a more prolonged inflammatory stimulus are apt to undergo a fatty metamorphosis leading to the disintegration of the cell. But whilst these are the changes most prone to occur during life—especially in internal parts or organs—I fully believe (with other observers) that after death, or when death is close at hand, such particles may undergo an internal change fitting them

* Dr. Sigerson writes: "The author of this paper, having opened the dense shell of a cocoa-nut, and cut through its oily albumen, both perfectly intact to all seem ing, found in the milk a web-like plant, a kind of *Achlya*." (*Monthly Microscopical Journal*, Aug. 1870, p. 11.)

† The so-called *Trichothecium roseum*, for instance.

‡ *Disease-Germs*, 1870, p. 72. Dr. Beale's mode of accounting for these facts will be subsequently referred to.

for independent life just as milk-globules are able to individualise themselves, and grow into embryo *Penicillia*. The union of two, three, or more of such granules in linear series has been watched by MM. Béchamp and Estor (*Compt. Rend.*, 1868). At first the granules form chaplet-like series, but these gradually tend to become more cylindrical, so as to produce ordinary *Bacteria* and *Vibriones*. Similar phenomena have been testified to by Signors Crivelli and Maggi (*Rendiconti di Lombardo*, 1868). These observers watched the union of vitelline granules, and saw them gradually fuse into bodies which in all respects resembled *Vibrio bacillus*, whilst these in their turn gave rise to distinct *Leptothrix* filaments. Precisely similar changes were subsequently followed out by the same observers within epithelial cells taken from the back of the tongue of a diabetic patient. In this case the granulations of the epithelial cells, by their union in linear series, formed the rudiments of the future independent organisms. Moreover, Dr. Hughes Bennett has for several years asserted that such changes habitually take place, and has always laid much stress upon them, since it was in part owing to the occurrence of phenomena of this kind that he was induced to propound his "Molecular Theory of Organisation".

On the other hand, we may watch all the stages by which epithelial cells in an apparently healthy condition become filled with the minutest granules, which subsequently develop into well formed *Bacteria*, just as particles similarly productive of *Bacteria* may be seen to appear within the substance of dying Amœbæ. If healthy looking epithelium cells from the inner side of the cheek be mounted and kept in a warm damp chamber, in the course of from twelve to twenty-four hours a multitude of isolated and motionless specks make their appearance and speedily develop, within the substance of the cell, into well formed *Bacteria*. This takes place when no notable amount of *Bacteria* exists in the surrounding fluid; and, indeed, from the mode of appearance, distribution, and development of the particles within the cell, it is obvious that, on the "germ-theory", we should have to believe that each epithelial cell which goes through this transformation is saturated with as many invisible germs of *Bacteria* as would correspond to the motionless and scattered organisms which are subsequently imbedded in its substance. In their earliest stage these units do not multiply; and before the contents of the cell fluidify, their relative positions are maintained and may be well observed.

Thus, then, we have the possibility of independent organisms arising within unhealthy or dying cells, either by means of a heterogenetic modification of some already existing particles or globules, or by a process of new birth in the fluid or semi-fluid matter of the cell. By one or other of these modes, we believe that the various Fungi and other allied organisms, which are so frequently met with in the bodies of animals as well as of plants, are capable of arising *de novo*.

In the moist mucous membranes, *Bacteria*, *Vibriones*, and *Leptothrix* are most abundant; and, more rarely, larger Fungus-germs occur, which soon develop an abundant mycelium, as where *Oidium albicans* is produced in the affection commonly known by the name of "thrush". These various organisms exist abundantly enough in almost all the mucous membranes of the body, more especially when there is some unhealthy mode of action going on in the part; and their prevalence in these situations is far more dependent upon the presence or absence of such conditions, than upon the degree of exposure of the part to the possible contaminating influence of germs derived from without. Some of those which are least exposed, are most prone to throw off the organisms already mentioned, as well as Monads and other more animalised forms.*

Fungus-germs and rudimentary mycelia are also frequently met with upon and in the superficial layers of the skin of man and the lower animals, where they represent the best known characteristics of certain familiar diseases.

Here again, as in the case of the mucous membranes and of the general parasitic diseases, there is the possibility that such growths may be occasioned by actual contact with some disseminated and all-pervading Fungus-germs. We know, indeed, that these parasitic diseases are contagious; that persons free from such maladies may become affected, provided the infecting germs fall upon suitable situations and find these in a condition favourable to their growth. Even here, however, the conjoint influence of predisposing and exciting causes of disease must come into play. So that the question is whether, in certain cases, the "predisposing" causes may not be sufficiently potent to generate the disease, without the aid of any "exciting" cause in the form of pre-existing Fungus-germs. Much evidence of a general character, in addition to the many facts and observations already alluded to, tends to favour this view—more especially in the face of the almost

insuperable difficulties which beset those who are exclusive advocates of a "germ-theory".

In reference to the above-mentioned skin-diseases in the human subject, Dr. Tilbury Fox (*Skin-Diseases of Parasitic Origin*, 1863, p. 43) calls attention to the fact of the extreme frequency with which the hair-follicles are the seats of the first manifestation of the morbid product; and says that, for the most part, "the very first spot at which any perceptible fungus can be detected is a little way inside the follicles, near the opening of the sebaceous glands." Thence the fungus-growth extends in various directions—into the follicle itself, into and upon the hair, and into the immediately adjacent portions of skin. In other cases, as in chloasma, it is the substance of the epithelial cells over the chest or abdomen which is the seat of the fungus spores and filaments.

Certain altered states of secretion from the hair-follicles, under the influence of derangement or peculiarity of constitution, seem to be most favourable to the occurrence of nearly all the cutaneous diseases which are attended by the presence of vegetable parasites. In the case of chloasma especially, we constantly see that its presence is associated with the occurrence of certain favouring conditions. Thus it is met with most frequently in persons of delicate health, who wear flannel next the skin, and who do not have recourse to sufficiently frequent ablutions. The increased moisture from retained secretion, and the increased heat, seem sufficient in certain persons to engender this disease of the skin. It is one of those affections in which we have the least reason for supposing contagion necessary, and in which we least frequently find any evidence of it. In epithelial cells affected in this manner we frequently find a number of highly refractive particles of various sizes, having a close resemblance to fatty globules and granules, though they are probably composed of a combination of fatty and albuminoid matter.* These globules and particles are frequently indistinguishable from precisely similar-looking bodies, which, by their subsequent growth, prove to be germs of fungi. It seems to me, therefore, highly probable that, under certain conditions, it is these oleo-albuminoid globules, already in existence, which actually undergo some internal changes whereby they are impelled to develop, after the fashion of ordinary milk-globules, into Fungus-germs.†

This view derives additional support from the fact that, on other occasions, Fungus-growths of various kinds have been found, in different kinds of animals, in internal cavities of the body which are wholly closed—just as they have been found within similar cavities occurring in the fruit of many plants. The Rev. M. J. Berkeley says (*loc. cit.*, p. 260): "The strongest case I have met with is the development of a yellow mould within the cerebral cavity of golden pheasants, which soon proved fatal;" whilst Dr. Murie‡ has found Fungus-growths within the abdomino-pleural membrane of a Kittiwake gull, of a great white-crested cockatoo, and of a rough-legged buzzard. Well developed Fungi have been found, moreover, within the uninjured eggs of birds and serpents by Rayer, Robin, § Cantoni, and others (*Rendiconti di Lombardo*, Nov. 1867); and this not merely just within the shell and its lining membrane, but growing from the surface of the yolk itself. It would also appear, from the experiments of Harless (*Zeitsch. für wissensch. Zool.*, 1851, t. iii, p. 308), that Fungi may be constantly engendered in unbroken eggs by exposing them in a damp oven to a continuous temperature of about 104 deg. F.

Nay, Fungi have been found even within the eye of man himself. Helmreich records the case of a gentleman who, having previously suffered from an inflammatory affection of the eyes, became rather suddenly troubled with images of a constant character floating before the left eye. On examination, a morbid product was detected floating in

* Such particles are always most abundant in the white secretion with which some of the sebaceous follicles are apt to become filled. If a specimen of this be mounted in a drop of distilled water and flattened by a covering glass, fungus-filaments will often be found to develop from all parts of it in the course of two or three days, if the specimen have been enclosed in a wide-mouthed stoppered bottle, containing a little water, so as to prevent evaporation of the drop of water in which the specimen is mounted.

† In accordance with the more generally received hypothesis, however, we should be required to believe that the Fungi in the different diseases were specifically distinct, and that spores of the different varieties were so uniformly diffused through the atmosphere as to be always ready to infect any suitable nidus. The difficulties of this hypothesis were felt by the Rev. M. J. Berkeley, who accordingly wrote in his *Introduction to Cryptogamic Botany* the following passage: "It is true that in many cases the Fungi may be of very common kind, or under disguised forms; but this is what might readily be supposed, for it is very rarely the case that such peculiar matrices as the human skin or mucous membrane should nourish Fungi absolutely peculiar to themselves. It is in such cases far more easy to believe that the common *Penicillia* or *Aspergilli*, which are notoriously indifferent about their matrix, provided the proper chemical conditions be satisfied, are the real antagonists" (p. 238). This view was afterwards ably taken up by Dr. John Lowe and Dr. Tilbury Fox, who endeavoured, with considerably success, to demonstrate the convertibility of the several forms of Fungi met with in skin-diseases, and their relationship to the forms above mentioned. This was decidedly a step in the right direction.

‡ See abstract of his communication in *Report of British Association*, 1871.

§ See his *Végétaux Parasites*, Plate II, Figs. 5 and 6, and Plate IV, Fig. 9.

* See Dr. Gros on Vaginal Animalcules, in *Bull. de la Soc. Imp. des Nat. de Moscou*, 1845, p. 426; and Mr. T. R. Lewis, in his previously quoted "Report".

the posterior chamber just in front of the lens. Soon afterwards the lower part of the cornea was punctured, with the hope that the separated foreign body might be washed out with the evacuating aqueous humour. This actually occurred, and it was then found to be a minute coniferoid-looking growth, bearing series of spores arranged in chaplet-like fashion.*

We will merely mention the fact of the existence in India of a terrible disease, known by the name of 'Madura foot', in which a fungus, closely allied to some *Mucors*, seems to extend outwards from the bones and ligaments, growing destructively through all the tissues of the part. In exhibiting a specimen at a recent meeting of the Pathological Society, Mr. Jabez Hogg also expressed his belief that this fungus takes origin from some altered tissue-elements in the foot itself, rather than from externally derived germs.

The occurrence of such growths within closed cavities not exposed to the incidence of external germs, and their not more frequent presence in the ramified air-chambers of birds (where germs might so well lodge and grow), tells just as much against the supposition of their external derivation, as does the comparatively rare and occasional presence of the growths in these cavities militate against their being developments of some out of the multitudes of minute and invisible germs with which the tissues of higher organisms are believed by a few persons to be permeated. This latter view is only countenanced by the occurrence of facts which are considered by others to be equally capable of receiving a totally different explanation.

Certain insects, moreover, are not unfrequently the seats of remarkable parasitic growths, which, although they may after a time manifest themselves externally, either take origin in, or at all events start in their development from, some internal portion of the body, and are unknown to exist in any other situations. This is the case, for instance, with the strange *Sphærea Robertsii*, which grows from a caterpillar (*Hepialus virescens*) in New Zealand; with *Sphæria entomorphiza*, which appear in pairs from the middle line of the back, at the junction of the thorax and abdomen, in many of the West Indian beetles; and with many other anomalous growths which have from time to time been met with upon members of the insect world.

[To be concluded.]

OSSIFICATION OF THE THYROID CARTILAGE, FOLLOWED BY NECROSIS: TRACHEOTOMY AND DISCHARGE OF BONE.

By W. J. FYFFE, M.D., Surgeon-Major, Assistant Professor of Medicine Army Medical School, Netley.

PRIVATE J. CURTIS, 31st Regiment, aged 38, was admitted to the Royal Victoria Hospital, Netley, at the 12th December, 1871.

Previous History.—He had been nearly twenty years in the army. He served five years at the Cape of Good Hope in the 73rd Regiment, and three years in India, including active field-operations during the mutiny in 1857-8. Having completed his first term of ten years' service, he claimed his discharge, and shortly afterwards he re-enlisted in the 31st Regiment. While serving in Portsmouth in 1866, he contracted a sore on the penis. The character of this sore is not stated in his official medical documents, and it is therefore impossible to say whether it was a true syphilitic chancre or a soft non-infecting sore. He was treated with iodide of mercury, but not to the extent of salivation; and as none of the constitutional effects of syphilis appeared for a period of four years afterwards, it is difficult to connect the symptoms which followed with a syphilitic taint.

In June 1870, he began to suffer from hoarseness, loss of voice, and dysphagia. These symptoms were followed by the formation of a large abscess in front of the neck. He described this swelling as having been as large as his closed fist, and occupying the space between the lower jaw and the middle of the thyroid cartilage, and bulging as far forward as the chin: it was extremely tender and painful. This abscess was opened in two places. One incision was made to the right of the thyroid cartilage, on a level with the pomum Adami; the other opening was made at the inner edge of the right sterno-mastoid muscle, near the sternal end of the clavicle. A large quantity of pus was discharged, the tumour rapidly disappeared, and the incisions healed up; but the opening of the abscess was not attended with any alleviation of the symptoms of aphonia or dysphagia.

His condition on arrival at Netley was as follows. He had great

dyspnoea, evidently produced by mechanical obstruction to inspiration; each effort to inspire was accompanied by a harsh stridulous noise; the voice was reduced to a whisper; he was unable to lie down; the countenance was anxious, but there was no abnormal rapidity of pulse or increase of temperature. On examining the neck externally, there appeared some thickening of the skin over the thyroid and cricoid cartilages; the space between the thyroid and cricoid was diminished, and the crico-thyroid membrane shortened. On placing the finger on the cicatrix of the upper incision already alluded to, a distinct depression in the right ala of the thyroid could be felt, indicating loss of substance at that point; but no pain or tenderness on pressure could be detected over any part of the cartilaginous skeleton of the larynx.

On examination with the laryngoscope, the ary-epiglottidean folds were found swollen; the left one was displaced and pushed over to the right side. The vocal cords appeared healthy; but immediately below the left vocal cord a whitish tumour appeared protruding into the cavity below the cords occupying the posterior third of the opening, and partially blocking up the rima.

As the case was urgent, everything was prepared for the operation of tracheotomy, in case any alarming paroxysm of dyspnoea should supervene. A violent paroxysm having set in at 1 o'clock next morning, the trachea was opened by Surgeon-Major Mackinnon, C.B. The operation was followed by instant relief to the patient; he breathed quietly and easily through the tube, and there was a cessation at once of all dyspnoea.

For several days the patient progressed most satisfactorily; but, on the morning of the seventh day after the operation of tracheotomy was performed, he was suddenly seized with a violent fit of coughing, which lasted for about two minutes, and during which he brought up into the mouth a piece of bone; the paroxysm of coughing ceased immediately afterwards. This fragment was very foetid, light, and friable, and had somewhat the conformation of the inferior and posterior angle of one of the alae of the thyroid cartilage. It was at first conjectured that it might possibly have been a piece of mutton-bone, which might have been accidentally lodged in the larynx when the laryngeal symptoms first appeared. But the patient positively declared that no such accident could have happened without his cognisance; and the shape and appearance of the fragment gave no support to this conjecture. The question was, from what bony structure could this have been detached. The os hyoides was examined and found intact. The spine was examined from behind, and no tenderness could be discovered indicative of disease of any of the cervical vertebræ; and, in addition to this, it was unlikely that a necrosed portion of the body of a vertebra would find its way into the larynx, on account of the strength of the structures lying in front of the spinal column. Besides, the sequestrum did not present the large cancellated structure characteristic of the body of a vertebra. There appeared, therefore, to be but one solution of the question—namely, that ossification of some portion of the cartilaginous frame of the larynx had taken place, followed by necrosis, and resulting in the large abscess already alluded to. Separation of this sequestrum most probably took place then, which, failing to find its way through the incisions in the neck, became fixed in the larynx, and produced the urgent symptoms which rendered tracheotomy necessary.

The patient is now convalescent; the opening in the trachea is completely healed up; he speaks no longer in a whisper, but with his natural voice; the swelling of the ary-epiglottidean folds has disappeared; the rima is, however, slightly narrowed, and there is a little stridor on deep inspiration.

It is possible, although I think very doubtful, that the pathological conditions existing in this case may have had a syphilitic origin. It appears to me more probable that the affection is one more allied to those described by Dr. Wilks at page 205 of his work on Pathology; where he remarks "that, although many chronic affections of the larynx have a syphilitic origin, there is, however, an affection independent of syphilitic taint attended with pain and swelling about the larynx, followed by abscess either within or without; very commonly one ala of the thyroid becomes converted into bone, dies and exfoliates, originating probably first in perichondritis, then ossification, and finally necrosis."

There can be no question that in this case the patient, on admission to Netley, was in imminent danger, and that the promptitude with which the operation of tracheotomy was performed by Mr. Mackinnon saved his life.

Note.—Since the above account of this case was written three weeks ago, the patient has been again attacked with urgent laryngeal symptoms, and it was considered desirable to open the trachea a second time, which has been done to his great relief. Most probably there is still another sequestrum making its way into the larynx, although nothing can be seen by the laryngoscope to indicate its presence.

* See Robin's *Végétaux Parasites*, 1853, p. 370. He adds: "Après l'opération le malade se trouva bien, et continua ses occupations sans gêne." Robin also mentions a case (p. 364) in which a larger though somewhat similar growth was met with by M. Gubler growing beneath the skin of the back of the hand of a young man suffering from a gun-shot wound. Several white bleb-like elevations of the skin were produced, which were found to have been occasioned by the growth in question.