importance to my relation of facts; but, as sceptics of the kind must be rare, I proceed to narrate very briefly three cases bearing on the question at issue.

CASE I.—In 1856, I attended a lady in her seventh confinement. Labour formerly had been always quick and easy: so it was in this case up to the completion of the first stage. The os was well dilated, the membranes were ruptured, and the hairy scalp presented. There the matter ended, however. I had prophesied a speedy termination, and, not being then so well acquainted with the difference between meddling and muddling as I now hope I am, I waited till tired nature could endure no longer, and went home to bed. Next day, and the next, and the next, matters were unchanged. The patient got up and moved about her room, although it seemed as if one good expulsive pain might complete the process. Not until the tenth day was there any recurrence of pain, when a living child was born, before even a messenger could be despatched for me. Every other point was normal.

CASE II was of a precisely similar character. It occurred six years later, and the patient was, oddly enough, also in her seventh labour. Here I allowed matters to go on naturally with my eyes open, and as a matter of scientific curiosity. The result was absolutely similar, with the exception that in this case twelve days intervened between the full completion of the first stage and the sudden delivery of a living child. I have not met with another case where nature has attempted this freak, and if I did, I should probably not fall in with her dilatory proceedings.

CASE III was of quite a different kind. In June 1868, I was asked to see a lady in the end of the seventh month of her first pregnancy. She suffered so much from sickness and consequent exhaustion, that I felt compelled to recommend the induction of premature labour. This I endeavoured to bring on by the introduction of a catheter between the uterus and membranes. In attempting to do so, I accidentally ruptured the membranes a few inches above the os uteri. A very large quantity of fluid escaped, and the uterus, which had formerly been large for the period, collapsed and hardened. From that time the sickness ceased, and, though there was frequent dribbling of fluid, there were no other signs of labour. The patient went on for six weeks, and was then easily and rather quickly delivered of a living child, which, however, survived only a few hours. From its appearance, I fancied that the date of pregnancy had been somewhat miscalculated. From the time of the rupture of the membranes and cessation of sickness, I made no further attempt to follow up the induction of labour.

These three cases alone seem to me to settle positively and affirmatively the question at issue. They prove indisputably that, after the escape of all or nearly all the liquor amnii, and the actual uncovering of the lower segment of the foetus, labour may be delayed for at least twelve days without injury to mother or child. Can any one affirm that twelve days is the physiological limit? They also prove that the membranes may be even artificially ruptured, a very large quantity of fluid escape, and labour not supervene for six weeks, the phenomena being highly advantageous to the mother, and apparently not detrimental to the child. No one can dispute that the general law is otherwise, and the rarity of such exceptions as I have narrated only tends to further its proof.

THE EFFECTS OF A BELLADONNA PLASTER.

By J. BOWER HARRISON, M.D., F.R.C.S., M.R.C.P.,

Corresponding Member of the Epidemiological Society of London, etc.,
Higher Broughton, Manchester.

BELLADONNA plasters are found in so many instances to be the means of affording relief in painful nervous affections of a local kind, that they are doubtless very largely employed. It will be interesting, then, to relate a case in which the poisonous effects of belladonna were fairly attributable to the use of one of these plasters.

On November 17th, 1871, I was called to see a lady who was taken very ill on her return from Bury, a town in the neighbourhood of Manchester. She was in a state of great confusion of mind, and appeared in many respects like a person under the effect of intoxicating liquor. Her speech was altered; and her ideas were so far distracted that she had the most strange fancies, imagining that the pictures on the walls were moving, and that a medallion profile which hung before her was continually opening its mouth. As much scarlet fever was prevailing in the neighbourhood, I naturally supposed that she was about to suffer from the same complaint, and gave my opinion accordingly, recommending that an able nurse should at once be procured. It was

stated that, when the lady reached Manchester, she felt so confused that she was scarcely able to direct the cabman where to drive her, and did not, in fact, know at what place she had arrived.

On visiting her the next morning, she complained of a soreness and dryness of the throat; and I noticed that the fauces were unnaturally red. A rash subsequently showed itself on the body; so that I concluded that she was suffering from the scarlet fever.

Still there were some circumstances, which have now escaped my recollection, that made me rather hesitate in my opinion; and I felt somewhat embarrassed in making up my mind as to the nature of the case. In the course of inquiry, however, it came out that the lady some little time previously had put a large belladonna plaster over her back, and that this plaster had occasioned so much irritation that she had been obliged to remove it. The removal, nevertheless, had not been effected without considerable abrasion of the cuticle. Notwithstanding this lacerated state of the skin, it appears that the plaster had been reapplied before her journey to Bury. A ray of light now seemed to dawn upon me, and I began to suspect that the case was one of belladonna-poisoning. In corroboration of this view, I found that the pupils of the eyes were inordinately dilated. I then directed that every trace of belladonna should be carefully removed from the skin.

I need not go into further details. Suffice it to say, that in a short time she began, though slowly, to recover, without the rash being followed by any desquamation of the cuticle, or any affection of the kidneys.

The daughter of my patient, an intelligent young lady, at my request drew up the following brief narrative.

"Dear Dr. Harrison,—My mother was comparatively well on the morning of Thursday, November 16th, and went out of town to visit some friends. She complained, it is true, during the day, of some giddiness; but this was not so great as to be noticed by her friends. On her return home, however, she became suddenly confused in her mind—so much so that she was unable to tell where she was on reaching Manchester, and with considerable difficulty was capable of directing a cabman to drive her home. On reaching home, her family noticed a complete change in her manner. Her speech was affected, and she was unable to walk steadily. She fancied the pictures on the walls were moving and speaking, and that people were passing in and out of the room in the most singular manner. She thought that she saw insects on the bed, and sometimes butterflies with beautiful wings. She afterwards complained of soreness and dryness of the throat; and a rash showed itself on the body. The delirium continued, with the exception of a lucid interval of about two hours, from Thursday until Saturday morning, when it suddenly disappeared. The rash also soon declined, together with the soreness of her throat. At this period, you began to suspect that a belladonna plaster, applied to an abraded surface, had been the cause of her malady; and, after its complete removal, the health slowly recovered. My mother being, however, of a delicate constitution, was a considerable time before she regained her usual health.

"Believe me, dear Dr. H., yours sincerely,

"JANE —."

In the *BRITISH MEDICAL JOURNAL* for Jan. 27th, 1872, there is a lecture on Disease of the Valves of the Heart by Dr. George Johnson, in which he remarks (p. 93): "A belladonna plaster worn over the heart often proves an efficient local anodyne. Belladonna, remember, cannot be applied to an *abraded surface* without risk of absorption and alarming symptoms of poisoning." In the *ASSOCIATION MEDICAL JOURNAL* (Sept. 16th, 1853, p. 818) will be found the report of a case in which all the symptoms of poisoning by belladonna arose from the application of a weak solution of atropia in water to the conjunctiva.

Some interesting cases of belladonna poisoning, from its external application, are related by Mr. Wm. F. Morgan in the *BRITISH MEDICAL JOURNAL* for December 1st, 1866. "I was sent for," he says, "in haste, a few years ago, to see a gentleman who was supposed to have impending some serious affection of the brain. I found him in a state of mental confusion bordering on delirium, with his lower extremities semi-paralysed, his pupils widely dilated, and vision much impaired. The pulse was very frequent; the skin hot and dry, and covered with a *scarlet rash*, which occasioned intolerable itching. The rash extended throughout the fauces and throat, accompanied with a feeling of great dryness and constriction and intense thirst. I was told he had been suffering from lumbago, and of his own accord applied a large belladonna plaster to his loins. This being removed (it was at least a foot square), he speedily recovered."

"The other case was the mother of a medical friend, who had some painful affection of her leg, for which he strapped it with belladonna plaster. He came to me soon after, in much alarm, requesting me to see his mother immediately, as he feared she was threatened with apoplexy. Her symptoms were those of belladonna poisoning, in a less

degree than in the former case, and without the characteristic *scarlet rash*; and, on removing the plaster, she was soon all right."

The same journal (November 24th, 1866) relates from the *London Hospital Reports* two cases of poisoning by the external application of belladonna. "The first was that of a nobleman, for whom a liniment containing two drachms of liquor belladonnae in two ounces of soap liniment had been ordered. After using it, he hastily summoned his physician, who found him with widely dilated pupils, cerebral excitement, and rapid pulse. Another liniment, without the belladonna, was prescribed, and the doctor left. The next day, he was again hastily summoned, and found his patient with a solicitor, three keepers, and a 'mad doctor', who had already signed a certificate of insanity. It turned out that the liniment containing belladonna had, by mistake, been applied a second time; and hence a return of the cerebral excitement, rather exaggerated. Dr. Brown explained that the patient's state was attributable to the belladonna, and that the symptoms would speedily disappear, as indeed they did."

Another case is related by Dr. Fraser, of a servant girl who was taken to the London Hospital by her mistress, "who was afraid the girl was going out of her mind. The girl was extremely restless; would wander about the room; stand first on one foot, and then on the other; but was quiet for a moment when spoken to, and answered a question with an effort, generally breaking off in the middle, as if she had forgotten what she had to say. It was noticed at once that her pupils were widely dilated and fixed. It was subsequently discovered that she had pains in her breasts for some days, and had gone to a chemist, who had given her a lotion composed of half a drachm of extract of belladonna in an ounce of water, with directions to apply it on cotton-wool covered with oiled silk. She applied some in the evening, and again in the morning; and in the afternoon the symptoms appeared which induced her mistress to take her to the hospital. The girl expressly denied having swallowed any of the lotion. She recovered slowly." (*Chemical News*.)

A case somewhat similar, but of a milder kind, occurred in my own practice. A young woman recently confined, being unable to suckle her child, applied to me for some means of removing her milk. I prescribed a small quantity of extract of belladonna dissolved in olive-oil to be applied to the breasts. A day or two afterwards, she complained of dryness and soreness of the throat—of confusion of mind and vision—as well as of an irritable rash which spread itself over the body. I immediately suspected that the belladonna was the occasion of these symptoms, and on its removal they rapidly disappeared.

From the foregoing observations, then, it may be well to remember that even the external application of belladonna is occasionally productive of mischievous effects.

ANÆMIA OF THE OPTIC NERVES TREATED BY GALVANISM.

By R. J. PYE-SMITH,
Late Dresser in the Eye-Wards of Guy's Hospital.

VERY little success has hitherto attended the treatment of amblyopia and amaurosis associated with anæmia of the optic nerve (white optic disc). I have, therefore, drawn up the following account of a few cases treated by galvanism, in the hope that its publication may lead to further experiment on the subject.

Bearing in view the experiments of Professor Helmholtz on the healthy eye,* Mr. Bader suggested the application of galvanism to the optic nerve in cases of white optic disc. And for the purpose of testing its therapeutic value, the following cases have been received into Guy's Hospital.

Before giving a short account of each case, I had better describe the mode in which the galvanism was applied, and the effects it produced on the healthy eye.† A battery of Daniell's elements was used, the electrodes terminating in small sponges soaked in warm water. The negative electrode being placed on the skin behind the ear, the positive electrode was applied to the closed eyelids of the same side, or to the brow. On closing the circuit with ten elements, the healthy eye perceived a round central illumination, generally described as about the size of a sixpence, or of the tip of the finger, irregularly circular, and of a blue tint. The impression lasted about five seconds, and then gradually faded away. On interrupting the circuit, a momentary flash was perceived over the entire field.‡ These effects were produced

when the eyes were open, though more marked when they were shut. The modifications observed in the patients with white optic discs will be mentioned in the account of each case. The number of elements employed was determined by the flashes perceived, the smallest number being used that produced in each case any illumination, and where no flashes were perceived an unbearable amount of burning at the electrodes, or the occurrence of giddiness, served to limit the number.

Ophthalmoscopic examination during the passage of the current showed in all cases the veins of the retina to be congested after the first second (during which they sometimes appeared to be partially emptied), and to remain so till the interruption of the circuit, when they gradually resumed their natural fulness. This effect was observed even when no flash was perceived by the patient.

The patients were galvanised every Tuesday, Thursday, and Saturday. The current was usually passed six times at each sitting, for eight or ten seconds each time, with intermissions of an equal period.

The following cases are all of men, Mr. Kiddle having undertaken the galvanism of the women admitted with the same disease.

CASE I.—J. I., aged 42, a wheelwright, was admitted December 12th, 1871. His general health was good, except for an attack of quotidian ague twelve years ago. He had been in the habit of smoking about two ounces of shag per week. His sight began to fail nearly five years ago. He went to the Ophthalmic Hospital, and attended as an out-patient for three years, taking medicine daily. The sight continued to fail during this period, but the patient was able to keep at work till the end of that time, and till the last few months of it, to read. He frequently had severe occipital headache. He came to Guy's Hospital as out-patient nearly two years ago, when both optic discs were found to be white; he had attended ever since, giving up smoking, and taking one-third of a grain of pilula opii three times a day. His sight had continued to become worse. During the last year he had felt ill and nervous. Since his sight began to fail, he had had about half a dozen attacks of frequent shivering followed by vomiting, lasting for a day or two, and then passing off. On admission, he was very tremulous. He could just distinguish windows with each eye, but could not count the panes. He considered the right his best eye. He stated that he saw best in a dull light, and that he was able occasionally to see the time by a large clock for a minute or two, and then the sight became dim again. The tension of the right eye was (?) slightly above the normal. The pupils were dilated and sluggish. On ophthalmoscopic examination, both eyes were found hypermetropic. The right optic disc was well defined, bluish-white, with very white centre; its vessels were slightly displaced to the nasal side; the arteries were small and few. The left optic disc was well defined, bluish-white; the arteries were very small and few; the veins few and half empty. The retina was rather anæmic. There was small floating opacity in the vitreous body. Both eyes were galvanised twenty-three times—i.e., on twenty-three different days; each day, as before stated, the current was passed about six times through each optic nerve. A battery of twenty elements was used. A flash was usually perceived on closing the circuit, and a "dark flash" on interrupting it. Occasionally slight headache was produced, but the trembling disappeared in a week or so. After the fourth application, the patient was able to see the seats in the grounds; after the seventh, he could count the panes of a window upwards, but not from side to side; after the seventeenth application, he was able to read "Operation Room," painted in letters about three inches high over a door in the ward. He could now always see the time by a large white-faced clock, and could see his way about the grounds much better than at first. On January 18th, he was seized with one of his attacks of vomiting, shivering, and loss of appetite. The sight was a good deal impaired for several days, but the galvanism was continued as he was very anxious for it. His sight was examined again on January 31st. He could now see the paths in the grounds, people as he met them, a white stone sundial out of doors, the time by the clock, and could count the panes of a window in both directions with either eye. Ophthalmoscopic examination showed no marked change.

CASE II.—W. C., aged 30, a soldier, was admitted December 15th, 1871. His general health was good. He had served in India and China. He had been accustomed to smoke about an ounce and a half

side of the field, and was accompanied by a less distinct and somewhat larger illumination in the centre of the field. These may perhaps represent the optic disc (well defined and to the temporal side) and the region of the yellow spot (central). That they are not due to the form of the electrodes, is shown by the fact that small V-shaped electrodes produced the same circular illumination. It may be mentioned that this illumination was not perceived by patients whose eyes had been excised; though one of three such patients experimented upon did perceive the flash over the entire field on interrupting the circuit, and all three of them perceived distinctly the flash over the entire field produced in healthy eyes on interrupting the circuit when passed in the opposite direction (i.e., the negative electrode being on the eyelids). In one of these cases the eye had been completely excised seven years before.

* Helmholtz on *Physiological Optics*.

† Both the method of application and the effects on the healthy eye were substantially those described by Professor Helmholtz.

‡ In my own case, the defined illumination was placed somewhat to the temporal