the same process occurs under the skin, as in one of the two cases, it goes by the name of phlegmonous lymphangitis. I said the passage of the puerperal ichor through the larger lymphatics caused various degrees of inflammation; and the error of Cruveilhier and the pathologists of his time was to suppose that puerperal fever could not be attributed to the lymphatics unless they contained pus. In my paper on lymphangitis in pelvic pathology, I have cited cases of puerperal fever in which the enlarged lymphatics were found to contain not pus, but a fetid brown fluid, similar to that found in the womb, containing a bit of putrid placenta. I have, moreover, shown that when filtered putrid lochial fluid is injected into the womb of recently delivered animals, this gives rise in all to the same typhoid symptoms; but, on opening the animals, it is found that sometimes the lymphatics contain pus, and at others a fetid sanious fluid like that found in the womb.

SPINA BIFIDA.

By E. NOBLE SMITH, L.R.C.P.Lond.

MR. BURTON'S communication upon this subject induces me to again express a doubt that a case of spina bifida, in which the cord communicates with the sac (*i.e.*, adheres extensively to the back of the sac), can be cured by an operation which causes the sac to contract and shrivel up. The success attending Dr. Morton's treatment seems, in a very great measure, due to the modifications which he has introduced. This fact I fully recognise, but I trust I may be allowed to express a doubt as to the spinal cord being connected with the sac in the manner shown in the diagrams given below, or in the preparation, a drawing of which accompanies Sir Astley Cooper's cases reported in the *Medico-Chirurgical Transactions*.

In answer to a former letter of mine, Dr. Morton observes (*BRITISH MEDICAL JOURNAL*, November 21st, 1874) that, as all his cases have been successful, he had not had an opportunity of making a *post mortem* examination of an injected case. Five consecutive cases treated according to Dr. Morton's plan recovered. A sixth is now reported, which has failed.

I now venture to analyse these five successful cases.

CASE I (*BRITISH MEDICAL JOURNAL*, April 6th, 1872). *Spina bifida in a child two months old.*—"Its covering was thin, so that it was quite translucent; and at the child's birth it was about half its present size; moderate pressure upon it did not cause much inconvenience or suffering; and the child had never had convulsions, and seemed in all other respects quite healthy." It was cured by Dr. Morton's injection. In this case, there is no evidence of the cord communicating.

CASE II (*BRITISH MEDICAL JOURNAL*, June 15th, 1872). *Spina bifida in a child fourteen days old.*—"Dr. Morton's assistant reports, that 'the tumour was as large as a middle-sized orange, and cylindrical in shape. In some parts it was reddish, and at others bluish in tint. It was semi-transparent, and somewhat wrinkled on the surface. It became tense when the child cried. On looking through it, several striæ were seen passing over its internal surface.' There is no proof that these striæ were nervous, and, as the child does not seem to have suffered at all from convulsions or paralysis, I think it is highly probable that they were only inflammatory adhesions. This opinion is strengthened by the fact that the tumour 'at birth was small, but had increased in size', and that 'some time ago it showed signs of ulceration, but these healed up, leaving thickened cicatrices of different colours'. It was cured by Dr. Morton's injection.

CASE III (*BRITISH MEDICAL JOURNAL*, April 26th, 1873).—Dr. Watt reports a case in which the spina bifida was a sessile semi-transparent flesh-coloured fluctuating tumour, as large as an ordinary sized hen's egg. It was extremely sensitive, and, "when viewed by transmitted light, cords were seen passing near the surface". There is much more evidence in this case in favour of the cord communicating to some extent with the sac, for the child moaned almost constantly. "Its legs dangled, and no voluntary attempt was made to move them", and "the sphincters seemed very weak". This tumour was injected with Dr. Morton's fluid, and the tumour contracted into "a cake of firm condensed tissue", but "the legs never moved together, but, when the soles were tickled, one or other leg was drawn up a short distance". There is no absolute proof, however, that the cord communicated with the sac, nor does there seem to have been a complete cure.

CASE IV (*BRITISH MEDICAL JOURNAL*, January 31st, 1874). Reported by Dr. Watt.—The child made an excellent recovery after injection with Dr. Morton's fluid. The tumour "was very sensitive to the touch, became tense when the child cried, and palpation discovered a perfectly fluid condition", which is equivalent to stating that the cord did not communicate with the sac.

CASE V (*BRITISH MEDICAL JOURNAL*, October 24th, 1874). Re-

ported by Dr. Morton. In this case, the tumour was situated in the cervical region, and no evidence is given of the spinal cord being involved in the sac.

The accompanying cuts represent two preparations in the Museum of the College of Surgeons, of which Mr. Flower kindly allowed me to

make drawings, and I cannot understand how such cases could be cured by making the tumour contract and shrivel up.

ON JABORANDI.

By SYDNEY RINGER, M.D., Professor of Materia Medica and Therapeutics in University College, London; and W. MURRELL, L.R.C.P.

SINCE a paper published in conjunction with Mr. Gould, the present writers have made ninety-four additional observations on the effects of jaborandi. The results, whilst confirming the previous statements, add a little to our knowledge of the action of this interesting drug. We gave the medicine sixty-eight times to patients, who took it at their own homes at bedtime. In fifty-nine instances, both perspiration and salivation occurred; in five, perspiration without salivation; in four, salivation without perspiration; thirteen complained of pain over the pubes, often very severe, with a strong desire to pass urine; thirty-one experienced pain in the head, often accompanied with giddiness and dulness.

On seventeen occasions, we gave the drug at nine o'clock in the morning, in a warm room, before a large fire. Of these cases, both perspiration and salivation occurred in eleven; in two, perspiration without salivation; in one, salivation without perspiration; three experienced pain over the pubes; six felt severe pain in the head; six nausea or sickness; and, in one instance, the drug caused a good deal of drowsiness; in four cases, the sight became affected; and finally, in three cases, the jaborandi produced no effect. On nine occasions, we gave the drug to patients who immediately left the hospital and returned home, during the late severe weather. Both perspiration and salivation occurred in eight cases; in three, pain over the pubes; in six, severe pain in the head; in four, nausea; in six, much sleepiness; and, in one instance, the drug produced no effect.

In by far the greater number of observations, both perspiration and salivation were profuse; but sometimes the perspiration or the salivation or both were only slight. The observations were repeated several times on certain individuals, and we found that the effects of the drug varied considerably, not only on different people, but even in the same person. Some yield with difficulty to the drug, and require a large dose, even ninety grains of the leaves. In others, whilst exciting considerable perspiration, it affects the salivary gland but little, or not at all; on the other hand, in some it excites profuse salivation, but scarcely affects the perspiration. In respect to its influence on the brain, the sight, the urinary organs, the effects of jaborandi are even still more uncertain.

As might be anticipated, cold, exposure, and food influence the effects of the drug. Thus it acts less freely on the skin of persons dressed or exposed to cold; moreover, when administered on a full stomach, the drug is more slowly absorbed, and its effects are less marked. In experimenting, therefore, on persons dressed or exposed to the cold, we were obliged to give a larger dose to get the desired effects. The pain over the pubes was often very severe, especially on two occasions, when it was felt across the navel and passed downwards along the penis, and down the urethra. With few exceptions, the pain subsided immediately on passing water. The headache was generally referred to the forehead over the eyes, though sometimes it involved the whole head, and on one occasion was limited to the left parietal region. The character of the pain was generally dull; most persons calling it a "weight over the eyes". In one case, it was paroxysmal. When the drug was given in the daytime, the pain lasted from half an hour to several hours. Giddiness and a stupid feeling were often associated with the headache. This effect on the head, we believe, has before not been noticed. On