wards during micturition at the umbilicus, welling up into the umbilical cup, filling it, and running over in such quantity that the parents had no reason to entertain any doubt as to the nature of the fluid. The urine was passed with difficulty by the urethra, and fell down from the orifice. No tumour was visible at the umbilicus; the prepuce was long, contracted, and adherent to the glans. Tincture of perchloride of iron had been applied to the umbilical aperture to produce contraction, but without avail. I operated for the phimosis according to Dr. Gordon's plan of slitting up the prepuce, as it seems to be very simple and efficacious. A nevus-needle was passed between the glans and the prepuce, and out through the prepuce at the corona. The prepuce having been grasped longitudinally beneath the needle by a pair of strong forceps, as much of it as was thus embraced was cut away by running a knife through the tissues along the lower border of the forceps. The mucous membrane was separated with difficulty from the glans, then cut freely, and its edges fixed to the skin by two sutures. Dr. Redfern saw the child lately, and has written me to say that, since the operation, the mother has been "quite pleased to see the urine projected to some distance from the body in the natural manner. Very little urine has come through the umbilicus; but at one period some blood oozed away," which was readily stopped by the application of tincture of perchloride of iron. "When I last saw it," he continues to say, "there was scarcely any appearance of ulcer or opening of any kind."

In some instances where the cure after circumcision has not proved complete, it might be necessary, in addition, to operate directly on the urachus according to the plans already mentioned; but neither in patent urachus nor in umbilical fecal fistula can we reasonably expect a cure, so long as there is an obstacle to the passage of urine or fæces by the ordinary route.

PROLAPSE OF OVARIAN CYST.

By A. W. STOCKS, M.R.C.S.,
Surgeon to the Salford Royal Hospital.

E. I., AGED 45, a fustian-cutter, pallid, slightly built, married, had three children; she was last confined about twenty-two years ago. She menstruated regularly up to two years ago, irregularly till eight months ago, and not at all since. About eleven years back, a tumour about the size of a walnut appeared at the anus when getting out of bed, becoming larger on exertion. It was accompanied by faintness and uneasiness, especially when sitting down. She was always constive, and could neither micturate nor defæcate unless she replaced the tumour manually. She had lately experienced difficulty in coition. The lump had increased in size during the last ten months, and come down always while at work, producing a good deal of pain in the hypogastrium, more particularly when she was in the upright position. She felt best when lying down. The prolapse was large, being about the size of a small cocoa-nut.

March 15th, 1872, 12 noon. When she was placed upon the operating-table, and under the influence of chloroform, for the purpose of having the simple prolapsus ani, as it was supposed to be, relieved by operation, defæcation took place, the contents of the bowel being expelled in such a peculiar manner as to lead to the further and more precise examination of the tumour. It was then discovered that the orifice of the bowel, instead of being at the most dependent part, was on the posterior aspect, and about half way between the edge of the anus and the lowest part of the prolapse. On introducing the finger into the rectum, a large mass was found to occupy the anterior fold of the prolapsed bowel, of globular shape, and capable of being encircled easily at its base. *Per vaginam*, the cervix uteri was found tilted forward and to the right side. The uterine sound could only be passed three-quarters of an inch, and a finger passed into the rectum could be easily approximated to the one in the vagina over the tumour, clearly showing an absence of continuity between the uterus above and the mass below. Moreover, on rubbing the tips of the fingers together, a hardened cord could be felt slipping between them. The conclusion, therefore at once arrived at was that this cord was the Fallopian tube, and that the tumour was a small ovarian cyst, which had fallen through Douglas's pouch, became entangled with the prolapsed rectum and protruded through the anus, dragging the uterus itself out of its normal position. It was evident that to give permanent relief it was necessary that this tumour should be removed. Accordingly, an incision was made in the anterior aspect of the prolapse parallel to the axis of the bowel, and, after breaking down some slight adhesions posteriorly, a small ovarian cyst was easily turned out. The pedicle was divided, after being secured by a strong hempen ligature, the end

of which was left hanging out of the wound, and the wound was drawn together by an uninterrupted suture. There was considerable hæmorrhage, and the flaccid bowel was left outside the anus.

At 5 P.M., she had slight vomiting, the skin was cool, hæmorrhage slight, and the prolapsed bowel was swollen. She expressed a feeling of relief and lightness about the rectum, and it was noticed that the ligature had slipped off the pedicle and lay outside the wound. The catheter was passed, and she took an opiate.

March 16th. The countenance was good, pulse 120, temperature 98.2 degs. She had vomited frequently up to 7 A.M.; most probably from the chloroform. There was slight pain in the hypogastrium and right iliac fossa.

March 17th. There was no tenderness. The prolapse was supported by a T-bandage. She went on well until the 24th, when she passed a solid motion involuntarily, as she did daily for some time.

April 5th. The tumour was much smaller.

April 9th. The prolapse was easily returnable.

April 16th. She walked about and felt more comfortable than she had done for a long while back.

April 23rd. The protruded portion of the bowel was about the size of a walnut. Defæcation was fairly under her control, and she could draw the prolapse back without manual assistance.

1875, July 15th. The prolapse still existed, but was accompanied by no discomfort in, or obstruction to, either the act of defæcation or micturition.

The tumour (the dried preparation of which is in my possession) was an unilocular ovarian cyst, and contained about $5\frac{1}{2}$ ounces of brown, slightly viscid fluid, specific gravity 1.025. The ovary and fimbriated Fallopian tube were attached to it, and on it were found marks of slight adhesions.

This extraordinary case is of perhaps more interest from its singular pathological character, rather than from any peculiar practical surgical value. The numerous factors, each of which must have furnished its own particular but necessary influence in bringing about such an unusual result as a prolapse of an ovarian tumour through the anus, will, in all probability, be long looked for ere a similar instance will be recorded.

POISONING BY TOBACCO.

By E. R. MORGAN, M.R.C.S., L.S.A., Medical Officer of Health to the Neath Union.

THE history of the following cases of poisoning by tobacco is, I venture to think, of sufficient interest to merit a place in the JOURNAL. The manner in which the poisoning was effected is, as far as my knowledge goes, quite unique, not that I mean to say poisoning by tobacco is unknown; on the contrary, I know many cases are on record. By some means or other, some tobacco got into a well, the water from which was drunk by three persons, and in each very severe symptoms of poisoning were set up, causing death in one instance; how the tobacco was discovered, will be described more fully hereafter.

It happened in a large village called Ystalyfera in Glamorganshire, which at the time was in the district over which I had jurisdiction as medical officer of health; hence I took great interest in the cases, and investigated the facts upon the spot. The other details were kindly furnished me by Dr. Thomas.

On Wednesday, March 10th, 1875, Dr. Thomas was called to visit a man, aged 55, living in a row of five houses, called "Cwmtawe". He found him lying helpless in bed with very confused ideas, unable to swallow, with complete loss of power over his lower extremities, hands trembling, and skin clammy and cold; heart's action very feeble; pulse hardly perceptible; he did not vomit, but had constant nausea: it ought here to be stated that he had been a habitual smoker for many years. With such symptoms, Dr. Thomas's suspicions were immediately aroused; and forthwith he made minute examinations in the garden for any vegetable poisons, and as carefully inquired of the wife of the patient whether she had procured any drug recently from some itinerant quack or chemist; but from these sources nothing could be elicited to account for the symptoms, and so for the time his mind was obliged to rest satisfied that the case, in all probability, was one of vegetable poisoning, but what that poison was he was then unable to say.

The same morning a child, a nephew, aged four years, was taken ill, but not with severe symptoms. However, on the next day, the doctor's attention was drawn to him. The lad was very sick, prostrated, with severe pains in the bowels; their action was arrested, necessitating the use of enemata; throat parched; pupils dilated, fixed and staring; pulse scarcely perceptible at the wrist; head thrown back, almost