was then put on infusion of jaborandi (one drachm to six ounces), and this treatment was continued for forty-four days. The dose was from time to time increased till he took several times daily a dose enough to produce in healthy people very profuse perspiration and salivation. As will be seen, the jaborandi had no effect on the quantity of urine; but his mouth was moister and more comfortable, and, though the jaborandi never produced copious perspiration, it kept the skin comfortable, and, after exercise or eating, or hot drinks, he perspired rather freely. The sweating of the feet soon ceased after taking jaborandi. The medicine did not make him spit. After it had been clearly proved that the jaborandi was useless, he was put on ergot for the third time with considerable success. The results of the treatment will be seen in the following table.

Date.	Quantity of urine in litres.	Treatment.	Date.	Quantity of urine in litres.*	Treatment.
Aug. 12 ..	9.0600		Aug. 30 ..	10.0600	
" 13 ..	10.0600		" 31 ..	10.0500	Four drachms every 3 hours.
" 14 ..	8.0500		Sept. 1 ..	11 ..	Five drachms every 3 hours.
" 15 ..	11.10		" 2 ..	10.0500	Six drachms every 3 hours.
" 16 ..	10.0250		" 3 ..	11 ..	Five drachms every 3 hours.
" 17 ..	10.0100		" 4 ..	9.0600	
" 18 ..	10.0100		" 5 ..	11.20	
" 19 ..	11.600		" 6 ..	10 ..	One ounce every 3 hours.
" 20 ..	11 ..		" 7 ..	10 ..	
" 21 ..	10.0100	Two drachms of in- fusion of jabor- andi every 4 hrs.	" 8 ..	11 ..	
" 22 ..	11 ..		" 9 ..	10.0500	
" 23 ..	10.0600	To be given every 3 hours.	" 10 ..	11 ..	
" 24 ..	10.0250		" 11 ..	11 ..	
" 25 ..	11 ..		" 12 ..	10.0500	
" 26 ..	10.0600	Three drachms every 3 hours.	" 13 ..	11.0500	One ounce every 2 hours.
" 27 ..	10.0600				
" 28 ..	11.0250				
" 29 ..	11.0600				

From the 14th to the 30th inclusive, the amount varied between 10.0500 and 9.0250

Octr. 1 ..	10 ..	One ounce and a half every 2 hrs.	Octr. 16 ..	6.0300	
" 2 ..	11 ..		" 17 ..	6.0600	
" 3 ..	11 ..		" 18 ..	6.0500	
" 4 ..	11 ..		" 19 ..	5.0800	
" 5 ..	11 ..	Ergot begun.—30 minims of liquid extract of ergot every 3 hours.	" 20 ..	6.0200	
" 6 ..	12.0800		" 21 ..	6.0200	
" 7 ..	12 ..		" 22 ..	5.0900	
" 8 ..	11.0800		" 23 ..	5.0700	
" 9 ..	11.0500		" 24 ..	5.0400	
" 10 ..	11.0500		" 25 ..	3.0500	
" 11 ..	11 ..		" 26 ..	2.0800	
" 12 ..	7.0900		" 27 ..	1.0600	
" 13 ..	7.0700		" 28 ..	1.0100	
" 14 ..	6.0500		" 29 ..	0.0700	He left the hospi- tal, and the ergot was discon- tinued. He came as an out- patient in a week. The urine had not increased.
" 15 ..	6.0400		" 30 ..		

When he was in the hospital, the urine was free from albumen or sugar. The specific gravity varied between 1000 and 1006; but as the quantity became less, it rose to 1017.

It will be seen from the foregoing notes that this well marked case of diabetes insipidus was uninfluenced by jaborandi, whilst on three occasions the quantity of urine was very greatly reduced by ergot.

CHLOROFORM AND ANÆSTHESIA.

By JOHN SMITH, M.D., F.R.C.S.E.,
Surgeon-Dentist to the Queen for Scotland.

SINCE the publication of Dr. Lauder Brunton's communication, in which he alludes to the administration of chloroform in insufficient quantity as a cause of danger, some attention seems to have been drawn to the subject in respect to this and other points.

I wish to make a few additional remarks on the matter; and that I am so far qualified to do so, will perhaps be admitted from the fact of an experience in the use of chloroform ever since its introduction in 1847, and a practical acquaintance with it, more or less, in every variety of case in which it ever has been given—over and above that of between three and four thousand times administering it in my own department of surgery. I trust my making such a statement may not be set down to any other motive than to show that I do not speak without sufficient data to entitle me in a certain measure to do so.

First of all, I entirely agree with Dr. Lauder Brunton's views regarding the degree of anæsthesia required for operating safely. In a

communication read by me before the Medico-Chirurgical Society here ten years ago, and published in the *Edinburgh Medical Journal* for 1866, I make the following remarks, which will be seen closely to coincide with those of Dr. Brunton.

"In all cases, anæsthesia, to be either satisfactory or safe, should be complete. I have repeatedly noticed patients, who have had chloroform on different occasions for such operations as came under my own department, express themselves as much less uneasy after a full dose—much less exhausted; and I may mention my own personal experience of this agent to bear out such a statement. So far it is more satisfactory; but it is also more safe. We know that one of the benefits of anæsthesia is the prevention of shock during an operation. We know also that shock tends to interfere with the heart's contractions, by the influence of the cerebro-spinal system on the cardiac ganglia through the par vagum. And this tendency does not always depend on the magnitude or importance of the operation. The operation may be a minor one, and the suffering inflicted very slight. The patient may be unconscious of pain, and yet the heart may, through the ganglionic system, be affected by the operation, and syncope result. . . . There can be no reasonable doubt that, of two patients equally under chloroform, if one be operated on in any way worthy of the term, and the other not, although neither has actually experienced pain, yet the one wakens up in a very different condition from the other: the one has suffered more or less from shock, the other has not."

In the same paper, allusion is made to the fact of the sudden diminution of the heart's action, on the operation being commenced, even during the state of anæsthesia. This I have myself frequently remarked even in tooth-extraction, and the fact has been over and over again observed in the greater operations. It is this tendency to sudden syncope which indicates the necessity for all operations being, if possible, performed with the patient in the recumbent position. In my lectures at Surgeons' Hall here in 1854 and subsequent years, and in the *Handbook of Dental Surgery* published as my class-book, this was, so far as I am aware, inculcated for the first time as applied to dental surgery; while, again, in the paper I am now quoting, I find it stated that "syncope during anæsthesia, as at all other times, is most apt to occur when the patient is in the upright or semi-upright position; and it need scarcely be here said that, in order to diminish the chances of a complication so serious, the horizontal posture should in all cases be the one adopted during the exhibition of chloroform. Such, however, is far from being the position generally chosen for minor operations. The preponderance in these operations of death associated with chloroform is, no doubt, attributable in a great measure to a contrary practice, it being very generally considered not worth while in such cases to make a patient lie down."

With regard to the subject of death from the administration of chloroform, much has been stated calculated to create a very unfair prejudice against it. No agent possessing such power of abrogating sensation can possibly be free from a certain amount of risk in its employment. Thus much, therefore, is to be understood as admitted in speaking of chloroform. But a great deal beyond this has been attempted to be made out. Every death now occurring during an operation in which chloroform has been given is set down to its effect, and sensational paragraphs too often appear in our daily press spreading such ill-authenticated reports. In fact, no deaths under such circumstances are now-a-days attributed to "shock" and other causes which used, before the days of anæsthesia, to be set forth in explanation of such cases, which, by the way, were far more numerous than they now are. But these cases, although fewer, still do occur; and that chloroform is unwarrantably blamed in them is, I fear, equally true.

Dr. Brunton mentions a case where chloroform was to have been given, and in which the patient suddenly died during the operation without its use. As I happen to have been present on that occasion, I may mention that the non-administration of the anæsthetic was not due to the bottle being broken, but to Professor Simpson, who was to have administered the chloroform, not arriving in time. Here the patient died on the first incision being made through the skin over the hernial sac. Shortly before this, again, Dr. John Argyll Robertson here had met with a similar case—strangled hernia—where the patient died while the groin was being shaved, and where no anæsthetic had been given. And, since the introduction and general use of chloroform, did time and space permit, many such cases could be cited where circumstances rendered the administration of any anæsthetic impossible. Even in tooth-extraction, this has occurred; and Mr. Holmes mentions the case of a patient whom the dentist found dead in the chair on returning from a side table, where he had been selecting the instrument to be used. There is very little doubt that, had chloroform been employed

* A litre is = 35¼ fluid ounces.

in these cases, it would have been subject to strong suspicion, if not to actual blame, as the cause of death.

Lastly, there is an important point connected with the numerous alleged deaths from chloroform which seems to be overlooked or set aside, as undeserving of notice; and that is the extreme discrepancy prevailing in the account of *post mortem* appearances. In the case of any other poison producing death, we find certain specific and uniform indications of its peculiar toxic action; whereas in those ascribed to chloroform the *post mortem* appearances range from those presented by death from syncope, on the one hand, to those produced by asphyxia, or those of the very opposite characters to syncope, on the other; arguing it to be extremely probable that one or other of them was during, and not owing to, anæsthesia from this agent. I will add a single other quotation on this topic from the paper to which I have so largely ventured to refer. It is as follows.

"In the fifty cases collected by the late Dr. Snow, the condition of the heart was very various. In nearly twenty, it was either gorged or tolerably filled with blood. In others, it was empty, but stained with blood, showing that probably it had been emptied only at the *post mortem* examination. Dr. Berend of Hanover details, among others, ten cases, in all of which the heart was empty, flaccid, collapsed, and doubled upon itself. In other cases from time to time recorded, the heart was firmly contracted. In many it was diseased, and in not a few the cause of death was found in some other organ altogether, such as the spinal cord. Again, the opinions as to the mode of death are not less various. Dr. Sibson, in 1848, attributed the fatal result to death of the heart; Dr. Snow, to what he termed cardiac syncope; Dr. Black, to asphyxia; Caspar, to neuro-paralysis; Sansom, to both syncope and asphyxia, etc. A conflicting mass of evidence, from which it appears impossible to derive any other conclusion than that the deaths described were due to no one cause, but had been produced in a manner widely different, according to the circumstances of the case."

A word here may be said regarding nitrous oxide as an anæsthetic. Although in the daily use of this agent, I fear that we can scarcely yet place it in comparison with chloroform, allowing that its mode of action were similar, which is very doubtful. The limited sphere of its application in medicine and surgery, the condition of patients under its influence, and some of its possible bad effects not being immediate, but more likely to show themselves at an after date, preclude its being used as anything like a fair standard by which to judge of chloroform. It is an excellent, an invaluable agent in its own place; but neither in medicine nor surgery is it ever likely to supersede such anæsthetics as ether or chloroform. In fact, if such were attempted, it seems very doubtful whether its present small rate of mortality would not increase in rather a startling manner; while its efficiency as a substitute would be found at fault in many ways.

CASE OF ACUTE POISONING WITH CHLORAL-HYDRATE.

By DAVID YOUNG, M.D., Florence.

A FEW days ago, I was called to see a gentleman who had just recovered from an attack of delirium tremens. He was feeble looking and nervous, and wished me to prescribe for persistent sleeplessness. For many months previously, he had taken chloral-hydrate in doses of twenty-five grains, with excellent results; but, since his last attack, the medicine had failed to act as before. I suggested a slight increase of the dose, and ordered the following mixture.

R Chloral hydrat. $\mathfrak{z}\text{ij}$; glycerini $\mathfrak{z}\text{ij}$; syrupi $\mathfrak{z}\text{ss}$; aquæ q. s., ad $\mathfrak{z}\text{iv}$. Misce.

A sixth part to be taken at bedtime, and half a dose more in two hours, if necessary. I was hastily summoned in the night; and, when I reached the hotel, found the patient in a deep stupor, the breathing slow and difficult, and the surface of the body, especially the extremities, very cold. In reply to my question if he had taken the medicine as ordered, his wife told me that he had come home about three hours before, and had apparently been drinking. He said, just before he went to bed, "I will now take the doctor's draught", and poured out, as she thought, one dose. Shortly after taking it, he became very restless, complained of being breathless, and of severe cramps in the legs. On attempting to get up for a drink of seltzer water, he staggered and fell on the floor. His wife became alarmed; and, after lifting him up, discovered that the bottle which had contained the chloral was empty; he had taken the whole—180 grains.

I was immediately sent for; and, on arriving, found him as above

stated. Acting on the suggestions given by my friend Dr. Lauder Brunton for the treatment of this accident, I ordered blankets and hot bottles to be procured at once. In the brief interval which elapsed before they were got ready, I noticed the following points in the condition of the patient. His breathing was slow and difficult; respirations 11 in the minute. Pulse 76, feeble and irregular. The temperature in the axilla was 35.6 deg. C. (96.08 Fahr.). The first sound of the heart was indistinct, and its action feeble. The eyelids were opened with difficulty. The conjunctiva was deeply injected, and the pupil firmly contracted. The surface of the body was very cold.

The lower limbs were wrapped in warm flannel, and hot bottles applied to different parts of the body; and a gutta-percha water-bag was filled with hot water, and laid over the cardiac region, and the patient carefully covered with blankets. The effects of the warmth were decided, and soon apparent. The temperature slowly rose, the breathing improved, and the heart regained its power. Half a teaspoonful of Darby's extract of meat, with a teaspoonful of brandy in a small cupful of hot water, was now ordered to be given every two hours.

Six hours after the application of the warmth, and nine after the taking of the chloral, the patient's condition was noted as under. Breathing was tranquil; respirations 16 per minute. The heart-sounds were distinct; its action was somewhat tumultuous. Pulse 92, full and soft. The pupil was contracted. The conjunctiva was injected, and he complained of muscular pains in the arms and legs. The patient continued in a state of great drowsiness for fifteen hours. He was easily roused, but immediately went off again to sleep.

In two days, he was quite well. I have no doubt that the effect of the application of the warmth, in this case, was to prevent another being added to the list of those whose death has been caused by chloral injudiciously taken.

A NOTE ON ANGINA LUDOVICI.

By F. MURCHISON, M.A., M.B. Edin.,

Assistant-Physician to the Crichton Royal Institution, Dumfries, N.B.

Two years ago, during my residence as medical officer in Harris, in the Hebrides, some cases of what I now believe to have been the *Angina Ludovici*, occurred among the natives of the island. The disease was different in its symptoms and peculiarities from any throat-affection that either my experience or reading had brought under my notice. I vainly tried to diagnose the malady, and as vainly read whatever medical literature of the past and present I had access to; but I could find nothing, having a name, similar to it. It attacked children and young persons of various ages. The youngest patient under my care might have been three years of age; the oldest was not certainly over thirty. The symptoms were uniformly of the same type, save that they were modified according to the diathesis and idiosyncrasy of the individual invaded. As a rule, the older the patients, the more intense were the sufferings, and the better developed were the throat-tumours that occasioned them. It is worthy of mention that the anæmic, pale, and pasty fair-haired, suffered in the proportion of six to one of the plethoric and healthy-looking; and that the disease among the latter did not occasion the prostration which it did among the former. Indeed, the cases among the absolutely robust were so few and far between, that I was induced—in the absence of a more plausible one—to adopt the theory that the sickness was one of the numerous manifestations of a scrofulous habit of body, incited into activity by the prevalence of very cold weather, insufficient clothing, want of cleanliness, and inappropriate food. All the cases were purely idiopathic, and independent of any local lesion. They all occurred in winter, and in close succession; but they were not confined to one isolated locality. One island seemed to favour the growth and development of the affection more than any other situation; but the six cases that occurred in it seemed to have arisen independently of infection for their origin, and rather to have commenced in an unhealthy locality, destitute of proper sanitary arrangements. No two cases occurred in the same family, although isolation of the sick from the healthy was not enforced; so that I am induced to believe that the disease was occasioned by climatic and insanitary influences rather than by contagion.

The disease began with a chill, a stiffness of the muscles that move the tongue and throat, a feeling of general *malaise*, discomfort and prostration, loss of appetite, and, in a word, with the feelings that are frequently the precursors of the specific fevers—symptoms which preceded the development of the disease in every instance. This incipient stage was followed by another of much distress and discomfort to the patient. The vicinity of the submaxillary gland, the gland itself, and