time. The intercurrent attack of pneumonia was probably due to purulent infiltration from the ear. The mastoid was implicated by extension of the suppuration from the meatus and tympanum.

[To be continued.]

EFFECTS OF VARIOUS DISEASES ON THE WEIGHT OF THE BRAIN IN 2,050 SANE AND INSANE ADULTS OF BOTH SEXES.

By ROBERT BOYD, M.D. EDIN., F.R.C.P.L.,

Late Physician-Superintendent of the Somerset County Lunatic Asylum, etc.

THE following record of the effects of various diseases on the weight of the cerebrum, cerebellum, pons, and medulla, in the sane and the insane, is taken from the necrological registers kept by me for thirty years at the St. Marylebone Infirmary and Somerset County Lunatic Asylum. As far as I know, the subject is one which has not hitherto been referred to by any author, and I think it may be of sufficient interest to bring before one of the sections at the Edinburgh meeting of the British Medical Association.

The citation of figures must be wearisome to all, but it is the only means by which the facts can be brought clearly under notice: I will, however, endeavour to be as sparing as possible in this particular. The sane adults, and many of the insane—classified according to age—are included in my tables of the weights of the human body and internal organs in the *Philosophical Transactions* for 1861. From these cases, and others of the insane subsequently examined, it was found that, as regards the sexes, the average weight of the encephalon was from four to five ounces heavier in males than in females, and that there was a difference in height of an equal number of inches. The proportions of the different parts of the encephalon, estimated according to the height, in 400 males and 325 females, insane, were as follow—the average height being in the males sixty-six, and in the females sixty-two inches.

TABLE I.

CEREBRUM.		Cerebellum.	Pons and Medulla.	Encephalon.	Sex.
Right Hemisphere.	Left Hemisphere.				
.301	.302	.077	.016	.607	Males
.294	.295	.076	.016	.681	Females

From the above, it appears that the average weight of the cerebrum was .007 more in males than in females; the cerebellum .001 heavier in males than in females; the pons Varolii and medulla alike in both sexes; and that the left cerebral hemisphere was heavier than the right by .001 in both sexes. The encephalon was .016 heavier in males than in females. Inequalities in the cerebral hemisphere have been frequently observed, especially in epileptics; and cases by me have been published in the thirty-ninth volume of the *Medico-Chirurgical Transactions*.

The heaviest brains have not unfrequently been those of epileptic idiots. Hypertrophy of the connecting tissue of the brain is stated to have been found in such cases; and to the same cause may probably be attributed cases of convulsions and sudden death in children, of which there are instances in the tables referred to amongst the cases examined in the St. Marylebone Infirmary. In these cases of sudden death, apparently in healthy children, when the skull-cap was removed, the dura mater was observed to be extremely tense and shining, and, when divided, the cerebrum projected considerably over the edges of the skull, the convolutions were flattened, the ventricles devoid of fluid, and their sides pressed together, the other internal organs being natural.

The following table shows the average weight of the cerebrum, cerebellum, pons Varolii, and medulla, and the encephalon in the sane and insane of both sexes, commencing with diseases of the nervous system, including those of the brain and membranes, and also of the spinal cord; diseases of the chest, under three heads—pleuropneumonia, bronchitis and asthma, pulmonary phthisis; diseases of the heart and dropsy; diseases of the stomach and abdominal organs; and, lastly, fevers, including the eruptive and erysipelas. The mental disorder in the insane of both sexes is shown on the annexed comparative table. (Table III,

on next page.) Of the 2,050 cases (616 sane males, 653 sane females, 430 insane males, and 351 insane females), the principal causes of death were as follows.

TABLE II.

Diseases.		SANE.		INSANE.	
		Males.	Females.	Males.	Females.
Nervous	Brain	Per cent. 17.3	Per cent. 25	Per cent. 32.5	Per cent. 15.7
	Spinal cord	.7	.1	14.4	7.4
	Totals	18	25.1	46.9	23.1
Lungs.	Pleuritis and pneumonia	22.1	17.3	21.6	17.4
	Bronchitis and asthma	11.1	13.6	3.7	8.8
	Phthisis	28.7	14.8	13.7	22.8
	Totals	61.9	45.7	39	49
Heart and dropsy		5.5	9.6	3	3.4
Abdomen		7	13.6	11	19.1
Fevers		7.6	6	.1	5.4

From the comparative table of the effects of various diseases on the weight of the brain and its several parts in the sane and insane of both sexes, it appears that, taking the total numbers, the encephalon was heaviest in the insane; the difference being in the males three-quarters, and in the females one ounce and a quarter. The average weights are as follow.

TABLE IV.

Cerebrum.	Cerebellum.	Pons & medulla.	Encephalon.	
41.4	5.1	1	47.5	In sane males
41.8	5.4	1	48.2	In sane males
36.4	4.6	.9	41.9	In sane females
37.5	4.7	1	43.2	In insane females

These averages, taken from the whole of the cases, will serve to show what effect is produced on the average weight under the various diseases enumerated in the table.

With respect to diseases of the nervous system, which include diseases of the brain and membranes, as well as those of the spinal cord, we find the cases most numerous amongst the insane males, amounting in them to very nearly 47 per cent. of their whole number, and in females to 23 per cent. A considerable proportion of this excess in males (insane) is due to the frequency of general paralysis—a disease, as the symptoms indicate, usually arising from disease of the spinal cord. This fact, confirmed by the microscopical observations of Mr. Gulliver, was first stated by me in 1848, also in subsequent annual reports of the Somerset Asylum, and in the *Journal of Mental Science*. The fact has since been confirmed more or less by other observers at home and abroad.

In diseases of the brain and membranes, the encephalon was in the males one ounce below the average above given in the sane, and two ounces in the insane, and in the females about three-quarters of an ounce below the average. In both classes, four-fifths of this loss was in the cerebrum in males and in the sane females; in the insane females, the entire loss was in the cerebrum, and in them the cerebellum was one-tenth of an ounce above the average. It will be seen that there is a considerable difference in the weight of the cerebrum in diseases of the brain and spinal cord, owing to cerebral wasting usually accompanying paralysis of the insane.

In the four cases of disease of spinal cord in the sane males, the encephalon was half an ounce above the average, the increase in weight being due to the cerebellum and pons Varolii. In the insane males a very different state of the encephalon was found; it was three-and-a-quarter ounces below the average, being the lowest average of that organ in any of the various diseases: the loss of weight was entirely in the cerebrum. In the females, also, the loss of weight was entirely in the cerebrum, and amounted to 1.1 ounce.

The form of mental disorder in the foregoing (282) cases of cerebro-spinal disease was in the following ratio.

Mania	19.4 per cent. males; 23.4 per cent. females.
Melancholia	4.4 " " 12.3 " "
Fatuity and dementia	12.9 " " 12.3 " "
General paralysis	35.3 " " 29.5 " "
Epilepsy	25 " " 16.5 " "
Idiocy	3 " " 6 " "
100	100

TABLE III.—Comparative Table, showing in both Sexes in various Diseases the Average Weight of the Cerebrum, Cerebellum, Pons Varolii, Medulla, and Encephalon, in 2,050 Cases of Sane and Insane Patients, examined in the St. Marylebone Infirmary and Somerset County Lunatic Asylum.

Mental State.	Diseases.	MALES.					FEMALES.					ASYLUM CASES.		
		No of Cases.	Cerebrum.	Cerebellum.	Pons; Medulla.	Encephalon.	No. of Cases.	Cerebrum.	Cerebellum.	Pons; Medulla.	Encephalon.	Mental Disorder.	Males.	Females.
Sane	Brain and membranes	107	40.6	5	.9	46.5	163	35.9	4.5	.8	41.2	Mania	33	15
												Melancholia	8	8
												Dementia	11	8
												Fatuity	7	
												General paralysis	28	8
Insane	Do. do.	140	40	5.2	1	46.2	55	36.8	4.8	1	42.6	Epileptic mania	48	12
												Idiocy	5	4
												Total	140	55
Sane	Spinal cord	4	41.3	5.6	1.1	48	1	32.2	4.5	1	37.7	Mania	6	4
												Melancholia	1	2
												Dementia	8	2
												General paralysis	43	16
Insane	Do.	61	38.7	5.3	1	45	26	36.4	4.8	.9	42.1	Epileptic mania	2	1
												Idiocy	1	1
												Total	61	26
Sane	Pleuropneumonia	136	41	5.1	1	47.1	113	36.9	4.6	.8	42.3	Mania	35	14
												Melancholia	15	20
												Dementia	9	14
												Fatuity	7	2
												General paralysis	7	
Insane	Do.	93	40.8	5.2	1	47	61	36.8	4.8	.9	42.5	Epileptic mania	15	9
												Idiocy	5	2
												Total	93	61
Sane	Bronchitis; asthma.....	68	40.6	5	.9	46.5	89	36.9	4.6	.9	42.4	Mania	6	9
												Melancholia	1	7
												Dementia	4	6
												Fatuity	3	4
												General paralysis	1	
Insane	Do. do.	16	41.5	5.3	1.1	47.9	31	37.9	4.8	1	43.7	Epileptic mania	2	4
												Total	16	31
Sane	Phthisis	178	41.1	5.1	.9	47.1	97	36.8	4.6	.9	42.3	Mania	31	35
												Melancholia	7	18
												Dementia	10	8
												General paralysis	2	
Insane	Do.	59	40.7	5.2	1.1	47	80	36.9	4.7	1	42.6	Epileptic mania	4	13
												Idiocy	5	6
												Total	59	80
Sane	Heart; dropsy	34	42.8	5.3	1.1	49.2	61	37.6	4.7	.8	43.1	Mania	6	4
												Melancholia	3	1
												Dementia	1	5
												General paralysis	1	
Insane	Do. do.	13	42.2	5.1	1	48.3	12	39.2	4.7	1.1	45	Epileptic mania	2	1
												Idiocy	1	1
												Total	13	12
Sane	Abdominal organs	43	41.4	5.1	1	47.5	89	38	4.8	.9	43.7	Mania	24	40
												Melancholia	4	11
												Dementia	9	12
												Fatuity	5	2
												General paralysis	2	
Insane	Do. do.	47	42.6	5.4	1.1	49.1	67	38	4.8	1	43.8	Epileptic mania	2	1
												Idiocy	3	1
												Total	47	67
Sane	Fevers, etc.	46	42.2	5.2	1	48.4	40	36.8	4.7	.9	42.4	Mania	1	15
												Melancholia		2
												Dementia		1
Insane	Do.	1	48	6.5	1	55.5	19	38.2	4.7	1	43.9	Idiocy		1
												Total	1	19

As regards the diseases of the lungs (in the deaths assigned to pleuropneumonia—131 in males and 113 in females in the sane, and 93 males and 61 in females insane)—in the sane males there was a loss of four-tenths of an ounce in the cerebrum, whilst in the sane females there was a gain of half an ounce in the cerebrum as compared with the average; in the insane males the cerebrum was six-tenths of an ounce below the average; in the insane females the same organ was three-quarters of an ounce below the average, while the cerebellum and pons Varolii together were of the average weight.

The cases of bronchitis and asthma were in number about one-half of the foregoing. There was a loss of eight-tenths of an ounce in the cerebrum, and of one-tenth each in the cerebellum and pons in the sane males, whilst in the sane females the cerebrum was five-tenths of an ounce above, and the cerebellum and pons were equal to the average. In insane males the cerebrum was three-tenths of an ounce below, and the cerebellum and pons together equal to the average; and in the females of this class the encephalon was again five-tenths, and the cerebellum one-tenth, above the average weight; so that, whilst in both

sane and insane males there was a loss of weight, there was, on the contrary, an increase in the females of both classes.

Of the three subdivisions of lung-disease, the cases of pulmonary phthisis were most numerous in the sane, amounting to 178 in the males and 97 in the females; also in the insane females to 80. In the insane males, pleuropneumonia was most frequent. The loss of weight in the cerebrum, as compared with the average in these cases of phthisis, was seven-tenths of an ounce in sane males, and there was an increase of one-tenth each in the cerebellum and pons; whilst in sane females the cerebrum was four-tenths of an ounce above, and the cerebellum, pons, and medulla, equal to the average. In the insane, the decrease in the cerebrum was eleven-tenths in the males, and six-tenths of an ounce in the females; and the cerebellum and pons in the males were one-tenth of an ounce below the average: in the females, the cerebellum and pons Varolii were of the average weight.

In disease of the heart, the encephalon attained the highest average weight in each sex and in both classes, being in males one ounce and three-quarters, and in females one ounce and a quarter above the average in the sane, the increase in weight being in all the parts. In the females it was confined to the cerebrum; in the insane males the increase in the cerebrum was four-tenths, and in the females one ounce and three-quarters; in the cerebellum, three-tenths in males; in the pons Varolii and medulla, one-tenth in females.

In the diseases of the abdominal organs in the sane males, the several parts of the encephalon were of the average weight; in the females the cerebrum was one and a half, and the cerebellum a quarter of an ounce above the average; in the insane males the cerebrum was eight-tenths, and the pons one-tenth of an ounce above the average; and in the females the cerebrum was five-tenths, and the cerebellum one-tenth of an ounce, above the average weight.

In fevers, the encephalon in the sane males was nine-tenths of an ounce, and in the females five-tenths of an ounce above the average; in the insane females the cerebrum was seven-tenths of an ounce above the average. There was but one male insane classed as a case of fever; the encephalon, unusually large, weighing fifty-five ounces and a half.

Of the diseases of the abdominal organs in the insane, including fevers, in females there were 133 cases; and the forms of mental disorder in the 47 males and 86 females were in the following ratio per cent.

	Males.	Females.
Mania	51.2	64.1
Melancholia	8.4	15.6
Dementia and fatuity	29.8	17.2
General paralysis	4.2	—
Epilepsy	—	.2
Idiocy	6.4	2.4
Totals	100	100

In the whole number of insane—430 males and 351 females—the forms of mental disorder were in the following proportions per cent.

	Males.	Females.
Mania	37.7	40.4
Melancholia	8.8	18
Dementia and fatuity	20.8	16.4
General paralysis	15	10
Epilepsy	12.5	10.9
Idiocy	5.2	4.3
Totals	100	100

It may be observed, in conclusion, that the "assigned cause of death" was often difficult to determine in the insane, as it was frequently—in pulmonary disease, for example—associated with chronic inflammation of the cerebral membranes.

1. In diseases of the nervous system in the males, the numbers were 18 per cent. in the sane, and as many as 47 per cent. in the insane; in the females of each class, the numbers were about 25 per cent., the cerebrum being below the average weight. The larger ratio in the insane males was due in great measure to the fatality of general paralysis in that class, which is usually accompanied by inflammation of the spinal cord. This statement was first published by me in 1848, and subsequently confirmed by further examinations—some made by Dr. J. Ogle, at the Pathological Society of London.

2. The three pulmonary diseases, especially phthisis, were most frequent in sane males: in the sane and insane females the numbers were nearly equal. The weight of the cerebrum was in the males below, and in bronchitis and asthma, in females, above the average. In diseases of the vascular system, the brain was above the average, and the numbers in the sane were more than double the number in the insane.

3. The diseases of the abdominal organs were a third more in the insane than in the sane; and the brain was above the average weight,

as it also was in fevers, which were most frequent in the sane, the insane being comparatively free.

From these observations on the different parts of the encephalon, it may almost be said that the variations in weight were confined to the cerebrum.

NOTES ON THE CLIMATE OF ALGIERS.

By WILLIAM THOMSON, M.D.,
Resident English Physician, Algiers.

At this time of year, when many invalids in this country are beginning to ask themselves or their doctors where they ought to go to spend the winter months, I venture to bring forward the claims of Algiers as a health-resort of no small importance; and, in doing so, I have endeavoured to make my remarks as short and concise as possible, and to avoid all semblance of vaunting the comparative merits of the place: a practice which, I regret to say, is too common amongst some of my professional brethren. Having been in many climates, I have never as yet found, and never expect to find, a perfect one; certainly Algiers is not; but I merely wish to describe it shortly as it is, and give a few short remarks from my own small experience as to the cases most suitable for sending there, in the hope that they may be of some small use to my fellow-practitioners at home, from many of whom I have received letters, asking for information regarding Algiers. My personal experience of Algiers and its climate only extends back as far as the last two winters; but the meteorological notes which I have made use of are taken from the published reports of M. Bulard, the director of the observatory at Algiers, who has laboured hard and conscientiously in making observations there during the last fourteen years, which observations can be far more implicitly relied on than stray ones made by casual residents.

The town of Algiers lies in 36.47 deg. N. latitude, and 3.4 deg. E. longitude, almost exactly five hundred miles S.S.W. of Marseilles, with which town it has communication thrice weekly by the steamers of the Messageries Maritimes and Valery Companies, which do the passage in thirty-four hours. The town lies on a low range of hills called the Sahel, a prolongation of the Lesser Atlas mountains, which runs out to the sea here in the shape of a spur in a north-easterly direction, forming the western extremity of the Bay of Algiers, the breadth of which from here to its eastern extremity, Cape Matifon, is about sixteen miles. As the town and suburb of Mustapha, the favourite residence of the English visitors, lie on the south-east side of this range, they are thus protected from the north-west wind, the most prevalent one at Algiers, but which, from its passage across the sea, has no resemblance to the dry piercing mistral of the South of France. The form of the town is triangular, the base of the triangle being the French part of the town, and lying along the sea, whilst the cone comprises the old Arab quarters, and stretches up the hill, ending at the old fort or kasbah, the former residence of the Deys of Algiers. The town and suburb of Mustapha face the south-east, so that they enjoy a good exposure to the sun nearly all the day, whilst the view across the bay, with the fine range of the Djurjura Mountains in Kabylia, is of great beauty. The season for invalids may be said to be from the beginning of November to the end of April, though many English residents who are in the habit of staying on later say that May is the pleasantest month of all in Algiers. I should advise no invalids, however, to arrive there before November 1st, as the heat in October is sometimes still very trying. With that view, I should recommend all intending visitors to Algiers who leave England in the beginning of October (and I most decidedly think none, with few exceptions, suffering from pulmonary complaints should ever delay their departure later than the 10th of that month) to cross over to Paris, where the temperature is usually warmer, and where the air, at any rate, is free from the fogs to which London becomes liable about this time, and which are often so harmful; after spending a week or so in Paris, they can then move gradually southwards by stages, such as Dijon, Macon, Lyons, Avignon, Nismes, Arles, etc., to Marseilles, where they embark for Algiers. I would here offer another word of advice on the subject of dress, and would most earnestly advise all, especially invalids, to take plenty of warm clothing with them, as, though the temperature of Algiers during the winter is, on an average, much the same as our summer in England, yet slight variations in it are much more felt than the same variations at home.

The average mean temperature during the winter months in Algiers may be said to be 55.0 deg. Fahr.; the mean daily range between 10 deg. and 12 deg. Fahr. The mean height of the barometer at sea-level is 30.0 deg., and the daily variations are very slight, except