more so to vie with each other in the common cause of humanity. Then, perhaps, they may come to acknowledge the truth of the saying of Cicero in his oration *Pro Ligario*, "Neque enim ullâ aliâ re homines propius ad Deos accedunt, quam salutem hominibus dando".

EPIDEMIC CEREBRO-SPINAL MENINGITIS: REPORT OF *POST MORTEM* EXAMINATION IN THREE CASES OF THIS DISEASE.

By JAMES RUSSELL, M.D., F.R.C.P.,
Physician to the General Hospital, Birmingham.

THE two *post mortem* examinations which I here record represent the sole occasions on which I have had an opportunity of making a dissection after death from epidemic cerebro-spinal meningitis; they will, therefore, be conveniently detailed apart from the clinical history of other cases which have lately fallen under my care at the Birmingham General Hospital or in private practice, leaving an opportunity of inserting hereafter the clinical details of these cases.

The two examinations which follow afford confirmation of the truth of Mr. Simon's observations on this subject, that "for practical purposes, the state of the covering membranes of the nervous centres may be regarded as a mere index of changes more or less distinctive which these centres, in their own intimate composition, have at the same time undergone; and hence it is that the essential phenomena of the disease during life consist in disturbances, more or less grave, of the functions of those all-important organs" (Reynolds's *System of Medicine*, 2nd ed., p. 513).

CASE I occurred in the person of a girl, aged 14, who was admitted on June 22nd, on the eighteenth day of the disease, and died on July 16th. I shall merely record here that the origin of the disease was acute, and dated from June 5th; it was referred to the child having fallen into some water and remained in her wet clothes. Certainly, she felt ill the same evening, and shivered and vomited the following day; the vomiting ceased in three days. The headache was the prominent symptom; it was seated in the occipital and frontal regions. The intellect was clear from the second or third day. The symptoms assumed somewhat of an intermittent type, but, during the last ten days of life, settled into a state of apathy, with much depression of vital power, ending in torpor and death by asthenia. The urine was very phosphatic. The temperature manifested a tendency to morning remissions and evening exacerbations, very decided in two quarter periods, very imperfectly marked in the other two quarters. The maximum temperature never exceeded 102 deg.

POST MORTEM EXAMINATION.—No external marks on the body; a loose fibrinous riband in the longitudinal sinus; fluid blood in this and in the lateral sinuses; very marked injection of the minute vessels over the entire surface of the brain; but the arachnoid and pia mater retained their normal transparency, and could be stripped off the brain with ordinary facility. To this statement there are the following exceptions.—Just at the anterior extremity of the superior vermiciform process of the cerebellum was a collection of not very recent yellow lymph in the pia mater, over a space not half an inch square; a similar deposit at the base of the cerebrum covered the chiasma, and a few broad streaks existed in the membrane at each Sylvian fissure, running for a short distance along the vessels. The lateral ventricles had plainly been subjected to distension; at the present time, they contained a moderate quantity of opaque highly albuminous fluid, in which Dr. Saundby found blood-corpuscles and small round granular nuclear cells. Their ependyma was sufficiently thickened to float in water as a consistent semi-transparent membrane. The surface of all the nervous masses in the ventricles showed minute branching vessels filled with blood, and the surface-tissue was rather soft. The blood-points throughout the white tissue of the brain were large and numerous, and on the surface of the convolutions the perivascular spaces were very distinct. The grey substance throughout the cerebrum was deeply tinged with pink; this pink colour, very apparent in the convolutions, was especially striking in the large ganglia. The ganglion-cells in the corpora striata were well formed. Dr. Saundby examined some of the minute vessels of the grey matter of the brain, and reported them simply loaded with blood. In all other respects, the brain-substance was healthy. The spinal cord presented nothing abnormal, excepting injection of the pia mater, which, though considerable, was less marked than in the brain. There was the same strongly marked pink colour of the grey matter as already noticed on the cerebrum. The tissue of the cord was of

normal tenacity; the rest of the nerves were healthy. The muscles of the back presented a granular appearance in the sarcolemma. There were fibrinous clots in the heart. The spleen was small but healthy. Nothing else was presented worthy of note. The blood was not unusually fluid.

CASE II.—The patient was 24 years of age. He was admitted for pleurisy. He had a very cachectic appearance, and gave a confused history of some foregoing illness; but neither in his description nor in the symptoms he presented was there anything to lead to the supposition of peculiarity in his case. He was recovering from his pleurisy, when, with no apparent reason, he was attacked with purposeless vomiting and retention of urine; the vomiting continued for three days and then ceased; but the man became rapidly apathetic, then torpid, and died asthenic on the sixth day of this last illness. The retention of urine passed into involuntary evacuation. Inquiry from the friends elicited a manifest history of cerebro-spinal meningitis, and ample confirmation was afterwards obtained from Mr. Bradley, who had attended the patient. Mr. Bradley described to me the symptoms of the simple form of the disease. The attack commenced on the 7th of last February; a brief relapse occurred; but convalescence was sufficiently established to permit of Mr. Bradley taking his leave on March 23rd. Of the subsequent attack of pleurisy, Mr. Bradley knew nothing; the patient came into the hospital on May 13th and died on the 29th.

POST MORTEM EXAMINATION.—The arachnoid and pia mater had undergone remarkable thickening over a limited space at the base of the brain, presenting a tough semitransparent membranous web covering the chiasma, stretching across the interpeduncular space, enveloping the pons Varolii and medulla oblongata and enclosing the cerebral nerves at their origin. The adhesion of the membrane to the pons and medulla was so close that it was impossible to separate it without damaging the nervous tissue. It covered the fourth ventricle and thence extended down the cord. The thickening extended on the under surface of each cerebellar lobe for about an inch in breadth, thence gradually ceasing; it appeared also as a broad band running across the anterior edge of the cerebellum, above the mesocephale. A very scanty deposit of firm yellow lymph existed in the pia mater, upon the chiasma and beneath the parietal layer of the arachnoid upon the basilar process. In all other parts, the membranes of the brain were quite healthy. The ependyma of the lateral ventricles was much thickened, and could be raised as a semitransparent membrane of considerable tenacity. The lateral ventricles were largely dilated, but no unusual quantity of fluid was found in them at the time of the *post mortem* examination. The tissue enclosing the ventricles had an unusually fibrous appearance, as though it had been subjected to stretching. No abnormal appearance existed in the brain, except that its tissue was generally softer than usual, but not in any one part particularly. The pia mater and visceral arachnoid of the cord were much thickened through the entire length, closely encircling the cord with a dense envelope, which bound down the roots of the nerves and adhered to the cord itself along the posterior surface only. The apex of one lung was broken up, and there were remains of fluid in the pleural cavity. No other morbid change existed in the body.

CASE III.—In a third *post mortem* examination on a patient of my colleague (Dr. Fletcher), who died after an illness of six weeks, the principal appearances consisted in the presence of a soft flocculent lymph over a very limited space extending from the chiasma to the posterior edge of the pons Varolii; of a small patch of the same material over the root of the right suboccipital nerve; and of two or three similar patches within the subarachnoid space of the upper dorsal region of the spinal cord. The cerebral arachnoid was more resistant than usual, and was opaque where stretching across the sulci: in the upper dorsal region the spinal arachnoid was also opaque. The ventricles were distended. The condition of the cerebral tissue closely resembled that in my first case. One kidney was excessively atrophied, thereby accounting for the presence of a considerable amount of albumen in the urine during life.

RECURRENT EPISTAXIS FROM MALARIAL INFLUENCE.

By E. M. SINCLAIR, M.D., Surgeon-Major Royal Artillery.

WITH reference to the paper by Surgeon-Major J. H. Porter of Netley, on Intermittent Hæmorrhage from Malarial Influence, read before the Royal Medical and Chirurgical Society in February last, the following case of intermittent epistaxis from the same cause may prove of interest.