under tension, the circulation gives way at its weakest point. And, unlike a chain, the tension is greatest at this particular point.

No other vascular region is equally exposed to those sudden or rapid flows which so sorely try the resisting force of the vessels. There is abundant reason to conclude that ovulation, or at least an attempt at ovulation, proceeds during pregnancy. Hence is explained the fact that abortion and the bleedings associated with placenta prævia mostly occur at menstrual epochs. It seems a happy provision of Nature that the break-down happens at this place. The system, unable to bear the burden of pregnancy, saves itself by rejecting the ovum. If hæmorrhage do not take place from the uterus, abortion attending or not, or from some other mucous membrane, thus obeying the physiological law of hæmorrhages, extravasation would be the more likely to take place either into a serous cavity, as the peritoneum, into the parenchyma of the internal organs, or into the brain, producing apoplexy.

Depending partly upon the altered dynamic conditions of the circulation, and partly upon the altered constitution of the blood, various inflammations are apt to arise during pregnancy, but more especially soon after labour. These I can only glance at. But it must be obvious that this rapidly developed proclivity to inflammations must illustrate the causes of inflammation in the non-pregnant. Upon the same conditions depend essentially the phenomena of thrombosis and embolism. Indeed, in no other state do we find so many or such instructive examples of this blood-disturbance as in pregnancy. Did time permit, I could adduce evidence to show that, for the production of thrombosis, arterial or venous, the addition of a third factor, the entry of poisonous matter into the blood, is at least generally necessary.

What is the mode of genesis of that singular disease, *acute yellow atrophy of the liver*? Conjecture is nearly all that can at present be offered; but, since a very large proportion of all the recorded cases have occurred during pregnancy, it is surely in the direction to which I am pointing that a solution must be sought. Taking note first of the known modifications wrought in the blood by pregnancy; secondly, of the increased strain thrown upon the large glands, especially upon the liver; thirdly, of the liability of the blood to be invaded and altered in its chemical form by poison; fourthly, of the peculiarly impressionable explosive condition of the nervous centres—we may see dimly looming in the distance the path that shall lead to the synthetic thought that will reveal the truth.

One of the most curious and least understood phenomena of pregnancy is *pigmentation*. Many isolated facts may suggest lines of speculation to account for the production or deposit of pigment in certain cases; but the hope of discovering the combining links that shall produce a true theory of pigmentation must lie in the observation of pregnant women, who present so many examples of the process. That pigmentation is intimately dependent upon blood-conditions, cannot be doubted; but facts are not wanting to show that the nervous system is concerned in the process. In the investigation of this problem, there lies the promise of a rich harvest of physiological and pathological discovery. Here I cannot help citing a most remarkable experiment due to the perils of the sea recorded in the touching narrative (*Blackwood's Magazine*, 1876, by F. Wordsworth) of the sojourn of forty-nine persons for nearly seven months on one of the Crozet rocks, who had escaped from the wreck of the *Strathmore*. After living some time on sea-birds and their eggs, exposed to wet, cold, and fatigue, it was found that the dark-haired persons had grown fair and the fair-haired had become flaxen, with pink and white complexions; and that, conversely, the pristine pigmentation was regained when return was made to the ordinary diet and conditions of life.

In this place, I take the opportunity of calling attention to a clinical fact not previously, at least to my knowledge, recorded. My late resident-assistant at St. George's Hospital, Mr. Lacy, observed in two cases of postpuerperal inflammation that the temperature rose at every menstrual epoch two or three degrees, sinking again on the subsidence of the flow. I have since made other observations in point. It is highly probable that we may in this fact discover indications in the treatment of inflammations in women. An associated fact in one case is very interesting. It was a case of pyæmia with arthritic complications. At the menstrual periods, the temperature rose to 101.5° and hæmoptysis attended. She had never had hæmoptysis before.

But the warning hand of inexorable Time reminds me that I must respect his authority and your patience. The subject I have feebly sketched, or rather pointed to, is inexhaustible in matter for observation and in its theoretical and pathological applications. I shall not have failed altogether if I incite others to work and think in this direction. I will not weary you with any further speculations, but will simply state one or two propositions which may fitly conclude this address.

1. The physiological phenomena evoked by pregnancy must be

studied and grasped as a whole; that is, as forces called into combined action to accomplish a definite object. We cannot arrive at a just idea of the state of any one organ, or system of organs, without taking into consideration the state of all the rest in their absolute and relative conditions. This may seem a somewhat obvious proposition, but I cannot call to mind a systematic treatise on medicine or even on obstetrics in which more than a faint glimmer of the lesson it propounds is manifest.

2. This general law may be stated. Since, in pregnancy, nearly every organ and the whole organism are specially weighted, and undergo extraordinary development and functional activity, so any defect or fault inherited or acquired, howsoever latent, will be liable to be developed or intensified under the trial. Hence pregnancy is the great test of bodily soundness.

3. Whether there be any latent defect or not, but more especially if there be a latent defect, the history of pregnancy supplies illustrations throwing a vivid light upon many problems in pathology complementary to that derived from the ordinary methods of clinical observation and pathological research. Pregnancy is a more luciferous experiment than was ever devised in the laboratory.

TWO CASES OF ADDISON'S DISEASE WHICH ENDED WITH CEREBRAL SYMPTOMS.*

By P. H. PYE-SMITH, M.D., F.R.C.P.,
Assistant Physician to Guy's Hospital, etc.

IT is not necessary before a medical society in England to defend the reality of Addison's disease, or to trace the steps by which our present knowledge of it has been acquired. Here at least pathologists have not been blind to the facts, which no difficulties in explanation can alter, and one might rather complain that this last and most striking discovery of Dr. Addison has diverted attention from his other and no less remarkable services to medicine.†

I may, however, remind you that, in 1855, the original short treatise on the *Constitutional and Local Effects of Disease of the Suprarenal Capsules* was published. Of its eleven cases, only the first six were undoubtedly genuine; but the doubts which reasonably existed on a subject incompletely worked out were, or ought to have been, set at rest by the publication in the *Guy's Hospital Reports* for 1862 of twenty-five genuine cases, with a commentary by Dr. Wilks, in which the characteristic lesion of the adrenals‡ was distinguished from others to which they are subject, and the clinical features of the disease were fully described. In 1866, Dr. Greenhow was able to publish in the seventeenth volume of the *Pathological Transactions* no less than one hundred and twenty-eight cases of morbus Addisonii verus; and the same physician has since added to these about fifty more, and has embodied the results of his investigations in the Croonian Lectures delivered last year before the College of Physicians, and afterwards published in this JOURNAL as well as in a separate form. To these lectures I may refer for a more detailed account of the literature of the subject.

The point to which I desire particularly to direct the attention of this Society is the occasional rapid termination of the disease, when apparently in a very chronic stage, by the sudden access of cerebral symptoms: a mode of death of which the following two cases are examples.

CASE 1. §—A boy, aged 14, came among my out-patients at Guy's Hospital last March, whose appearance had already led the house-physician to a correct diagnosis of the case. Several years before, he had suffered from caries of the tarsus, and Mr. Bryant had removed some dead bone with good result. Though somewhat thin and delicate in appearance, he appeared to be a healthy boy until a few months before I saw him, when his appetite began to fail; he complained of headache and pain in the back, and occasionally vomited. Meanwhile,

* Read before the Hunterian Society.

† Especially his recognition of the disease since known as idiopathic or pernicious anæmia, his classical paper on pneumonia, and his long-forgotten but now accepted doctrines on phthisis.

‡ I venture to use this word, familiar to comparative anatomists, as a more distinctive and convenient term than suprarenal capsule. The organs were described as *glandulæ* by their discoverer Eustachius, in 1564; by Wharton, a century later, as *glandulæ ad plexum* (sc. *solarem*); and it was the Danish anatomist Bartholinus who gave them the doubly misleading title of *capsulæ atrabiliarie*. Since they are neither true (secreting) glands nor lymphoid structures, since they do not form black-bile, and are only capsules when the medulla has decomposed into an apparent central cavity, it is better to use a term which is sufficiently short and characteristic, and does not involve any theory as to their function.

§ This case has already appeared in a paper on Morbus Addisonii, which I published in *Virchow's Archiv* for 1875.

the skin had gradually become dark, until, when I first saw him, the boy looked like a half-caste. His face was of a brunette complexion, and would have looked healthy enough but for the contrast with his light brown hair, hazel eyes, and English features. The back of the hands was of the same colour, the trunk and limbs lighter, except the loins and bend of the joints; the penis and scrotum were very dark. This characteristic distribution of pigment in the disease is strikingly different from that produced by exposure to air or sun, from the melasma of prurigo senilis, or the discoloration of jaundice; but it is precisely like that seen in the darker races of mankind, for in them also the genitals and the areola of the nipples are darkest, the face darker than the trunk, and the palms and soles the lightest parts. A microscopic section also of the skin of a patient affected by Addison's disease shows precisely the appearance of a similar preparation from a negro: pigment-granules deposited in the Malpighian layer of the epidermis, and, less abundantly, in the papillary layer of the cutis. The boy, though somewhat slender in build, was fairly nourished, and the mucous membrane of the mouth was not anæmic; there were one or two dark patches on the inside of the lips. The temperature, the blood, and the urine were normal, and the chest and abdomen gave no indications of disease. I took the patient into the hospital, and gave him the twentieth of a grain of phosphorus thrice daily. During the two months following, there was no notable change, except that the colour of the skin gradually deepened and several minute dark brown spots, like pigment-nævi, appeared on the face and trunk. There was some nausea and a feeble rather rapid pulse. The urine gave a slight precipitate with tincture of galls, but never contained albumen. One morning, he complained of severe pains in the head. He lay for about three hours screaming with pain, and then sank into an unconscious state till the evening, when he died. There were no convulsions and no paralysis, nor was there previous vomiting. At two in the afternoon, the temperature was 96.2; the pulse rapid and almost imperceptible at the wrist.

After death, besides the usual condition of the adrenals, which were shrunken, hard, and caseous, there was no other important lesion found. The heart showed the usual fatty degeneration of the ventricles. The lungs were normal; but there were slight old adhesions of one pleura and one of the bronchial glands contained a little caseous material. The lymph-follicles of the stomach and intestines (lenticular and solitary glands) were large and distinct even for a boy of 14. The thymus was also rather large. The connective tissue around the adrenals was thickened; but, on dissecting out the solar plexus, its ganglia appeared to be normal. The brain was large, weighing fifty-eight ounces; the fornix and septum lucidum were soft, probably from *post mortem* change, as there was other appearance of disease in the brain or membranes. There was no tubercle found, nor any disease of the vertebrae, tarsus, and other bones.

CASE II.—J. H., aged 27, was a patient of Mr. R. H. Cooke of Stoke Newington, to whom I am indebted for the following notes. He was a clerk, who was always taking medicine, and was thought by his friends to have little else the matter with him.* For several years, he had become darker in complexion. The exact time we could not ascertain; but his brother knew that the change had gone on for at least seven years, and Dr. Gooding of Cheltenham, whom the patient consulted a year before his death, believes that he then stated that his skin had begun to darken eight years before. One day last August, he went to bed in his usual health, and next morning was found cold and insensible on the floor of his room. He was put to bed and rallied enough to take a glass of wine and answer questions, but could not remember the attack, which must have come on in the early morning. He was soon afterwards seized with convulsions and, after a series of well marked epileptic fits, died comatose in the course of the following night. The fatal attack thus lasted about twenty-four hours. No urine was passed during this time. The pupils were sensible to light. We were only allowed to examine the kidneys and adrenals. The former were healthy and surrounded by a quantity of fat; the latter were shrunken, the one dense and contracted, the other soft and caseous. There was no thickening or adhesion of the surrounding connective tissue. The discoloration of the skin was characteristic in its distribution, the body looking like that of a mulatto. The face was very dark and the genital organs quite black. There was a fair amount of subcutaneous fat.

REMARKS.—It is not uncommon, during the extreme exhaustion which attends the last stage of morbus Addisonii, when uncomplicated by phthisis and not cut short by an intercurrent disease, for the patient to sink into a half-conscious condition, lying quiet, with half-open eyes,

slow gentle respiration, and almost imperceptible pulse. The state is apparently the same as that seen in some cases of diabetes, more frequently in fever, and occasionally in cirrhosis of the liver and in Bright's disease. It is thus described by Addison himself:—"With respect to the first mentioned form of cerebral disorder connected with renal disease, that of quiet stupor, it is in its most exquisite form probably the least frequently met with. The face is pale, the pulse quiet, the pupil natural, or at least obedient to light; and, although the patient may lie almost completely motionless, there is no paralysis; for, on attentively watching him for some time, he will be observed slightly to move all the extremities. By agitating him and speaking loudly, he may sometimes be partially roused for a moment, but quickly relapses into stupor as before; or it may not be possible to rouse him at all. There is little or no labour of respiration, no stertor, and no convulsions." (*On the Disorders of the Brain connected with Diseased Kidneys*: Sydenham Society's Edition of Addison's Works, p. 188.)

But, when this condition comes on as the most frequent termination of uncomplicated disease of the adrenals, it is always preceded by extreme and increasing prostration, with vomiting and syncope; while the sudden and more violent cerebral symptoms occurred in both the above cases during an almost latent state of the usual constitutional symptoms. The last patient seen by Dr. Addison himself (No. 18 in Dr. Wilks's paper, *Guy's Hospital Reports*, 1862), a lad aged 15, was delirious, with slight convulsions before death. In the thirty-seven cases given in detail in the appendix to Dr. Greenhow's volume, most died in the state of extreme prostration without loss of consciousness described by Dr. Addison, three were delirious, one of these and two others died comatose, and in three cases there were epileptic convulsions. In one of the last, there were found congestion of the brain, excess of serum in the ventricles, and a diffused state of the fornix and septum lucidum; but Dr. Greenhow does not regard these appearances as at all explaining the cerebral symptoms. Softening of the central parts of the brain, fulness of the vessels, and slight excess of serum in the ventricles, are so commonly found that, in the absence of all signs of meningitis and of microscopic evidence of inflammation in the brain, they are commonly and justly regarded as due either to *post mortem* changes, or to the obstruction of respiration which may have existed shortly before death.

I am, therefore, disposed to look on these violent and fatal symptoms as purely functional; that is, dependent on some molecular change in the grey matter of the brain, not on "metastasis", meningitis, or any inflammatory or tubercular lesion. They should take their place with the vomiting and cardiac disturbance, which are both independent of any anatomical change in the stomach and head, and are correctly looked upon as neuroses. From Addison himself to Dr. Greenhow, the best observers of the disease have attributed its characteristic symptoms not directly to the affection of the adrenals, but indirectly, through the implication of the nerves of the solar plexus. A long-continued peripheral irritation of the great branches distributed to these organs probably affects the nervous centres somewhat as they are affected in the vomiting of pregnancy, in the syncope which follows a slight blow on the epigastrium, or in the convulsions which, in some cases of epilepsy, can be traced to a distal source of irritation. A German pathologist has even supposed that all the symptoms of Addison's disease, pigmentation included, are due to a primary affection of the nervous system, including both the motor and the trophic centres, without any necessary lesion of the adrenals. But, in the first place, the single case on which this ingenious theory is based was clearly not one of Addison's disease at all, the distribution of pigment being very different; secondly, it was complicated with scleroderma; and, thirdly, the elaborate macroscopic and microscopic investigation of the organs after death is, unfortunately, deficient in one important point—the state of the spinal cord.*

It is abundantly clear that the characteristic features of the disease during life go with the characteristic lesion of the suprarenal capsules found after death. But that this connection is through the nervous system seems to be indicated by the character of the symptoms themselves as well as by the large supply of nerves and ganglia to the organs, by far the largest in the body in proportion to their size. We can scarcely expect beforehand that the quasi-tubercular lesion of the adrenals should in itself produce the symptoms of Addison's disease; for they may be destroyed by cancer or hæmorrhage, or waxy degeneration, without any apparent result. Moreover, both their histology and the history of their development lead us to regard them as obsolete and functionless organs, at least after birth. Either their use is during foetal life, when they attain their full size, or even that development must be regarded as a "recapitulation" of a lost stage in

* A prescription containing bismuth and hydrocyanic acid was found after death, which pointed to his having suffered from nausea or vomiting.

* See Virchow's *Archiv.* vol. I, p. 566.

phytogenesis. In the adult, Riolan maintained them to be "inutiles, tamquam vietas et marcidas, instar vasorum umbilicalium; adeoque usum querendum in foetu".

The unfortunate term "gland" has led some writers to imagine that a "secretion" might be retained in the blood when their supposed function is injured, and others to look for leukaemia: a condition which, I believe, never exists in Addison's disease. They are, we know, glands in no other sense than is the glans penis. They have neither the structure of a secreting gland, nor that of a lymph-gland; and, whatever the term "blood-vascular gland" may mean, they have certainly no connection in structure, physiology, or disease with the thymus, the spleen, or other lymphatic structures. Like the thyroid and the pituitary body, they are probably rudimentary organs, survivals of what once performed functions in the individual or the race. Their importance depends upon the mischief they may do through the abundant nerves and ganglia they contain. If rapidly and completely destroyed, as by cancer, or if only chemically altered, as by waxy degeneration, no symptoms ensue; but a chronic inflammatory change, leading to caseous degeneration and contraction, keeps up a continued irritation, which, sooner or later, produces the various neuroses that make up the symptoms of the disease.

CASE OF OVARIOTOMY.

BY GEORGE BUCHANAN, M.A., M.D.,

Professor of Clinical Surgery in Glasgow University.

Reported by Dr. FARQUHARSON, Coatbridge.

MRS. R. of Culhill, near Coatbridge, aged 40, had given birth to seven children; no miscarriages. Her previous health and family history were good. Three years ago, she was treated for what the doctor designated inflammation of the liver, from which she suffered very acutely. Eleven days after the commencement of the attack, she was delivered of a child; labour, she believed, being brought on by the application externally of hot fomentations to the region of the liver, violent movements of child having been felt at that time. However, she recovered from this illness completely.

In 1872, she felt a lump about the size of a fist, two inches above Poupart's ligament, on the left side. It was so movable, that she could carry it across to the corresponding place on the opposite side. For this she consulted Dr. Simpson of Edinburgh, who diagnosed that it was an ovarian tumour. It gradually increased in size, became fixed and fluctuating, and displaced the viscera. From its great bulk and weight, it was a source of great inconvenience and discomfort, and Dr. Farquharson thought it proper to tap it in the usual way on January 7th, 1873. The fluid that was drawn measured ten Scotch pints. She experienced a little faintness, but was much relieved afterwards. Again she was tapped on July 27th, when the fluid measured nine pints. During all this time menstruation was regular, though a little scanty.

Dr. Buchanan operated on November 21st. She then measured fifty-six inches round the abdomen at the umbilicus. She was very emaciated, and suffered much from the bulk of the tumour. The longitudinal incision measured nearly nine inches, through which, after separating adhesions, the tumour was extracted. It weighed 10¾ lbs., and was multilocular; the largest cyst containing fluid of a pale straw colour, the smaller ones a viscid material.

The patient did well during the operation, which occupied three-quarters of an hour, commencing at 1 P.M. on November 21st. At 3 P.M., her pulse was 109; at 3.30 P.M., 106; at 5.30 P.M., 108; at 12.45 A.M., 98; at 3.15 A.M., 108; and so it remained till the night of November 22nd.

On the night of the 21st she complained of pain in the right shoulder. At 12.30 A.M., her urine was drawn off, which measured four ounces, and was high coloured. At 5.30 A.M., she got an opiate enema, which was repeated three times during the next day. Ice only was given. She complained of little pain.

November 23rd, 10.30 P.M. Pulse 114. The urine was drawn off; it measured five ounces. The skin was cool; the tongue reddish and gently moist; thirst great. She had snatches of sleep from 3 P.M. Ice was continued. Pain was felt occasionally in the right shoulder. The left leg was powerless. She complained of pain in the bowels and round the loins from flatulence. There was no vomiting or sickness. She had lain in one position till now, when the pillows were turned. An enema of tincture of opium was repeated at 7 P.M. The patient had a cheerful look, and seemed in good spirits.

November 24th, 8 A.M. The urine was drawn off; it measured three

ounces, and was paler than previously. Menstruation had appeared. She passed some wind, which afforded much relief. She had not the slightest pain. Last night she slept three-fourths of an hour, which was the longest sleep she had had. Thirst was still great; pulse 102. She was ordered a teaspoonful of milk with ice every hour. She had a sensation of hunger to-day, having had no food or brandy from 2.30 P.M. of the 21st till 9 A.M. of November 24th. The skin was always moderately warm. There was no excitement.—2.30 P.M. Pulse 96. She passed more wind. She could move the paralysed limb freely. She complained of hunger, and was ordered a tablespoonful of milk with ice every hour. The other symptoms were unchanged. The urine measured seven ounces.—10.30 P.M. The urine measured eight ounces. She was restless; pulse 114; no fever. She was still hungry. The flatulency was annoying, otherwise she was well and cheerful.

November 25th, 8 A.M. She slept one hour last night, which was the longest sleep as yet. A drachm of tincture of opium was given at bedtime. The urine measured nine ounces. She was still hungry. There was a slight smell from the wound. Skin natural; pulse 90.—5 P.M. Urine twelve ounces; pulse 96. She was ordered a tablespoonful of chicken-soup every hour.—12 P.M. Urine eight ounces; pulse 96. The opiate was repeated. She had no pain, but felt rather chilly.

November 26th, 7 A.M. Urine eight ounces.—2 P.M. Urine seven ounces; pulse 80. The wound was dressed; it had a slight smell. She was still hungry. She passed more wind. There was a tingling felt in the left leg. She felt happy and comfortable. The wound looked well. There was very little grumous pus. Slight tenderness was felt over the left iliac region.—10.30 P.M. Urine eleven ounces; pulse 78. Milk and soup were continued, with ice. The opiate was repeated. She was very anxious for food. The symptoms were unchanged.

November 27th, 1.30 P.M. Pulse 90. Iced milk was given every hour; and she had a cup of beef-tea with a half slice of bread toasted. Tea and toast at 5 P.M.; after which she slept for half an hour, then feeling very comfortable.—10 P.M. She had no pain; tongue clean. There was no thirst to-day. She had improved much on beef-tea; and was ordered porridge and milk in the morning. The urine was natural.

November 28th, 1.30 P.M. She enjoyed the porridge, which was followed by a cup of tea and a little toast an hour afterwards. Beef-tea and toast were given at 2 P.M., and tea and toast in the evening, milk and ice being given between. Pulse 84. She felt stronger, very well and comfortable. The urine still required to be drawn off, and was quite natural.

November 29th, 1.30 P.M. The ice was stopped; pulse 84. She felt stronger. The food was similar to yesterday's, only increased in quantity. The urine was the same.

November 30th, 1 P.M. Pulse 90. She still enjoyed food, and felt well. She passed urine herself to-day for the first time; it measured four ounces. The wound was dressed; a small quantity of dark fluid was found around the clamp. The wound was healed; the edges at the clamp were healthy and granulating. Carbolic oil dressing was used. The fluid evidently originated from the decaying pedicle. The appearance of the abdomen was natural, except that the part was drawn in where the clamp was attached. She had now no pain on being lifted. The tongue was slightly furred. The food was as yesterday, but increased in quantity. She had no thirst.

December 1st, 2 P.M. Urine normal; pulse 90. The wound was dressed. The discharge around the clamp was very slight. The patient felt very well. She was ordered a mutton-chop for dinner.

December 3rd. There was slight pain in the wound, but she felt stronger.

December 4th. The bowels were moved freely. She had slight tenesmus afterwards.

December 5th. The tenesmus continued. She was ordered a drachm of castor-oil. There was very little discharge from the clamp, but a little more pain when touched.

December 6th. The patient was quite well to-day. The bowels were moved without oil; pulse 84. The wound, which was healthy, was dressed. She felt pain on lying on the left side.

December 7th. The dose of castor-oil was repeated; the bowels were freely moved; she felt much lighter; pulse 96, a little weak. Her appetite was good; she slept well; she moved in bed with considerable freedom.

December 8th. The clamp came away; a circular sore was left, fully half an inch in diameter, healthy. The retraction of the abdomen was not so great. Pulse 96. She did not require an opiate last night. She slept well, and felt very hungry in the morning.

December 9th. The wound was dressed; it looked healthy; there was very little pus. The bowels were moved in the morning. She was taking more solid food.