the involvement of the hepatic region and subsequent death of the patient (Fig. 17).

Pins in the appendix have been noted by many surgeons, and may cause pericolicitis. In a collection of 63 published cases R. H. Fowler refers to a case of W. W. Keen, in which the tip of the appendix was solidly incorporated in the bladder. At the age of 7 the patient had dysuria, and a pin, which he believed he had swallowed, was removed from the urethra.¹³

Pericolic adhesions may cause intestinal obstruction, as in a case recently operated upon at midnight, in which a coil of small intestine had passed under a band of such origin.

CHOLECYSTITIS WITH PERICOLITIS.

Inflammation of the gall bladder, leading to localized peritonitis in its neighbourhood, is a common cause of pericolicitis in the hepatic region, and is invariable in severe or long-standing cases. The following will serve as an example of pericolicitis following cholecystitis of pregnancy:

The last child was 7 years old, and the pain, of five months' duration, was first noticed in the epigastrium and passed round to the back. It was of a severe shooting character, worse at night and on walking, doubling her up, and relieved by the vomiting which always occurred. She had had indigestion for two years. There was tenderness in the epigastrium, not over the gall bladder, which could not be felt. At the operation the gall bladder was thickened and mottled, but no stone was present in it or in the ducts. Marked pericolicitis was present, the ascending colon being adherent to the anterior abdominal wall and the liver adherent to the gastro-hepatic omentum. In particular there was a strong band of omentum adherent to the lumbar peritoneum (Fig. 18).

It will be seen, therefore, that pericolicitis is a frequent accompaniment of inflammatory affections of the appendix and gall bladder. Its persistence as bands and adhesions may give rise to symptoms after the successful treatment of the original focus, particularly in appendicitis. In wandering kidney it is now demonstrated that pericolicitis may coexist with encysted intraperitoneal collections of fluid. In the surgery of the right loin more attention will doubtless be paid to this associated condition by more complete examination of the region at the time of operation; and, by extended observation, the pathology of vague pains and after-pains will be more fully understood.

REFERENCES.

- ¹ Francke: Coli-infektion der Harnwege. *Mitteilungen aus den Grenzgebieten der Med. und Chir.*, xxii, No. 4, 511-674.
- ² Glantenay et Gosset: Le Fascia Péri-rénal. *Annales des maladies des organes génito-urinaires*, 1898, p. 118 et seq.
- ³ Morris, H.: *Surgical Diseases of Kidney and Ureter*, 1901, i, pp. 103, 104.
- ⁴ Carwardine: *Bristol Medico-Chirurgical Journal*, 1905, xxiii, p. 24.
- ⁵ Jackson, Jabez: Membranous Pericolicitis. *Transactions of the Western Surgical and Gynecological Association*, 1908; also *Surgery, Gynaecology, and Obstetrics*, 1909, ix, 278.
- ⁶ Sims, Parker: *Transactions of the New York Surgical Society*, February, 1912.
- ⁷ Lane: *Surgery, Gynaecology, and Obstetrics*, 1908, vi, 116.
- ⁸ Lane: *BRITISH MEDICAL JOURNAL*, 1911, ii, 914.
- ⁹ Mayo, C. H.: *Collected Papers*, vol. ii, 1911, p. 221.
- ¹⁰ Gerster: *Annals of Surgery*, 1911, ii, p. 325.
- ¹¹ Pilcher: *Ibid.*, 1912, i, p. 3.
- ¹² Stanton: *Ibid.*, 1911, i, p. 813.
- ¹³ Fowler, R. H.: *Ibid.*, 1912, ii, p. 433.

THE bequests made by Miss Alphonsine Sarah Jarry, of Settle, Yorkshire, included one of £600 to Halifax Infirmary.

WOLBACH and TODD, in the Third Report of the Expedition of the Liverpool School of Tropical Medicine to the Gambia in 1911, state that they have established the presence throughout the Gambia of that form of ulceration occurring in the tropics which is associated with the presence of spirochaetes and fusiform bacilli. In 17 cases the pathological history of the ulcers was studied; although they varied greatly in their gross appearances, their histological features had so much in common that they could be regarded as having a common etiology. The conditions observed excluded as etiological factors, syphilis, varices, acid-fast bacteria, blastomycetes, Leishmania, actinomycetes, and the streptothrices. The presence of spirochaetes in nine of their cases lends support to the view that they are the cause of the process. These spirochaetes corresponded accurately to the parasites described by v. Prowazek, Keysseltz and Mayer, Balfour, and other authors, and named by v. Prowazek *Spirochaeta schaudinni*. Subcutaneous inoculations made by the authors into white rats gave negative results.

A Lecture

ON

SOME POINTS ABOUT DRAINAGE OF THE ABDOMEN UNDER VARIOUS CONDITIONS OF INFLAMMATION.

By ARTHUR E. BARKER, F.R.C.S.,

SURGEON TO UNIVERSITY COLLEGE HOSPITAL, AND CONSULTING SURGEON TO THE QUEEN ALEXANDRA MILITARY HOSPITAL, MILLBANK.

THE subject chosen is one pre-eminently difficult to handle. Anyone dealing with it even in the most detached way is very liable to be misunderstood. I hope I shall be able to avoid this in the present instance.

I want to speculate from evidence before us as to whether the current views on drainage of the abdomen in inflammatory conditions may not in some cases be modified with advantage to-day, as they have required to be in the past.

I need hardly remind a senior audience that not so long ago stirring debates took place in our societies as to the employment of drainage of the abdomen in such simple non-inflammatory conditions as early and uncomplicated ovariectomies. Yet now such a measure is rarely if ever adopted; the wound is at once sewn up. The reason is obvious. Any effusion which follows such an operation nowadays is either absolutely aseptic or the few organisms which may have gained access to it from without are easily disposed of by the germicidal properties of the fresh exudation with its phagocytes, and are absorbed by the extensive lymph sac which the peritoneal sac may be described to be. The effect of discarding drainage in such simple aseptic cases was, of course, to reduce the number of weak scars and consequent ventral herniae, and to diminish the liability to subsequent dangerous adhesions about the track of the drain. A painful after-treatment was also avoided, as was also a great loss of fresh healthy serum which otherwise would be returned by the lymphatics to the system with benefit.

Appendicitis.

Let me now ask you to recall the earlier medical experiences of appendicitis before the operative treatment of the affection became established. From these it soon became clear that many acute cases could recover without opening the abdomen, and without any escape of pus or other effusion either externally or into the intestines. It was a fair deduction from such experiences that the peritoneal cavity could dispose of a large amount of septic inflammatory matter. Otherwise none of these cases could have recovered.

Later on it was recognized that in many cases the amount of septic matter was too great to be left entirely to the resources of nature, and abscesses were encouraged to burst under fomentations, or operations were undertaken to evacuate them. But even here much was left behind which was not carried off by drainage, and could only be destroyed on the spot. From our experience of these operations we now know that ventral herniae and kinks of the intestine were not uncommon sequelae. But the operations on these very ventral herniae themselves following upon extensive suppuration within the abdomen have taught us, perhaps, more than anything else of the marvellous powers the peritoneum possesses of arresting and repairing the damage done. We have undertaken a radical cure, perhaps, a year or so after we had actually seen all the coils of bowel in the right flank matted together by foul lymph, and have found in many cases no traces of adhesions and all the peritoneal surfaces polished and as healthy looking as if they had never been involved in so dangerous a septic inflammation. Another point must have struck many surgeons after making long incisions to evacuate appendiceal abscesses. The ends of the incisions have been stitched up, leaving a drain in the centre, and we have seen the ends heal practically by first intention, in spite of being bathed in pus under a fomentation for days in many cases.

* Delivered at the Medical Graduates' College.

From all this it has seemed to many surgeons, both at home and abroad, that perhaps we have been too wholesale in routine drainage when dealing with inflamed appendices, and that more discrimination is desirable. In other words that, after the removal of the inflamed organ and the septic products around it as far as possible by careful mopping, we might perhaps with advantage in some cases stitch up the abdominal opening after cleaning its edges, and obtain primary union with all its advantages. Here, of course, some traces of the septic matter would be left behind within the abdomen; but this, as we have seen from our experience noted above, can be disposed of by the defensive powers of the peritoneum.

Those who have taken this view can now point to numerous cases of acute epityphlitis in which inflamed and even gangrenous appendices surrounded by septic effusion have been dealt with and which have yet healed *per primam* after careful suture. In some of these cases the septic fluid from within has obviously flowed over the edges of the wound during the operative manipulations, and the inflamed lymph-covered organ has perhaps been in contact with them, and yet the result has been as perfect as if this had not been the case. Of course there remain a large number of cases in which thorough and prolonged drainage appears absolutely called for by common consent. This is our great difficulty, only to be overcome by experience. But it appears equally obvious that there are many in which drainage can be dispensed with to the great advantage of the patient, immediate and remote.

So far, then, as can be seen from clinical and experimental observation, a very considerable amount of septic matter is tolerated by the peritoneal surfaces, and can be prevented by them from doing irremediable injury when not in too great amount or overpoweringly virulent. That the virulence of the particular infection affects the issue is, of course, generally known. But that this general knowledge does not sufficiently affect our practice is what I am trying to bring out. We must all realize that every case of acute appendicitis is not the same affection, though the same organ is attacked in all. Let us take examples.

One patient is seized, and within a few hours has all the symptoms of profound toxæmia, which may be present to a degree almost precluding hope of recovery, with or without operation. In his case, if an operation is done, but little is found but a perforated vermiform appendix and a small quantity of a thin, greasy exudation from all the deeply injected peritoneal surfaces in view, but without a trace of plastic lymph. The intestines are distended, and more or less parased. All we can do here, if we operate at all, is to remove the appendix, and with it a source of further infection. But the toxic infection has got a firm hold, and the natural forces alone can arrest it, aided, perhaps, by measures to be alluded to presently.

Another patient has an apparently similar attack; the same time has elapsed, but the general condition in this case is fairly good; there are, in other words, no marked symptoms of toxæmia, or very few. But if we open the abdomen we find what might at first sight be taken for a worse local condition. There is an inflamed appendix, as in the first instance, and the changes around it look more severe. But they are localized. There is much plastic lymph about, and the intestines in view are little, if at all, injected, parased, or dilated. These are the cases which used to, and still do, get well with and sometimes without operation.

Now, these two conditions belong to entirely different categories. They are almost different diseases. In the first death will frequently follow, in spite of any operation which can be done for the local catastrophe. The general infection is too profound. The initial dose of poison has multiplied and saturated the whole system, either because the micrococcal strain was more than usually virulent, or because of a low standard of resistance in the individual.

In the second condition the virulence of the cocci is lower, is controlled by the greater vitality of the patient, and their effects are expended locally. It is unnecessary to refer to the particular cocci concerned in both processes. They may possibly be the same morphologically, but of a different strain as regards virulence.

Such differences in the cases we have to treat are seen daily. Is it rational that they should all be dealt with in

the same stereotyped way when we come to operate on them? I think not, and an ever-increasing number of operators are of the same opinion, to judge from the literature of appendicitis, both home and foreign. It must not be forgotten, too, that there are many degrees of severity between the two extremes just mentioned. And herein lies our difficulty—a difficulty which has led to a too uniform method of procedure. But we are growing every day in knowledge of the vital processes in action, and ought to be, and are, better able to discriminate as to the aims of operative procedure to be applied to each variety. We have come so far as to realize that the milder attacks may be left until they have passed off, and then operated on for the removal of the appendix without drainage. We are coming to know what cases of suppuration must be drained. We are getting to believe that some cases with actual pus formation need not be drained, and acting on this belief, have obtained results which contrast very favourably with those treated by prolonged and often troublesome drainage.

The question is, Can we, by careful clinical study, determine during an operation whether an acute appendicitis ought to be drained or not? I think such a thing is possible. Let us take a case or two by way of illustration. It is quite clear that in all so-called "interval cases," or *appendicite à froid*, no drainage is required. We close the wound after removal of the organ, and expect and obtain primary union. But on a recently inflamed organ we often find much discoloured and foul lymph, with adhesions to surrounding parts, and not infrequently more or less turbid fluid. Can we venture in such a case to close our wound after removal of the organ and careful mopping with gauze, and dispense altogether with drainage? Experience shows that we certainly can in a large proportion of cases, and my own lists show that this practice is increasingly common and successful. And yet, when we come to think of it, these conditions are septic, and it can hardly be claimed that the removal of the inflamed organ can be accomplished without more or less contamination of our incision edges. Why, then, do these adhere and unite faultlessly, as they certainly do in many cases? This can be explained in several ways: Either the inflammatory exudation on and around the vermiform appendix was practically sterile, or, if not, the amount of organisms already inhibited by the action of effused serum was not enough to overpower the active tissue cells of the freshly made wound and the plastic lymph which repairs it. In some cases we have found that the effused fluid in such cases poured out in response to the infection from the inflamed vermiform appendix is actually sterile. No organisms have been grown from it on the ordinary culture media. The appendix itself was undoubtedly infective, but the serous fluid effused around it was probably prophylactic, as we know it to be elsewhere, and inhibited the development of the organisms. It was not to be wondered at that here the wounds healed primarily. But what about other cases where there has been a small abscess round a putrid appendix? Here, too, we have had in some cases the best results from immediate closure of the wound. We must suppose, then, that though some of the pus did undoubtedly reach the edges of the wound, the same inhibiting forces had affected the cocci in it, and that they were overpowered by the patient's own living cells in the wound, just as part of a wound which is being drained, and is consequently covered by pus, may heal up *per primam*.

In some few instances we have underestimated the chances of infection, and some suppuration, usually very slight, has taken place in our closed wound. But it is notable that the peritoneal conditions have in no case been unfavourably influenced by this. When the wound has shown blemishes it has apparently been due to the accumulation between the skin and the deeper stitches of some blood clot which has proved a suitable soil for the growth of organisms. Usually the escape of a little turbid serum through the stitch holes or of broken-down blood clot from a corner of the wound has been followed by sound union, as though nothing had happened. In some very few cases the insertion of a subcutaneous drain has been necessary, and a stitch or two have worked out. But in the end union has taken place with apparently no more loss of time than had the wound been drained from

the first. Such blemishes will be inevitable occasionally until we have accumulated experience enough to know how far we may go with non-drainage, and what cases must have a drain from the first. So long as the intra-peritoneal conditions are not unfavourably influenced by non-drainage, the superficial processes are of secondary importance. Of course, if we follow the routine of draining all inflammatory intra-abdominal processes without exception, we escape the need of discrimination. But in this case there will be many more weak scars and ventral herniae to be dealt with subsequently, and probably more dangers from adhesions and infection from without.

If we endeavour to lay down formulae for our general guidance in such cases, they would be somewhat as follows: Given a case in which on opening the abdomen an inflamed vermiform appendix is found, around which there is a moderate amount of obviously plastic lymph shutting off the process, whether it be suppurative or not, and if outside this the abdomen contain a certain amount of clear effusion and the intestines in view are normal in appearance, if at the same time the patient's condition is good, his colour fresh and his pulse moderate in frequency and in quality, and if the offending organ can be removed with comparative ease, I am inclined to close the wound carefully without drainage. The clear fluid around the immediate area of trouble is probably prophylactic, and will often be found to be sterile.

But if, on the other hand, on opening the abdomen, a rotting or inflamed appendix be found with little or no plastic lymph anywhere, and the intestines in view are red, injected, and dilated from paresis and covered by a greasy exudation or immersed in a thin ichorous, discoloured effusion; and if this picture is associated with feeble, rapid pulse, great asthenia, loss of colour, perhaps frequent vomiting—in short, with clear signs of toxæmia—it is better to drain freely. The cocci in this case are particularly virulent and have led to a completely aplastic peritonitis, with toxæmia and consequent loss of general vitality. All that we can do here is to remove the exciting cause in the form of the peccant appendix (if this can be done without prolonged operation) and keep the wound open in the hope that some of the toxins, at all events, may escape from the peritoneal cavity by drainage. We are dealing here with a profound general poisoning, and the prognosis is extremely grave. Such cases must be spared from all prolonged exposure to the depressing effects of manipulations and the anaesthetic. In most all search for the appendix should be left for a subsequent operation unless it present itself at once. From evidence before us it appears more than probable that the effect of chloroform plays a large part in the general depression.

While writing the above I have seen 7 cases which illustrate the varieties alluded to on which I have operated with the last few days:

1. A mild quiescent case; appendix removed; no drainage.
2. A mild active case; appendix removed; no drainage.
3. A severe general case with much effused aplastic fluid in the right flank reaching to the diaphragm and producing cough, but with only moderate toxæmia and a relatively fair pulse; appendix removed; free drainage.
4. Severe inflammation with sloughing appendix and abscess, but general condition fairly good—organ removed; free drainage.
5. Severe attack with localized abscess and appendix imbedded in wall of caecum. Temperature high. Organ dissected out; drainage.
6. Intense inflammation; appendix imbedded in caecal wall and acute enlargement of mesenteric glands in the angle between the ileum and colon; caput coli and last two inches of ileum solid with oedema; appendix removed, no drainage. The patient had incipient pneumonia with much herpes on cheek and lips. Temperature very high before operation.
7. Intense inflammation of distal third of appendix with pus just about to burst through its serous coverings in many white patches; vascular adhesions all round; organ removed; wound closed without drain.

All 7 cases recovered. Of the 6 with fresh acute inflammation 3 were drained, and are practically quite well; 3 were stitched up without any drain, 2 of these healed without a flaw, one showed a perfectly healed skin incision with a little turbid serum from two stitch holes. A few days later: The wound is now dry, and has all the signs of primary union, being firm and soft. One of the cases (No. 4) in which drainage was employed would, I now think, have done equally well or better without drainage. Not so long ago all, except the first, would have

been treated with drainage, and would probably ultimately have developed ventral herniae.

PERFORATION OF STOMACH AND DUODENUM.

This same question—whether we ought to drain or not—confronts us after operations for several other acute inflammatory conditions within the abdomen. Nowhere is it more momentous than after perforation of the stomach or duodenum. Here after we have closed the perforation it is difficult in some cases to say whether drainage should follow or not. I have treated many in both ways, and have gained the impression that those whose wounds have been closed at once have suffered less from the catastrophe itself and the operation than those whose wounds have been drained. I do not think I have ever regretted not having drained. I have often concluded that drainage had been of no use if not actually injurious. One thing is, I think, pretty clear, namely, that if a drain be used the escape of fluid is only from the abdomen in the immediate neighbourhood of our wound. No effectual means of draining the abdomen have yet been devised. We ask ourselves, then, Is it worth while to drain an area not larger than could be covered by the closed hand, and probably for not longer than 24 hours? For by this time the field of operation will be shut off from the rest of the abdomen by plastic exudation. Indeed, I have often speculated whether, in some cases at least, our drain has not prevented the completion of this plastic process, and interfered with the apposition and union of the damaged viscus with the parietal peritoneum and other parts around. And if an aplastic peritonitis is the sequel to a perforation the fluid effused will gravitate into the flanks, and here it could not be withdrawn by any drain in the epigastric incision. Fresh wounds in the flanks might be made, but they are, as a rule, unsatisfactory in these cases. Indeed, a patient with aplastic diffuse peritonitis after perforation is in a desperate condition from which no amount of drainage will save him, and each fresh opening in the abdomen weakens him. His own powers of resistance alone can save him. I have seen collections in the flanks and pelvis appear and disappear in these cases without incisions.

From all these considerations I have been led for several years considerably to modify some older views about drainage in such cases. In many of them after careful mopping of the infected surfaces, however extensive, the wound in the abdominal wall has been completely closed in many cases. And I cannot recall one in which there has been reason to regret the practice. There is good ground for believing, as already stated, that fresh serum in the peritoneum and other serous cavities is germicidal, and further, dilutes the toxins produced by the bacteria. It is effused rapidly, and quickly absorbed, carrying the toxins with it, to be eliminated by the kidneys. The leucocytosis which is at the same time induced, acts further in destroying the septic germs, and between these two processes the infection is controlled if only the patient's general vitality is maintained. We believe we can assist in the cycle of events in several ways. For instance, copious subcutaneous or intravenous infusions of normal saline solution, or of 5 per cent. of glucose in distilled water, not only help to raise the vascular tension lowered by sweating, vomiting and toxæmia, but also further the dilution and elimination of the toxins by the kidneys; moreover, leucocytosis is said to be increased by these means.

The same processes for the destruction of bacteria are seen in operation in other analogous cavities of the body, notably the joints and spinal canal when infected through the blood stream. A joint thus recently infected may be found to swarm with bacteria. Subsequent aspirations may show—except in the most virulent infections, usually from without—an ever diminishing number of organisms, until at last they cannot be found either in smear preparations or by cultivation. The same is seen in cases of spinal meningitis. I have had several which have recovered perfectly. Two are recorded (and many more collected from other publications) in the *Transactions of the Royal Society of Medicine*, 1907, where suppurative meningitis of the dura, producing almost hopeless general conditions, recovered under no other treatment than repeated withdrawal of the highly infected fluid by repeated tapping of the lumbar sac.

There is much evidence, then, for the belief that the serous cavities of the body, being so important, are provided with specially powerful defensive works to protect them from bacterial invasion. It should be our aim to assist them surgically when they are in danger of being overpowered by the accumulation of bacteria and their toxins—sometimes by evacuation with drainage, sometimes by evacuation without drainage. We should always study to interfere as little as possible with the natural protective agencies. Overhandling, overdisturbance, overstimulation with chemical germicides, have long been recognized as harmful, and I feel sure that over-free use of drainage also often plays an injurious rôle. The mere fact that a wound in the epigastrium is kept open by the insertion of foreign material must more or less disturb the vascular balance within the abdomen; and the presence here of an open wound is more painful during the first few critical days than a closed one, on which the movements of respiration have less effect. The daily readjustment, too, of drains and dressings must cause a great addition of pain with all its depressing influences at the patient's weakest time, besides increasing the risk of added sepsis. I do not think there can be any question that a patient's general condition is more satisfactory with a closed wound healing more or less by first intention under a dressing untouched for days than with an open wound.

Of course an attempt to obtain primary union may not be quite successful in some cases. Invisible septic particles may have lodged in the wound and may infect the deeper or superficial stitches, and some of these may suppurate. But this, if it occur at all, will take place after some days, during which the peritoneal cavity has, at all events, been at rest. At the worst in such cases some of the deep stitches have worked out without delaying the general recovery longer, perhaps, than had the wound been drained. This is a small price to pay for getting the abdomen closed at once and giving the patient rest for the first few days while the catastrophe within is repaired. As a matter of fact, the number of cases of perforated gastric and duodenal ulcers treated by immediate closure of the abdominal wound, treated in my beds by myself and others, has largely increased, and primary union has been the result in an ever larger proportion of late. Some will continue to be drained after such lesions, but I think fewer and fewer.

Here, again, the character of the fluid met with when the abdomen is opened will help to decide the point in question. If much putrid food have escaped from the perforation and the resulting peritonitis has yielded an acrid acid fluid as tested at once by litmus paper, the case is a bad one. If, besides, there is deep injection of the intestines with distension, and little or no plastic lymph is seen, we may count on a virulent form of infection, which would inoculate our wound at once and prevent its healing. Here I should drain freely, but with small hope for the patient.

But in some cases of fasting stomach there is little extravasation through the perforation. Here we find a bland whey-like alkaline or neutral fluid with no deep injection or distension of the intestines, while they show an inclination to adhere by plastic lymph. In these cases, after careful cleansing of all the pockets of the peritoneum accessible to swabs on holders, I am now inclined to close the abdomen completely without drainage. Many such cases have healed without further trouble by primary union. For instance, in a case of duodenal perforation the diagnosis was very obscure, and an opening was made over the vermiform appendix, which was under suspicion. The whole peritoneal cavity was full of the bland fluid alluded to, and there was some free gas. Recognizing the true condition now, I opened in the epigastrium, found the perforation near the pylorus, and closed it. After thorough swabbing of the whole peritoneal cavity I stitched up both wounds without drainage, and permanent primary union without a flaw was the result, the patient being soon well.

PYOSALPINX.

Let us take now another acute abdominal condition. If on opening the abdomen a pyosalpinx with localized peritonitis is found, there need be less hesitation. Here the immediate closure of the wound may be done in the large majority of cases. In the first place, it is known to

bacteriologists that the pus from pyosalpinx is sterile in about 50 per cent. of ordinary cases. In this group there is no special reason for drainage if oozing from adhesions can be quite controlled. But in the other half of the cases the virulence of the bacteria mitigated by long residence in the body is low, as indicated by the plastic exudation and the absence of inflamed and paresed intestines, and a carefully sutured wound will heal *per primam* in the majority of cases in spite of the pus. I have seen this over and over again, and also after ruptured tubal pregnancy.

INTESTINAL GANGRENE.

Again, in another class of lesions the same question has to be considered. I allude to gangrene of the gut from internal strangulation. Here, except where there has been perforation and faecal extravasation, drainage should be avoided after removal of the damaged portion. The plastic processes appear to me to be more perfect without a drain, which may prevent the apposition of healthy structures around our line of suture. I could produce many instances to illustrate this point; one or two will suffice. A few years ago I had to remove four feet of small intestine black and gangrenous from constriction of an omental band. On opening the abdomen, over a pint of dark putrid flocculent fluid escaped. After the excision of the gangrenous coils and junction of the ends, the wound was completely sutured without drainage, and primary union followed. Her doctor tells me the lady has remained perfectly well ever since, with no signs of adhesions. Another similar case occurred to me in hospital a few years earlier. Here I removed 5½ ft. of small gut for gangrene with a couple of pints of dark putrid fluid. Perfect primary union followed without drainage. The patient was a feeble woman of 76, and was quite well when I last saw her two years after operation. In both cases the putrid fluid, one would have thought, ought to have interfered with primary union, but it did not.

In cases of removal of the caecum where we open up areolar planes it may be a different matter. Drainage may perhaps be required unless we are perfectly certain of our asepsis in removing the gut. To deal with more or less sepsis in the peritoneal cavity is quite another thing to arresting its progress in areolar tissue. But even here drainage may be discarded in many cases, as I know.

I have asked your consideration of these points and given you some of my experiences in the hope of drawing your attention to a subject upon which our conclusions are not yet fully crystallized and in an attempt to arouse a more careful clinical study which must always go hand in hand with bacteriological work. I am addressing experienced practitioners, not young student beginners who might be likely to draw wrong conclusions from the speculations offered them. These are of no recent growth. It must be more than twelve years ago since I ventured to publish a paper on this subject. And experience, in many cases combined with what we have learnt from bacteriology, has served to confirm and amplify the deductions then put forward.

THE Royal Dental Hospital, Leicester Square, has received from the executors of the late Captain T. B. Heathorn a donation of £400.

THE Association of Infant Consultations and Schools for Mothers, which is now doing its work as a department of the National League for Physical Education and Improvement (4, Tavistock Square, London, W.C.), has in view a competition in the subject of mothercraft, the competitors being women attending the various schools for mothers in London. It is proposed that each of the schools shall select by competition not more than six representatives, and that these shall subsequently take part in a competition held by the association itself for prizes in nine classes of work. Among the latter are dressing and undressing a baby, laundry work, cooking, cutting out and making garments, knitting, and cleansing babies' bottles. A prize will also be given for the baby evidencing the best condition and greatest care, provided that it is shown by a mother in possession of a favourable report as to her home for a period of not less than three months signed by a health visitor or other responsible person. The competition is to take place during the health week, April 6th to April 12th.