It has been suggested that local recurrence after local excision is strong evidence in favour of the presence of intramural permeation, even though such be undemonstrable.

That a typical carcinomatous ulcer may be formed as a result of invasion of the bowel wall from without is demonstrated by a case in this series.

On opening the abdomen, it was noted that the pelvic colon was densely adherent to the anterior and left lateral surface of the rectum at the bottom of Douglas's pouch. Separation from the uterus was regarded as an unsafe procedure, so this organ was also removed. On examination of the 12 in. of excised bowel the presence of two growths was revealed, though only one had been diagnosed. These were located on the bowel walls at the site of adhesion. The upper growth, in the pelvic colon, was of the whipcord type, and had completely encircled and constricted the bowel wall, giving rise to the symptoms of obstruction for which the patient was admitted to hospital. The lower growth, in the rectum, presented itself as a carcinomatous ulcer involving the anterior and left lateral walls of the bowel. It is difficult to escape the conclusion that the upper was the older growth and that the lower one had arisen as a result of contact.

In Fig. 7 is depicted a carcinomatous deposit in the pericolic connective tissue 3 in. above the upper margin of a growth. If in such a case a local excision had been practised with the removal of 1 in. of normal bowel above and below the growth, this deposit must necessarily have been left in close proximity to the intestinal junction. Local recurrence in such a case is readily explained. Clinical and *post-mortem* evidence also favours the view that recurrence is an extramural phenomenon.

Finally, it may be stated that in every case in the series the growth had spread to the tissues outside the bowel wall.

BIBLIOGRAPHY.

- Cole: The Muci-Carmin Staining Method, *Trans. Roy. Soc. Med.* (Path. Sec.), vol. v, 1911.
Cripps, Harrison: The Surgical Treatment of Rectal Cancer, *BRITISH MEDICAL JOURNAL*, October 5th, 1912.
Handley, Sampson: Recent Advances in the Surgical Treatment of Some Forms of Cancer, *Universal Medical Record*, 1912 (v).
Miles: The Treatment of Carcinoma of the Rectum and Pelvic Colon, *Glasgow Medical Journal*, February, 1912.
Paul: Address in Surgery, *BRITISH MEDICAL JOURNAL*, July 27th, 1912.

DISLOCATION OF THE METATARSUS, WITH AN ACCOUNT OF THREE CASES.

[WITH SPECIAL PLATE.]

By ARCHIBALD YOUNG, B.Sc., M.B., C.M.,

CONSULTING AND VISITING SURGEON, BROADSTONE JUBILEE HOSPITAL,
PORT GLASGOW; ASSISTANT SURGEON, WESTERN INFIRMARY,
GLASGOW; LATE SENIOR ASSISTANT TO THE REGIUS
PROFESSOR OF SURGERY, UNIVERSITY OF
GLASGOW, ETC.

FREQUENCY OF THE CONDITION.

DISLOCATION at the Lisfranc (tarsal-metatarsal) joint is an unusual form of injury, but it is probably much less rare than would be suggested by a consideration of the meagre literature on the subject. The comparative dearth of references in surgical literature may be realized when the figures given by Stimson in his treatise on fractures and dislocations are quoted:¹ "Malgaigne collected 21 cases of the various dislocations, Hitzig 29, Bayer 68, and Fischer 18 of the first metatarsal." Panse, quoted by Nasse and Borchardt, in the *Handbuch der praktischen Chirurgie* (Bergmann, Bruns, and Mikulicz),² "was able to collect, in the year 1897, 22 complete and 23 partial dislocations."

Considering the exposure of the foot to violence, these figures are undoubtedly small, and give almost certainly a false idea of the actual frequency of such a type of injury.

DEARTH OF RECORDS.

The relative dearth of records may in part be explained by the fact that minor degrees of displacement have been overlooked. The development of radiography in recent years ought to act now as an effective preventive of this type of error. Part explanation may also be found in the fact that the existence of these dislocations might readily be overlooked when they form mere incidents in a more severe injury to the foot and ankle, perhaps calling for immediate amputation.

Indeed, another aspect of this possibility is suggested by the fact that several writers of repute have put it on record

that these dislocations, when they do occur, are usually the result of crushing force of such severity as to destroy the vitality of the soft parts to a degree which renders amputation necessary.

Thus Griffith³ says they are usually "complicated by destruction of the soft parts, and frequently require amputation"; Wharton⁴ says, "These dislocations usually result from crushing forces which destroy the vitality of the soft parts so completely that amputation is required"; and even Erichsen⁵ states, "The question of amputation will always present itself in these cases, and must be determined on general principles, by the age of the patient, and by the extent of injury to the soft parts."

It is probable that such dislocations are neither so uncommon, nor, as regards the prognosis, of so grave a nature as the records and opinions referred to above would imply.

The varieties of dislocation are as follows:

1. Dislocations of single metatarsal bones.
2. Dislocations of several metatarsal bones simultaneously.
3. Dislocations of the whole metatarsus *en bloc*.

Any of these forms may be, and the second and third usually are, complicated by other injury, such as fracture, accompanying damage to the cuneiform bones or cuboid, with or without displacement, or damage to soft parts.

In Group 1 the first metatarsal seems to be by far the most commonly dislocated. In the cases reported it was almost always displaced upwards. As already quoted, Fischer⁶ was able to collect records of 18 cases of this injury. The recorded cases of isolated dislocation of the other metatarsals are very few in number.

In Group 2 cases have been recorded of quite a number of different combinations. In groups of two at a time there are records of the first and second being displaced together; also the third and fourth, and the fourth and fifth. The direction of displacement varied in each—sometimes upwards and inwards, at other times upwards and backwards. In groups of three at a time, there are records of 2 cases of dislocation of the first, second, and third metatarsals, these being displaced together upon the dorsum; in 1 case the displacement was downwards into the sole. The second, third, and fourth have been, in one or two cases, displaced together upon the dorsum.

Of dislocations of four metatarsals at a time, 4 cases are mentioned by Stimson, all having been dislocations of the first four metatarsals. In two of these the affected bones were displaced bodily upwards. In 1 case (reported by Malgaigne) the first three were displaced downward, and the fourth upward. In one (reported by Ely⁷) the first metatarsal was displaced inwards, the others upwards and outwards. This case of Ely's, it will be seen later on, seems to have been very similar to one of the three cases which I have to report in this paper.

Group 3 comprises those cases in which all the metatarsal bones have been displaced bodily at the tarsal-metatarsal joint.

Four forms have been described, according as the displacement was upward, outward, downward, or inward. Only one example of each of the two last-mentioned forms has been reported, and one, the inward form, is not free from doubt as to its correct description. The case of downward displacement was reported by Smyly.⁸ The injury was caused by the fall of a wagon which pressed the heel forward, while the toes were fixed.

Even the upward and outward displacements have not been met with, or recognized, with any degree of frequency. Hitzig, quoted by Stimson, gives the number of cases of upward dislocation he was able to collect as 11, the most usual cause being direct violence, but in 2 cases muscular violence was the agent, the patients sustaining the injury in making violent effort "to avoid falling, after having slipped while carrying heavy bundles."

This form of dislocation may be complete or incomplete, and it may be complicated by a part or the whole of one or more of the cuneiforms being displaced along with the metatarsals. Examination has shown in those cases examined *post mortem*, or where the injury was compound, just such damage—in the way of rupture of dorsal and plantar ligaments and of muscles, in the fracture or separation of cuboid and cuneiform bones, and in the fracture of one or more of the displaced metatarsals, or

their separation from each other in a lateral direction—as one might expect such an injury would readily produce or imply. The resultant clinical picture has varied a good deal in the different cases, the normal axis of tarsus and metatarsus being in some cases preserved, in others the metatarsus being displaced laterally *en bloc*, so leading to an angular deformity, in addition to the foreshortening of the foot. The proximal ends of the metatarsals bulge on the dorsum, forming a ridge corresponding roughly to the line of the joints, or proximal to this on the dorsum of the tarsus. In some of the recorded cases reduction was successfully accomplished. Even in the cases where it failed fairly good function was obtained in time. In only one of the cases was amputation done as a primary expedient.

Of the outward dislocation, Stimson states that 10 cases have been reported, by Langier and Lacombe, Tutschek, Mignot-Danton,¹⁰ Désprés,¹¹ and Bayer. It is pointed out by Stimson, as also it has been indicated by others, that in virtue of the manner in which the proximal end of the second metatarsal is locked between the first and third cuneiforms, such a lateral dislocation can hardly occur uncomplicated by fracture unless it be preceded by an upward or downward displacement. The proximal end of the second metatarsal has usually been broken, though in two cases it was the third that was fractured. In four of the ten cases quoted by Stimson reduction was effected.

PREPONDERANCE OF OUTWARD DISLOCATIONS.

The reasons for the preponderance of outward dislocation are worthy of brief reference. They are almost solely anatomical. Notice has already been taken of the fact that lateral displacement is possible only as a sequel to upward or downward displacement, or when associated with fracture. It is hardly necessary to point out further that where lateral displacement eventuates upon the vertical dislocation the latter is almost always dorsal in direction. The explanation of the preponderance of the outward displacement is not far to seek when regard is had to the anatomical relations of the cuneiform-metatarsal joint. The notch between the first and third cuneiforms into which the second metatarsal fits is not similar on its two sides. Its internal angle is about 90 degrees, while its outer is almost 120 degrees. This admits all the more readily of primary slipping of the in-fitting metatarsal towards the outer side. Further, the inner wall, formed by the first or internal cuneiform, is fully 1 cm. in height; while the outer wall, formed by the third or external cuneiform, is at most 0.5 cm. in height, so that here again is a condition which predisposes to the outward displacement.

Associated with the dislocation of the metatarsals, as has already been said, there may be fracture of the bases of the second or third metatarsals or of both. Also, there may be any degree of damage to, or comminution of, cuneiforms, and even of the cuboid. The degree of injury to, sprain, or rupture of dorsal and plantar ligaments, also of interosseous ligaments and muscles, will vary in different cases, but must always be considerable.

CAUSES AND METHOD OF PRODUCTION OF THE INJURY.

As regards the *actual forces causing* the metatarsal dislocation, the following facts are ascertainable from the records.

Falls of different kinds have caused the injury. For example, in several recorded cases, the first metatarsal has been dislocated by a fall from a horse, the actual displacement being determined by the pressure of the stirrup. A fall from a height—the patient alighting on the anterior part of the foot—has led to the upward displacement of the whole, or part, of the metatarsus, especially when the metatarsus has been forcibly plantar-flexed. Falls of heavy weights upon the upper part of the instep, driving the tarsus violently downwards, have led to the upward displacement of individual bones or groups of bones of the metatarsus.

Theoretically, any form of violence which is applied in the direction of abduction of the anterior part of the foot, the heel (or tarsus) being fixed, or which tends to produce forcible adduction of the heel (or tarsus) while the anterior part of the foot is fixed, may produce outward dislocation of the metatarsus. Two of the cases which I have to report are examples of the second mentioned type.

Conversely, it may be surmised that, did anatomical conditions permit, or encourage, the inward dislocation, it might be produced by violence tending to produce forcible adduction of the anterior part of the foot, the heel (or tarsus) being fixed; or violent abduction of the heel (or tarsus) the front of the foot being fixed.

Forces of complex character, and applied in irregular fashion, will have to be assumed in many cases where, as for example in my third case, one metatarsal is driven in one direction and the others in the opposite direction. It is possible that the tarsus may have acted as a wedge, driving the metatarsals apart; or some direct violence of a specially localized type, applied immediately to the sole, or to the dorsal surface of the foot, may have caused the bones to be driven in opposite directions. The comparative capacities of particular ligaments for resisting special strain, and the varying force of intrinsic muscles, as applied in special circumstances, may also require to be taken into consideration.

DIAGNOSIS.

The diagnosis of metatarsal dislocations should not be, nor is it, so far as my experience enables me to judge, difficult. More especially is this the case now that we are able to command the aid of the *x* rays to supplement what can be learnt from a careful consideration of the clinical pictures which such injuries present. Certainly, so far as the complete dislocations are concerned, there are no special difficulties.

Henry Morris¹² writing, in 1883, of the dislocations upwards and downwards, says that these "are readily diagnosed by the shortening of the foot, without a corresponding elongation of the heel, and by the projection and depression on the instep." Gould and Warren¹³ corroborate thus: "When backward dislocation occurs, the foot is shortened, and a prominence, with a groove either before or behind it, is seen in the plantar and dorsal displacements. The foot is usually somewhat adducted, and the hollow of the sole is flattened." Nasse and Borchardt¹⁴ speak of the symptoms of dorsal complete dislocation as being very characteristic (Smyly's case of plantar dislocation being the sole authentic recorded example of the other type), thus: "The whole foot seems shortened, being in the position of moderate talipes equinus, while the toes are held in the position of dorsal flexion by the tense extensor tendons. The arch of the dorsum of the foot is exaggerated; right across it there extends a tumour-like prominence, formed by the bases of the metatarsal bones."

Smith¹⁵ called particular attention to the peculiar alteration in the form of the sole of the foot in the upward dislocation. Instead of presenting the usual concavity, this becomes convex, both in its antero-posterior and in its transverse diameters.

As regards the diagnostic features of the lateral dislocations, Morris¹² refers to the projection at the inner or the outer border of the foot; Gould and Warren¹³ point out that there is not necessarily in these any shortening of the foot, but that there is "some adduction or abduction, according as the displacement is inward or outward." Nasse and Borchardt¹⁴ are more explicit. They say: "The lateral dislocations at the Lisfranc joint are easy to recognize. In the outward dislocation the point of the foot is mostly always abducted; on the inner edge the first cuneiform, and on the outer margin the base of the fifth metatarsal project markedly. In the inward dislocation the first metatarsal forms, on the inner margin of the foot, a definite projection; correspondingly, on the outer side there is a depression, behind which the cuboid projects."

These descriptions are quite in keeping with the findings in the cases which have come under my observation. It is, in the normal foot, so easy to make out prominent bony landmarks at the tarsal-metatarsal region, that any considerable departure from the normal regularity ought to be readily appreciated, provided that a careful inspection and manual examination be made; especially if comparison be made with the other (the uninjured) foot, and provided that the examination is not delayed long enough for swelling of the soft parts to so mask contours as to preclude accurate recognition of bony outlines. Parenthetically, it may be said, that in such gross displacements as we have to deal with in cases of complete dislocation the swelling of the soft parts must be very considerable to

completely obliterate and conceal such gross changes of bony relations. Practically, indeed, this is not, in my opinion, a substantial difficulty.

X RAYS IN DIAGNOSIS AND IN TREATMENT.

Further, as has been already said, we have now the all-important x-ray aid to an accurate diagnosis, and this aid ought never to be neglected where there is no insuperable obstacle to its use. Even if the diagnosis seems quite clear enough without its use, it will almost certainly determine a more accurate appreciation of the exact physical condition, and not improbably lay bare some detail which would otherwise be overlooked. It is, besides, a very real, if not absolutely essential, aid to estimation of the completeness or incompleteness of reduction of the displacement once this has been undertaken, and it may be laid down quite definitely that, whatever its value, in an individual case, from the point of view of diagnosis, as a guide or check regarding the efficiency of reduction, it ought never to be neglected. Both at the time of reduction, and at some period subsequent to this (say two or three weeks), the efficiency of the treatment should be confirmed by x-ray plates, taken both in the vertical and in the horizontal planes.

REDUCTION OF DISPLACEMENT.

Reduction of these dislocations should not present any insuperable difficulties, though the records of some of the earlier reported cases would seem to suggest that reduction might prove a formidable, and even an impossible, undertaking. Indeed, there are on record at least two cases in which death occurred as the result, so it would appear, of the attempts at reduction.¹⁶

In most cases, and certainly where any difficulty is encountered, an anaesthetic should be given—ether or chloroform. Under its influence there ought not to be much difficulty in manoeuvring the displaced bones into their proper position. In the case of dislocation of single bones, traction upon the displaced bone, combined with or followed by direct pressure on the projecting or displaced end, should in most instances readily effect reduction, and much the same may be said of dislocations of several bones together. If replacement prove difficult, open operation, under simple aseptic precautions, may quite readily be resorted to, and the bone or bones can be directly levered into position, or any part which cannot be fully reduced may be resected.

As for the complete dislocations, traction on the anterior end of the foot, combined with direct pressure on the dislocated bones, will generally prove effective. For the vertical dislocations this is certainly all that is ever likely to be necessary. In the case of the lateral dislocations the tarsus is first of all fixed, then traction on the anterior part of the foot is applied in such a manner as to exaggerate the pathological position. Simultaneously with this exaggeration of the malposition lateral pressure in the direction of adduction or of abduction will complete the reduction—adduction where the dislocation is outward, abduction in the case of the inward form (which, as already stated, is probably very rare).

In the three cases which I have to report reduction proved comparatively easy. In all three, chloroform anaesthesia was employed, and the actual reduction was accomplished by a combination of traction on the anterior part of the foot, adduction and abduction movements, plantar flexion followed by dorsal flexion, and direct pressure on projecting bones. In all three cases the general correctness of the replacement was at once verified by fluorescent screen and subsequently by x-ray plates. As stated earlier, it is probably wise to follow this course of immediate and later x-ray confirmation wherever possible.

TREATMENT AFTER REDUCTION.

As regards further treatment, it may be said that, once correctly reduced, these dislocations do not tend readily to redisplacement, but it is well to provide artificial fixation, by means of a plaster-of-Paris case, for several weeks. In the case of the upward displacements a period of two or three weeks ought to be ample, but where any strain is likely to be put on the foot this period may be prolonged somewhat with advantage.

In the case of lateral dislocations fixation should be maintained for a longer period, and this for two reasons: (1) on account of the greater damage to ligaments and soft tissues; (2) on account of the frequent coexistent fracture of metatarsals or cuneiforms. A period of six to eight weeks in plaster is regarded by most writers as essential, and it is generally advised that thereafter, and for some considerable time, an artificial support should be employed for the arch of the foot—such, for example, as a Whitman's brace.

In the less frequent cases, where time, care, and perhaps expense are of less importance, all forms of fixation may perhaps be dispensed with, and provided reduction has been effectively accomplished, reliance may be placed entirely on rest for a prolonged period, combined with gentle and systematic massage.

In two of the cases which follow, fixation by plaster was employed for three weeks, and no subsequent support was found necessary or advisable. In the third, the plaster case was kept on for four weeks, and was not thereafter reapplied.

It is obvious that early massage, so long as it is gently carried out, as also early active movements of the foot so as to prevent the stiffening and immobilization of the intertarsal and tarsal-metatarsal joints—so long as such movements stop short of strain on the injury-weakened ligaments—will be the most valuable and important factors in determining ultimate complete restoration of function.

In cases which have been recorded, quite tolerable functional capacity has in time been developed, even where replacement could not be at all, or at any rate completely, effected. Thus, in one case, recorded by Cock,¹⁷ of complete upward dislocation, combined with fracture of the first cuneiform and scaphoid, reduction was not accomplished, and yet the patient was back at his work—as a railway porter—within eight months of receiving his injury.

CASE I.

R. J., aged 30, male, admitted to the Broadstone Jubilee Hospital, Port Glasgow, on November 2nd, 1909, suffering from an injury just received. A heavy bale of yarn fell upon him from some height, causing, in addition to sundry abrasions and some injury to the chest, a somewhat severe injury to his left foot. The mechanism of production of the injury to the foot seems to have been as follows: The man got just sufficient warning of the falling weight to permit him to brace himself for the shock, but insufficient to allow of his evading it. When struck by the bale he had his feet widely separated, his back and shoulders somewhat bent, and his muscles tensely strung to resist the impact. He was struck in the upper dorsal region, and it may be supposed that the impact of the force was transmitted down the trunk and both lower limbs in such a way as to exert itself in maximum intensity at the tarsal-metatarsal regions of both feet. It may also be supposed that one foot got a larger share of the violence than the other. In any case, only one foot suffered to any extent. By reason of the wide separation of the feet, and the tense state of the limb muscles, the anterior parts of the feet, and especially of the foot which suffered, may be supposed to have been fixed—jammed firmly against the ground. The downward acting force would tend, therefore, to carry the whole tarsus and heel portion of the foot inwards—as actually happened. That this had indeed happened was at once obvious on examination—even on a quite superficial examination—of the foot. The internal cuneiform could be seen and felt bulging prominently on the inner side of the foot; the fifth metatarsal could be recognized projecting abnormally on the outer side. There was considerable swelling of the soft tissues and much pain. Crepitus was felt, suggesting some comminution of bone either at proximal ends of second and third metatarsals, or of cuneiforms, or of both. Examination by fluorescent screen, and subsequently by x-ray plate, confirmed the diagnosis of dislocation and determined the extent of displacement as well as of the accompanying fracture.

Considering the violence necessary to produce the dislocation, the amount of actual damage to bone seems small. In the accompanying reproduction (Fig. 1) of the radiographic picture the displacement is well shown, but the exact damage to bone is not quite definite. That there is some bone fracture is obvious, but whether this has been of the proximal end of the second metatarsal or of the projecting extremity of the third cuneiform is not clear.

As the man suffered great pain the dislocation was reduced without delay under chloroform anaesthesia. The reduction was easily accomplished by traction on the anterior part of the foot, followed by plantar flexion and forcible adduction. The correctness of the reduction was at once verified by fluorescent screen and confirmed by an actual plate exposure. The corrected foot was at once fixed in a plaster-of-Paris case, with the ankle at right angles, and the foot in a position of moderate inversion.

The plaster support was retained for three weeks. The patient had no pain to speak of, once the reduction had been

accomplished, and swelling quickly subsided; so much so that the plaster case was probably not of very great value after the first few days. At the end of three weeks it was removed and massage was instituted. After a week of this the patient was allowed to get about, but was advised not to put his full weight on the foot for a week or two more. The arch of the foot was well restored, and there was not then—nor did there appear later—any tendency to recurrence of displacement. Completely satisfactory function was restored in about ten weeks. No artificial support was employed subsequent to the removal of the plaster.

CASE II.

W. P., aged 46. Admitted to the Broadstone Hospital, Port Glasgow, on February 28th, 1911. The patient, a man, was injured in a shipyard by a falling plate, which slid down the front of the right leg, and pinned the anterior part of the booted foot, so that it had to be extricated subsequently by the man's comrades. Immediately the foot was thus caught by the fallen plate as in a vice the man fell over to the right, so that the heel portion of the foot was violently adducted, the anterior part of the foot being meantime firmly fixed. Probably there was also at the same time some degree of torsion violence, applied at the tarsal-metatarsal articulation. The mechanism of production of the injury was, thus, not very dissimilar from that in Case I, though in the latter the fixation of the anterior part of the foot was not produced in the same direct way.

The resulting lesion was quite obvious on admission and was recognized as a dislocation of the metatarsus by Dr. Alex. Butler, one of the surgeons to the hospital. The anterior part of the foot was markedly adducted and somewhat everted. The internal cuneiform was prominent at the inner side, the fifth metatarsal at the outer side. The base of the first metatarsal bulged slightly on the dorsum of the foot. There was considerable tension of the skin and soft tissues. No crepitus was felt. Pain was considerable, and to relieve this temporarily the foot was fomented till the following day, when, in ordinary course, I visited the hospital. I had no hesitation in coinciding with the diagnosis made by Dr. Butler. Skiagrams were obtained, which confirmed our joint diagnosis (see Figs. 2 and 3). One was taken from above downwards, the other with the foot in profile. The former, though quite a good picture, cannot be said to give much support to the idea of there being any accompanying fracture, either of metatarsal base or of cuneiform. The latter shows the slight dorsal displacement of the proximal end of the inner metatarsals.

Under chloroform anaesthesia the dislocation was reduced. What seemed to be quite definite crepitus was felt during reduction, but the exact site of the crepitus was not clearly established, though it was thought to be at the site of the in-setting proximal end of the second metatarsal. Once the reduction was complete crepitus could not again be elicited, and, in view of the fact that no crepitus was felt before reduction, and that the skiagram gives no definite indication of fracture, the suggestion may with some justification be made that in this case the added torsion element mentioned in describing the mechanism of production of the dislocation, may have had the effect of producing first of all a dorsal displacement such as would permit of the second metatarsal slipping over or past the third cuneiform without fracture. Fig. 3 in any case makes it clear that some dorsal dislocation had occurred. In this case there was rather more difficulty encountered in accomplishing reduction than there was in Case I.

After reduction the foot was encased in plaster in slightly over-corrected and somewhat inverted position. Ten days later the patient was dismissed with his foot still in plaster.

Three weeks after the date of reduction the plaster was removed. There was slight thickening about the inner side of the tarsal-metatarsal articulation, but the position of parts was all that could be desired. The plaster was not reapplied, but firm bandaging with flannel was continued for some time and massage was instituted. The patient was encouraged also to begin carefully to use the foot actively. Function was well restored after three months.

CASE III.

T. H., aged 24, male, admitted to the Broadstone Hospital, Port Glasgow, on August 28th, 1911. Injured by falling a considerable distance in a shipyard. His right foot was seriously damaged by the fall, but this was the only injury of consequence he received. Exactly how the foot was injured neither he nor his mates could definitely say, and so the actual mechanism of production of the lesion sustained could not be established. The only statement made—for what is worth—was to the effect that the man did not alight on the injured foot. In view of the nature of the damage produced this may perhaps be open to some doubt, for, as I have already suggested in the general statement which has gone before, the resultant displacement of the metatarsals certainly somewhat suggests that the cuneiforms were impacted violently against the metatarsus, driving the first metatarsal inwards and the second, third, and fourth outwards. In any case, on admission to hospital, the nature of the injury was quite evident. I saw, and dealt with, the case on the day the injury was received. The complete inward dislocation of the proximal end of the first metatarsal was quite obvious; it could be felt, and indeed seen, projecting on the inner side of the foot. The next three metatarsals could be seen and felt projecting outwards and on the dorsal aspect of the outermost cuneiform and cuboid, the skin being stretched tightly over them. It was easily determined, on examination, that the second and third metatarsals were fractured close to

their proximal ends, and part at least of the dorsal prominence was caused by the special projection of the sharp proximal ends of the distal fragments of these bones. The small proximal fragments were, however, also dislocated upwards. There was much bruising of the tissues of the dorsum of the foot, much effusion of blood into the tissues, and at one place the skin was almost torn through. Had reduction not been at once effected it is almost certain that the skin would have given way and sloughed.

The diagnosis was confirmed by means of the x rays. Fig. 4 shows quite clearly that the facts made out by ordinary methods of physical examination conformed with actuality.

Reduction was carried out without much difficulty on the same day, chloroform anaesthesia being employed, Dr. Butler anaesthetizing, and Dr. James Walker assisting me. The displacement of the first metatarsal was first corrected, and thereafter, by a reversal of the lateral pressure and adduction, the outer bones were readily replaced.

Plaster-of-Paris support was employed for four weeks; thereafter bathing, rubbing, and regulated passive and active movements being instituted. The result was in every way satisfactory, though stiffness persisted longer than in the other two cases.

Fig. 5 shows the complete correction of the displacement; it is a reproduction of a skiagram obtained about five weeks after the injury.

The publication of this paper has been delayed with a view to the searching of the records of the large Glasgow hospitals for cases of such injury. Only one case has been found in the last ten years' records of the Western Infirmary, which was at least labelled "Dislocation of metatarsus." The index title was all that one could obtain; access could not be had to any details, so that even this one case can hardly be counted. In neither of the other two hospitals could information be got from those likely to be aware of the occurrence of such an injury; in both, the radiographic experts could call to mind no example of such a condition as I have drawn attention to in this report.

This seems to make all the more remarkable the fact that, in a comparatively small hospital such as that at Port Glasgow, we should have had in so short a time, comparatively, three examples of a relatively infrequent lesion.

REFERENCES.

- 1 Stimson: *A Treatise on Fractures and Dislocations*, 1908, p. 839.
- 2 Nasse and Borchardt: *Handbuch der prakt. Chirurgie*, vol. iv, Part 2, p. 514. Pouse: *Laterale Luxation sui Lisfranc*, *Muench. med. Woch.*, 1897, Nos. 21-22.
- 3 Griffith: *A Handbook of Surgery*, 1904, p. 251.
- 4 Wharton: *Minor and Operative Surgery, including Bandaging*, 1905, p. 446.
- 5 Erichsen: *The Science and Art of Surgery*, tenth edition, vol. i, p. 714.
- 6 Fischer: *Deutsche Zeitschrift für Chirurgie*, vol. lxxiv, p. 55.
- 7 Ely: *Annals of Surgery*, August, 1906.
- 8 Smyly: *Dublin Quarterly Journal of the Medical Sciences*, 1854, vol. xvii, p. 317.
- 9 Stimson: *A Treatise on Fractures and Dislocations*, 1908, p. 841.
- 10 Mignot Danton: *Arch. gén. de méd.*, 1866, vol. ii, p. 405.
- 11 Després: *Bull. de la Soc. Anatomique*, 1878.
- 12 Morris: *A System of Surgery* (Holmes and Hulke), 1833, vol. i, p. 1108.
- 13 Gould and Warren: *Internat. Textbook of Surgery*, vol. i, p. 649.
- 14 Nasse and Borchardt: *Handbuch der prakt. Chirurgie*, vol. iv, Part 2, p. 515.
- 15 Smith: *Proceedings of the Pathological Society of Dublin*, vol. ii, N.S., p. 226.
- 16 Hitzig: *Berl. klin. Woch.*, vol. ii, pp. 39-42, 1865.
- 17 Guy's Hospital Reports, 1855, p. 280.

CAPTAIN W. S. PATTON, I.M.S., in No. 50 of the New Series of *Scientific Memoirs by Officers of the Medical and Sanitary Department of the Government of India*, gives a description of the work done by himself in the etiology of Oriental sore in Cambay during June, 1910. His researches were chiefly concerned with the carrier of the parasite, the *Leishmania tropica*, from person to person, and he believes that he has made out a case against the bug. *A Stegomyia* of unknown species and a *Phlebotomus*, probably *Cabu* (Annandale), as well as ordinary house flies and lice, were all studied, but with negative results. In bugs, however (*Cimex rotundatus*), the parasites were found to flagellate in the stomach if the temperature was below 25° C., and this happened also in the nymphs. Complete development apparently did not take place in the adult bug, as flagellates were never seen in them beyond the tenth day, but in the nymphs Patton believes this does take place. In addition to these facts he brings forward epidemiological observations and clinical cases of people who were bitten by bugs and afterwards developed sores. Though all this is very interesting and suggestive, it does not absolutely prove that the bug is the regular intermediate host. It may be so in Cambay, but it seems just as clear that in other parts of the world—Bagdad, for example—it is not. It is evident that more work on the subject is required.

MR. ARCHIBALD YOUNG: DISLOCATION OF THE METATARSUS.

Fig. 1.—Case I: Showing complete outward dislocation of metatarsus. Note prominence on inner side of foot caused by internal cuneiform.

Fig. 2.—Case II: Showing outward dislocation of metatarsus.

Fig. 3.—Case II: Showing slight dorsal displacement of proximal end of metatarsus.

Fig. 4.—Case III: Before reduction of dislocation. Note prominence on inner side of foot caused by proximal end of first metatarsal.

Fig. 5.—Case III After reduction of dislocation.