are claimed to be good, although it must be remembered that the percentage mortality from tetanus in Egypt is not so high as in the British Isles. The percentage mortality is less the longer the period of incubation, and where this is between three and four weeks the mortality in the British Isles is about 50 per cent. For an adult a dose of 6 c.cm. of a 25 per cent. solution in distilled water may be given. This contains $1\frac{1}{2}$ grams (about 22 grains) of magnesium sulphate.

As the injection of antitetanic serum has proved to be of very little curative value in tetanus, the injection of magnesium sulphate in this manner is certainly worth a good trial. The withdrawal of cerebro-spinal fluid is doubtless an additional advantage.

I have to thank Dr. Alexander Morison, into whose ward the patient was admitted, for his kind permission to publish this case.

CAESAREAN SECTION, WITH ESPECIAL REFERENCE TO THE RARER INDICATIONS FOR THE OPERATION.*

By W. H. MAXWELL, M.A., M.B.CAMB., F.R.C.S.ENG.

THE title of this paper was suggested to me by the circumstance that, while it has fallen to my lot to perform this operation four times during the last two years, in none of these four cases, curiously enough, has the indication for it been that for which I suppose something like 90 per cent. of Caesarean sections are performed—namely, pelvic contraction. The three cases upon which my remarks are based form an interesting series, in that each illustrates certain points of importance to the student of what one may perhaps be allowed to call the “higher obstetrics.” My personal experience of Caesarean section is relatively small; I have done the operation myself five times, and have had the opportunity of assisting at it in seven other cases. My own five cases have been unattended by a fatality; of the other seven, two died, the cause of the fatal issue being, as I shall point out later, apparently identical in both. In the first case that came under my observation the operation was done by my old friend the late Mr. E. H. Douty, of King's College, Cambridge, in Addenbrooke's Hospital in 1896.

For some reason or other, in South Africa we meet with few examples of the higher degrees of pelvic contraction; hence Caesarean section and kindred operations are called for somewhat rarely. I must make it clear that in speaking of Caesarean section I refer entirely to the classical operation of Säger and Porro—that is to say, what should be designated “anterior transperitoneal hysterotomy”—my experience of the various modifications suggested, of which “vaginal” Caesarean section seems the most extolled, being *nil*. With this premise I will proceed to a brief description of the three cases referred to.

CASE I.

Mrs. M., aged 40, married twelve years, had had four children, of whom three were living still, and two miscarriages. She was admitted into the Queen Victoria Hospital for Women early one morning in labour, and her arrival intimated to me by a telephone message from the matron to the effect that she had examined and “could not find the os.” Not, perhaps, without an inward smile, I obeyed the summons, and on making a vaginal examination likewise was unable to find the os. Finally, on introducing my arm, under light anaesthesia, nearly to the elbow, the partially dilated os could be made out lying right up in the lumbar region, the impression being that a very large rounded tumour was obstructing the pelvis and pushing the cervix upwards and backwards. The patient said (in explanation of a median abdominal cicatrix) that an operation had been done a year before in the General Hospital, but she was ignorant of its nature. Telephone inquiries elicited the fact that the operation had consisted in a direct “ventral fixation,” and the nature of the case was now evident, the “tumour” being merely the body of an extremely anteverted uterus, which, in fact, was almost in an inverted position. It was fairly obvious that extraction of the child *per vias naturales*, even dead, was wellnigh, if not altogether, impossible, and my colleague, Dr. Gordon Grant, who saw the case, confirmed this opinion. I therefore decided on Caesarean section. The patient was prepared as quickly as possible, the child, a

healthy full-term male, extracted without any difficulty, and the placenta and membranes removed; the womb was then drawn out through the abdominal incision and carefully sutured. Haemorrhage was negligible in quantity throughout and subsequent contraction was excellent. The mother made an uneventful recovery and was able to nurse the child as usual.

It was noteworthy in this case that, in spite of the almost incredibly anteverted position of the uterus, one was not troubled during the operation by any obvious adhesions or other sources of difficulty. Hitherto in performing the—often very useful—operation of ventral fixation or hysteropexy (in contradistinction to the “indirect” operation of Gilliam and its countless modifications) I have always had in my mind the fear lest it might—as it certainly does—interfere with subsequent pregnancy, unless great care be taken to pass all sutures well down on the anterior uterine wall; and not a few cases have been published in which recurring abortions seem to have followed the operation; but I had never realized before that such an extreme malposition of the uterus could result as to demand Caesarean section. About two years ago, however, there was published in the BRITISH MEDICAL JOURNAL¹ by Dr. Amand Routh a carefully collected series of eight cases similar to the one I have just described. I can only say that had I not seen the condition before delivery in this case I should have felt somewhat sceptical as to the validity of the indications for the operation. Possibly a similar suspicion might arise in the minds of many as regards the next case to which I have to refer.

CASE II.

Mrs. T., aged 34 years; married five years; one child living aged 4 years. Had had a “bad miscarriage” eight months previously. This patient was sent into the Queen Victoria Hospital one afternoon with a diagnosis of placenta praevia, by her medical attendant, Dr. Brebner. When I saw her soon after admission she appeared pale, but not markedly exsanguine; there were no pains and not the slightest sign of dilatation of the cervix; the general indications were those of about six and a half months pregnancy, the fetal heart being audible. She had bled a good deal, and I decided that something ought to be done on the following day.

During the night she had steady and almost continuous bleeding, and when I saw her at 6 a.m. she was in an extremely grave condition, partly, it appeared, owing to cardiac failure supervening on pretty severe haemorrhage. There was still not sufficient dilatation of the cervical canal to admit even the tip of a finger, and her condition was so bad that it seemed doubtful whether we could get her alive on to the operating table. Saline infusion (subcutaneous and rectal) improved the pulse a little, but it was tolerably obvious that any attempt at an *accouchement forcé* must result in death from the further haemorrhage inevitable during extraction of the uterine contents; I therefore decided upon Caesarean section as giving her a better chance. This was performed in the usual manner, saline infusion being kept up during the operation. The uterus was opened with the greatest possible rapidity after being drawn right out through the parietal incision; the child, followed by the placenta and membranes, was removed, with an astonishing absence of haemorrhage, in about three minutes from the start. The pulse practically disappeared, but was restored by filling the peritoneal cavity with warm normal saline solution while the stitching up of the uterine wound proceeded, another jugful being introduced just before closing the abdominal incision. The only loss of blood worth mentioning came from the placental site after removal of the uterine contents, but good contraction prevented any great loss.

For the first twenty-four hours it seemed very doubtful whether the patient would recover, as the cardiac condition gave rise to grave anxiety; but after this she improved so rapidly that she was able to sit up by the seventh day, and her recovery was not retarded by any complication. The child, of course, though delivered alive, was not viable. I should mention that the placenta, as far as a hurried examination allowed one to determine, was as nearly “central” in position as it could possibly be.

The great point of interest in this case lies in the consideration of *placenta praevia* as an indication for Caesarean section. The question has, of course, been raised often enough; and while it is usually admitted that on purely theoretical grounds Caesarean section should be the ideal treatment in certain cases, yet the general consensus of opinion appears to condemn this operation as a practicable line of treatment owing to the absence of any reliable method of gauging, in any given case of placenta praevia, how much or how little difficulty and danger the condition is going to cause. Any one who has had an even moderately extensive experience of this dangerous complication must have observed that many cases do extremely

* Read before the Witwatersrand Division of the Transvaal Branch of the British Medical Association.

well in spite of pretty severe bleeding, and that frequently the expected degree of trouble fails to materialize. The mortality *qua* the child is very high, no doubt because the very condition that plays so important a part in conserving the waste of maternal blood—namely, obliteration by thrombosis of the torn sinuses of the placenta *pari passu* with its detachment during delivery—is actually inimical to the child's chance of surviving; given, that is to say, a certain degree of placenta detachment the child, in spite of its being extracted with the utmost rapidity, is of necessity delivered in a too exsanguine condition to be capable of surviving; the drainage of blood from the placental sinuses during delivery means also, it must be remembered, drainage from the child's vascular system. It would appear from such considerations that those who have been foremost in urging the propriety of hysterotomy in cases of placenta praevia have for the most part had in mind rather the child's welfare, and if this be so the case I have recorded here is, I think, an interesting one in view of the fact that it illustrates the possibility of Caesarean section being desirable in the interest of the mother, though perhaps it must not be held to establish a precedent.

It is noteworthy that the late Mr. Lawson Tait strongly urged the adoption of Porro's operation in cases of placenta praevia, and recently I came across a remark of Professor Japp Sinclair's in 1901 to the effect that sooner or later placenta praevia would have to be considered as a possible indication for Caesarean section. Yet the statement was made only the other day, without qualification, "Placenta praevia is *not* an indication for Caesarean section"! Without wishing to emphasize unduly the value of my own judgement, I can affirm without any hesitation that in this particular case and at the particular stage at which the operation was performed no other kind of treatment would have availed to save this patient's life, though I am willing enough by all means to class such a case among those proverbial exceptions which are said to "prove the rule."

CASE III.

A woman, aged 42, was sent to see me when just five months pregnant for the first time, having been married two years. She complained chiefly of a good deal of pain about the rectum (though not markedly constipated) and some increased frequency of micturition. The first thing that struck me was that, while the menstrual history suggested five months' pregnancy, the uterus reached well above the umbilicus. Examination revealed a fibroid (evidently), about the size of a large orange, lying in the pelvis and pressing on the rectum, the cervix being pushed forward, considerably flattened, in front of the tumour against the pubes. It was quite impossible to push the tumour up, and at first, on account of the pain and pressure symptoms, I nearly decided on immediate operation, but was deterred from this course by the wellnigh certain prospect of having to do a hysterectomy; the position of the tumour gave very little hope of the possibility of a myomectomy with conservation of the uterine contents.

I put it, therefore, to the patient, who was extremely anxious to have a child, that it was worth while for her to try and hold on until full term, as she would then have a good prospect of a living child, and it was not likely that things would get very markedly worse during the next four months.

She followed this advice, and although at intervals of every few weeks she had rather severe attacks of pain and occasional slight bleedings, she managed to get along fairly comfortably up to within a week of the time reckoned upon for natural delivery—had that been possible—and two days before the time at which I had advised her to come into hospital I received a telephone message at 7 a.m. saying that she had arrived and was in labour.

Realizing the danger of delay, I operated at 9.30 a.m., the skin being prepared by the iodine method; no vaginal examination had been made for more than six weeks previously. The abdominal incision reached to about 2 in. above the navel, and swabs were then packed in between the abdominal wall and the uterus, the latter, which was studded with fibroids, being incised without eventration. The organ was so much distorted

by these numerous tumours, more particularly by one rather large intramural fibroid in front, that it was unusually difficult to hit off the uterine cavity, especially as it was now obvious that the membranes had already ruptured, evidently before admission, labour being, in fact, well on in the first stage. I feared at first that the child was dead; but it speedily revived and did well (male, weighing 5 lb.). I removed placenta and membranes, all of which were somewhat adherent, and then drew out the uterus. It was obviously futile, in view of the large number of fibroids of all sizes studding the uterus, to attempt multiple myomectomies, and the tumour in particular which had obstructed the pelvis, as shown in the diagram,* was situated so low down posteriorly that even after emptying the womb it would have been practically impossible to enucleate it. I proceeded, therefore, to remove the uterus by supravaginal amputation; both ovaries looked healthy and were not interfered with. The cervix had undergone such a degree of dilatation that it formed quite a broad stump, which, after approximating its cut surfaces with "mattress" sutures, was covered over in the usual manner with peritoneum. Haemorrhage gave no trouble—a somewhat unusual feature, to judge by my own experience, in amputation of the pregnant uterus, the cervical region often being the seat of a good deal of troublesome oozing.

The operation took over an hour and a half, but the patient's condition gave rise to no anxiety then or later, and she made an excellent recovery; plenty of milk appeared by the fifth day, and the child likewise did well. The obstructing myoma was subserous and almost pedunculated, but could not have been pushed up, being apparently incarcerated below the sacral promontory, on the one hand pressing upon the rectum and on the other pushing the cervix and bladder forwards against the pubes.

Diagrammatic representation of the condition of the uterus in Case III. A, Small fibroid, palpable before operation. B, Uterine cavity. C, Intramural fibroid. D, Line of section of cervix. E, Subserous fibroid obstructing pelvis.

In seeking to emphasize the points of chief importance which the cases I have referred to seem to illustrate, I venture to crave the indulgence of some of my colleagues whose experience of Caesarean section is possibly more extensive than my own, and I think sufficient justification for so doing can be found in the extreme importance to the obstetrician and his patients of a proper grasp of both the possibilities and the limitations of this time-honoured method of delivery; one would be tolerably correct in asserting that in no other department of surgery have such triumphant results attended the progress of aseptic technique. But it is of prime importance to understand that the modern diminution of the mortality of the operation is due, next to the development of a proper application of the principles of asepsis, to

the fact that the operation is now performed, not as a last resort—a desperate measure to save the mother's life—but either before the onset of labour or, at all events, before labour has proceeded very far. Even now, it seems to me, the cardinal point to remember is to operate early; Caesarean section done before rupture of the membranes (or, at any rate, before the end of the first stage) is one thing, Caesarean section done after the end of the first stage is quite another. Probably the mortality proper to the former operation is between 2 and 5 per cent.; that of the latter even now must be something like 15 to 20 per cent. And I must point out here that, especially if the rule be adhered to that vaginal examination just before the operation should be avoided whenever possible, one is very apt to discover that labour has progressed much further than one had suspected. In the only two fatal cases that have come under my own observation labour was found to be well on in the second stage; that is to say, not only were the membranes ruptured, but the child's head was in the vagina, and so, in order to release the head, the operator had to prolong his uterine incision down through the cervix—a proceeding which not only involves an otherwise unnecessarily long incision, but is almost certain to result in trouble, owing to haemorrhage from the numerous veins surrounding the upper part of the vagina. Proper apposi-

* The specimen was shown at the October meeting of the Branch.

tion of the edges of the uterine wound is much more difficult in this region, owing to the thinning of the uterine walls produced by the phenomenon known as "retraction." The whole operation is easier if the uterine incision can be kept high up, where the uterine muscle is thicker, though I can see no definite advantage in the transverse incision known as the *Fundusschnitt nach Fritsch*. Any one, too, who has had some experience of these cases will have noticed that the opening of the uterus is an easier proceeding when the membranes are intact.

Providing that one be lucky in avoiding the placenta, the uterine incision may be made slowly and deliberately with an extremely small loss of blood until the membranes bulge into the wound, when the amniotic cavity can be opened at leisure. The only disadvantage in cutting through the placenta is that one is obliged, in order to arrest the haemorrhage as soon as possible, to hasten the process of extraction of child and secundines, but it is scarcely worth while, as some have suggested, to try and work round the edge of the placenta instead of cutting through it. I can see little, if any, advantage in "eventration" before delivery of the child; a much longer abdominal incision is required, whereas, if the womb be left *in situ*, it is easy enough, by arranging a couple of large gauze swabs on either side between the uterus and the abdominal wall, to prevent the escaping liquor amnii and blood from entering the peritoneal cavity.

As soon as the placenta and membranes have been removed the uterus should be drawn right out and suturing begun. The manner of closing the uterine incision is of considerable importance. I personally prefer a not very coarse "chromicized" (Red Cross No. 1) catgut, and for the first row I use "mattress" sutures, as they are less liable to cut out than single interrupted stitches, each stitch taking a wide grasp of muscle and just missing the uterine mucosa. The second layer is a continuous suture ensuring proper approximation of the serous coat. There should not be more than a quarter—or at the outside three-eighths—of an inch between any two sutures of the first row.

The history of the development of the modern conservative operation is of some interest, because for a very long time it was the universal custom not to stitch up the uterus at all, and the mortality arose principally from infection of the peritoneal cavity through the uterine wound, and without any understanding of asepsis it will easily be believed that any attempt to close the uterine wound probably made matters worse. Hence it is understandable what a different complexion was put upon the operation by Porro's suggestion (first carried out in 1876) of removing the uterus altogether. Now, it must be remembered that the modern operation of supravaginal amputation is *not* the original method of Porro. His operation consisted in drawing out the uterus, transfixing with the "pin," and applying the Koeberlé *serre-noeud*, after which the body of the uterus was cut off; the whole operation could, of course, be done very rapidly, and the mortality was enormously reduced by this method. I believe that even now there are not a few cases, especially among those operated upon late in labour, where the mother's life could be saved by Porro's method.

It is no doubt hard for us to realize what Porro did for Caesarean section. His operation at once precluded the possibility of subsequent peritoneal infection and of haemorrhage, besides possessing the advantage already mentioned of extreme rapidity. The objection has been raised that the operation is "unsurgical." This is nonsense; the most "surgical" procedure is that which, under any given circumstances, gives the best chance of saving the patient's life, and, as regards safety, Porro's original method is not easily beaten. At the time when, in 1777, young Sigault of Paris performed the first successful symphysiotomy, public opinion as regards Caesarean section was at so low an ebb that the operation almost fell into disuse, but it was Baudelocque chiefly who demonstrated the fallacy of supposing symphysiotomy to be so very much the safer operation. So things continued without much progress until relatively late on in the nineteenth century, when Porro (of Milan) introduced his operation; immediately prior to this the mortality was well over 50 per cent. It is difficult to get satisfactory statistics, but it is probable

that Porro's method further reduced the mortality by about another 20 per cent. What we may call the final stage was reached when Säger of Leipzig not very long afterwards proposed the adoption of proper suturing of the uterine wound, and Säger's *conservative Kaiserschnitt* remains, with various modifications and improvements, the operation of to-day. With the advent of Säger's operation conservatism justifiably reasserted its claims, and at present the universal practice is to trust to careful suture and leave the uterus intact; the question as to whether sterilization should be effected must of course be decided by a consideration of the merits of the individual case.

In only one of my own cases did union by primary intention fail, and this occurred in a patient from whom I removed an acutely inflamed appendix early in labour (primipara), Caesarean section being done simultaneously because it appeared the best course in the circumstances. The appendix infected the uterine wound and a large abscess had to be opened and drained; the patient ultimately made a good recovery though a slow one.

Since the above paper was written it has, by a curious coincidence, fallen to my lot to operate on a second case very similar to one of those already mentioned; it seems worth while, therefore, to place on record a brief note of the case.

CASE IV.

A woman, married twelve years. Had had eight children, the last two and a half years ago. Prior to her admission to hospital she had been in labour for twenty-four hours; the membranes had ruptured early and repeated vaginal examinations had been made by an unqualified midwife and subsequently by two medical men. On admission the head of the child could be made out lying high above the pelvic brim, the pelvic cavity being nearly filled by a large firm rounded mass, which the examining finger could not pass (a fortunate circumstance, in all probability, for the patient and myself); the pains were fairly strong.

It was obvious that in this case Caesarean section was the cause *ex necessitate*; the usual preparations were therefore made, and the operation proceeded with. The only difficulty in the first stage of the operation arose from the scantiness of liquor amnii, but the child was extracted in a fairly healthy condition, only to succumb ten days later to the effects of a severe burn of its back produced by contact with a damp hot-water bottle. Haemorrhage was easily controlled after eventration, and it was then seen that the fibroid was a solitary subserous one, growing from a point about level with the os internum outwards and backwards under the right broad ligament; the shelling-out process occupied some considerable time, but it was possible to leave a good cervical stump with abundant peritoneal covering, the operation being completed in the usual way. The right ureter had a somewhat extraordinary course, and the right uterine vessels were difficult to secure.

The patient was eventually discharged practically well, but had to remain in hospital for nearly two months on account partly of a troublesome phlebitis arising in old varices of both legs and partly of grave failure of compensation in an old rheumatic heart with rather well-marked mitral disease, the existence of which was discovered subsequently to the operation. The wound healed *per primam*, and gave no trouble. The uterus with the tumour weighed, after removal, about 9 lb.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, 1911, i, p. 189.

THE eleventh international conference on tuberculosis will be held this year at Berlin in October (22nd to 25th). All communications relative to the congress should be addressed to the General Secretariat, Berlinerstrasse, 137, Charlottenburg.

A PAPER by Surgeon Grow, of the United States Navy, published recently in *Ophthalmology*, deals with a subject which has already received the close attention of the British Admiralty—the vision necessary for the men who train and point large naval guns. The United States naval authorities insist that gun-layers shall possess a vision of 6/4.5 in the sighting eye, and 6/6 in the other. In other words, a gun-sighter must have normal vision, for 6/6 is subnormal, although conventionally accepted as normal. The author concludes that all candidates for the original rating of gun pointer or trainer should have a minimum vision of 6/4.5 in the sighting eye, and of 6/6 in the other eye. Hypopia of over 3.00 dioptries should be a cause for rejection. The medical officers should carefully re-examine the eyes of these men once a year, and if any deterioration be present the man should be examined by an ophthalmic surgeon. A slight reduction may be accepted in the case of successful gun-layers who have served a considerable time. In all cases the navy "unlearnable vision test cards" should be used.