seem to me too obvious for discussion, and the same may be said in regard to the conditions of muscular hypertrophy or pseudo-hypertrophy met with occasionally in various cases of primary muscular dystrophy, as well as in the typical pseudo-hypertrophic group. Other diseases, such as the various forms of chronic oedema and elephantiasis in the lower extremities connected with lymphatic obstruction, are still less likely to be confused. Just at the commencement, when the atrophy of subcutaneous fat is still almost limited to the face, the condition might possibly resemble that seen in cases of "bilateral atrophy of the face"; indeed, it is just conceivable that certain cases of bilateral atrophy of the face may have been mild examples of lipodystrophia progressiva, modified by the male sex of the patients, males not being so inclined as females to special deposit of subcutaneous fat in the lower extremities. At the Clinical Society of London, in 1905,⁷ H. Batty Shaw showed a boy, aged 10 years, who had commenced to show bilateral wasting of the subcutaneous tissues of the face when he was 2½ years old. The boy was brought to the hospital because the mother feared he might have tuberculosis. A. F. Hertz and W. Johnson,⁸ at the Clinical Section of the Royal Society of Medicine, in January, 1913, showed a young man, aged 26 years, whose face had become progressively thinner during the last two years, so that his friends thought he must be consumptive. There was no weakness of the facial muscles, and otherwise he was well developed and strong. I instance these two cases of facial wasting in males to illustrate the kind of cases of "bilateral atrophy of the subcutaneous tissue of the face" to which I wish to draw attention in connexion with the diagnosis and pathological classification of "lipodystrophia progressiva."

I have given no photograph or drawing of the condition of lipodystrophia progressiva, but no such illustration is needed. For a typical case one need only imagine a grotesque figure the lower part of which seems to be modelled after an extraordinarily florid Venus of the ultra-Rubens style, whilst the face and upper part of the trunk might call up a picture of one of the forbidding Moerae, or their dreaded sisters, the Erinyes, or might bring to mind the popular idea of one of the witches in *Macbeth*.

REFERENCES.

- ¹ R. Cassirer, *Die vasomotorisch-trophischen Neurosen*, second edition, Berlin, 1912, p. 675. ² H. Campbell, *Transactions of the Clinical Society of London*, 1907, vol. xi, p. 272. ³ H. Campbell, *Proceedings of the Royal Society of Medicine, Neurological Section*, 1913, vol. vi, p. 71. ⁴ E. Hollander, *Munch. med. Woch.*, 1910, vol. lvii, p. 1794. ⁵ Pic and Gardère, *Lyon Médical*, 1909, vol. cxlii, p. 61. ⁶ *Nouvelle Iconographie de la Salpêtrière*, Paris, 1912, vol. xxv, p. 473 (with plate). ⁷ H. Batty Shaw, *Transactions of the Clinical Society of London*, 1905, vol. xxxviii, p. 222. ⁸ Hertz and Johnson, *Proceedings of the Royal Society of Medicine, Clinical Section*, 1913, vol. vi, p. 92.

THE estimated population of Jamaica in 1912 was 845,767, an increase of 6,520. The marriage-rate was 4.2 per 1,000 persons living, the birth-rate was 39.0 per 1,000, and the death-rate 22.1; the infantile death-rate per 1,000 births was 18.7 under 1 year old, and 27.4 under 5 years old. In 75.2 per cent. of the deaths registered during the year the causes of death were not medically certified. The principal causes of death were diarrhoea and enteritis, debility, phthisis, and enteric fever; 84 deaths were recorded as having been caused by "the vomiting sickness," 21 of these being medically certified. On April 1st, 1912, there were 127 names on the Jamaica Medical Register.

THE so-called sleeping sickness of Nyasaland, called by the natives kaodzera, differs in certain respects from the typical sleeping sickness, and the Royal Society's Commission finds that the trypanosome present in the disease, though it resembles *T. brucei* and *T. gambiense*, is a distinct species identical with *T. rhodesiense* (Stephens and Fantham). These observations were reported to the Royal Society, and have been published in the *Journal of the Royal Army Medical Corps* (May, 1913). The authors of the report are Sir David Bruce, Majors D. Harvey and E. A. Hamerton, R.A.M.C., Dr. J. B. Davey, Nyasaland Medical Staff, and Lady Bruce. The parasite resembles *T. brucei* more closely than *T. gambiense*. It is concluded that the human trypanosome disease of North-East Rhodesia and Nyasaland is not the disease known as sleeping sickness in Uganda and the West Coast of Africa, and it is suggested that the native name kaodzera might be used for this disease to distinguish it from the older-known sleeping sickness.

CHRONIC PNEUMOCOCCAL INFECTION OF THE LUNGS IN CHILDREN.

BY

P. L. SUTHERLAND, B.Sc., M.B.,
BACTERIOLOGIST, WEST RIDING COUNTY COUNCIL;

AND

A. A. JUBB, B.Sc., M.D.,
ASSISTANT MEDICAL OFFICER OF HEALTH, WEST RIDING COUNTY COUNCIL.

SINCE the beginning of the routine examination of elementary school children there has been much discussion on the prevalence of phthisis in children of school age, and while some medical inspectors consider that the disease is uncommon in school children, or at any rate not to be diagnosed always with assurance, others seem to believe that the cases are numerous and easily recognized. Indeed the amplitude of variation in figures given in the report of the Chief Medical Officer of the Board of Education compels the suggestion that the prevalence of phthisis in certain school areas may be not so much a matter of pathological fact as of clinical bias. For instance, the percentage of phthisis among the Staffordshire school children examined in 1910 was 0.151; the percentage in Worcestershire was 4.183; and yet the phthisis death-rate of these counties was the same, namely, 0.7.

It is, of course, the experience of all clinical observers to see children in whom the pulmonary signs suggest the presence of phthisis. These cases present the following features: (1) they are often the sequels, immediate or remote, of measles or whooping-cough; (2) cough, sputum, and emaciation are variable in amount or degree; (3) tubercle bacilli are not found in the sputum. It may be argued that these are cases of mild or chronic phthisis. We are of the opinion that phthisis is seldom mild or chronic in children; it is too often acute and progressive, and from observations made both clinically and pathologically we have come to the conclusion that many cases in school children diagnosed as, or suspected of being, phthisical, are not such, but are chronic infections due to some other cause.

In support of this we shall first compare the result of the examination of the sputum from (a) suspected cases of phthisis in adults, and (b) suspected cases of phthisis in children from the age of 5 up to and including the age of 15.

During the past eighteen months ending October, 1912, the sputum of 1,824 adults has been examined in this laboratory for the tubercle bacillus. Of these, 33.7 per cent. gave a positive result. During the same period the sputum of 230 school children has been examined for tubercle bacillus, and of these 9.1 per cent. were positive. In not a single negative case among the children was a positive result obtained by a subsequent antiformin examination.

To carry the matter to a more definite state the negative sputum was in the cases of 69 suspected children inoculated into guinea-pigs. Of the 69 guinea-pigs only 2 were found tuberculous when killed at the end of one month, 30 died in from one to nineteen days of peritonitis more or less acute, and 37 were normal when killed after one month. This gives a percentage of 2.9 tuberculous.

Negative sputum from adults was inoculated in only 18 instances. Of these, 2 guinea-pigs were found tuberculous, 7 died of peritonitis, and 9 were normal—a percentage of 11 tuberculous.

Results of Examination and Inoculation: Adults and Children.

(a) Adults:	
Sputum examined and found positive	33.7 per cent.
Guinea-pigs inoculated and found positive	11.0 "
(b) Children:	
Sputum examined and found positive	9.1 "
Guinea-pigs inoculated and found positive	2.9 "

There is a remarkably close resemblance in the proportion of the two sets of figures, the ratio in each case being nearly 3 to 1; and though the number of adult inoculations is small, the fact of the ratio between the microscopical and biological results being practically the same in adults and children may be accepted as far as it goes. That is to say, we may take the figures for the microscopical

examination of the sputa as being a fair approximation and index of the incidence of phthisis in adults and children. This is not to assume that the disease is over three times (33.9 to 9.1) more frequent in adults than in children, but we are entitled to say that tuberculosis of the lungs is less common in children than in adults, and less common, too, than is generally supposed.

In view of this conclusion, which is quite reasonable in the circumstances, the question is: What is the nature of these cases of suspected phthisis in children? The probable explanation is that they are chronic catarrhal conditions associated sometimes with varying degrees of bronchiectasis and are the chronic residue of pneumonic or bronchopneumonic inflammations which in turn have followed measles or whooping-cough, or indeed may have arisen alone. The cough, the types of breath sounds, the râles, and the sputum, can all be explained by such a condition, and where dullness is elicited it is no doubt due to pulmonary fibrosis. Dr. Theodore Fisher has already, in this JOURNAL, called attention to the existence of peribronchitis and bronchiectasis in children, and has emphasized the need of caution before diagnosing phthisis in such cases.

Ages of 22 Positive Cases.					
Age 15	...	7 cases	Age 11	...	3 cases
" 14	...	3 "	" 10	...	2 "
" 13	...	2 "	" 9	...	1 "
" 12	...	4 "	" 8 (and under)	0	"
			Total	...	22 cases.

The fact that a positive result was obtained in only one case below the age of 10 and in twelve between that age and 15 suggests an increasing liability to phthisis as the age of childhood is left behind.

Out of 22 cases we have been able to trace 17. Of these in eighteen months 7 are dead, 6 are "worse" or "past relief," and 4 are indefinite, which supports our belief that phthisis in children is more apt to be acute and progressive than slow and benign. It must not be forgotten, however, that many of these suspected cases may give a tuberculin reaction through being infected with tubercle elsewhere, whence an invasion may take place later into the already damaged lungs, but this does not invalidate our contention that many cases of phthisis in children are falsely so called.

The next point to be considered is the fate of the inoculated guinea-pigs. The inoculation of the sputum from both adults and children resulted in a large percentage in peritonitis—namely, 38.8 per cent. from adults and 43.5 per cent. from children. In the negative sputa of children the pneumococcus was found in large numbers and was by far the most abundant organism present. The peritonitis in all the guinea-pigs inoculated with the children's sputa was, with one exception, pneumococcal. The organism was present in abundance, and was isolated from the peritoneal exudate.

The fact that from adults and from children about 40 per cent. of the inoculated animals died of a peritonitis which was pneumococcal in almost all the children's cases and in the majority of adults' cases—such a fact suggests that the pneumococcus has a special significance and virulence in the causation of these cases of suspected phthisis. Speaking with regard to children it seems to us that such a lesion—namely, a chronic infection by the pneumococcus—is the true explanation of many of those cases of suspected phthisis which live on from year to year, no doubt in a more or less debile condition, but able to go about and put in an irregular attendance at school.

CONCLUSIONS.

1. Phthisis is not so common in children as certain recent reports would appear to imply.
2. Many cases of "phthisis" or "suspected phthisis" in children are really chronic pneumococcal infections of the lungs inducing bronchial and peribronchial lesions.
3. These pneumococcal cases in children form a large percentage of the cases cured or improved in open-air schools.
4. It is extremely likely that such cases in children are treated at tuberculin dispensaries as phthisis in which the tubercle bacilli are not found in the sputum.
5. These pneumococcal infections tend to be chronic and non-fatal, though there is always the liability to tuberculous infection.

THE TREATMENT OF ADENOIDS AND ENLARGED TONSILS WITHOUT OPERATION.

By W. STEUART, M.R.C.S., M.I.E.E., ETC.,
MEDICAL OFFICER, X-RAY DEPARTMENT, QUEEN'S HOSPITAL,
BIRMINGHAM.

CONSIDERING the efficacy of x rays in reducing the size of chronically inflamed lymphatic glands it is surprising that hitherto there has been no attempt, as far as I am aware, to treat enlarged tonsils and adenoids by this method.

As the present surgical treatment is open to several objections, an investigation of the possibilities of the Roentgen rays was undertaken with a view to providing at least an alternative method which would be suitable in a large percentage of the cases now operated on. From inquiries made it would appear that the average number of hospital operations in London for this condition alone is not less than 500 a week.

Observations were at first confined to cases that had chronically inflamed lymphatic glands situated near the angle of the lower jaw and having enlarged tonsils at the same time. The treatment of the glands involved the radiation of the tonsils, so that there was no question of experimenting on patients until justification was obtained to proceed further.

All these cases showed a corresponding diminution in the size of both glands and tonsils. Patients were then selected in whom the condition of "tonsils and adenoids" was well advanced, and characteristic symptoms were marked—that is, mouth breathing, snoring, deafness, discharge from the ear, etc.

Each of the following examples is taken to show some special feature, with a view to the summation of the results to be obtained by radiation.

CASE I.

F. B., aged 10, had the typical adenoid type of face, mouth open, lower jaw dropped. Adenoids were made out by digital examination to be an exceptionally large mass, and the tonsils projected well beyond the pillars of the fauces. There was a discharge from the left ear. It was stated that the child suffered from colds and earache; that she could not hear properly, and that she snored at night.

As the result of the first dose it was reported that she could hear better and snored less, but the discharge from the ear continued.

After the second dose, discharge less and snoring less.

After the third, discharge from the ear stopped, snoring stopped, but child breathes heavily when asleep. Hearing much better. Mouth still open, but much less than at first.

Progressive changes were to be noted at each successive visit, the tonsils became smaller, but, though the adenoid tissue round the pharynx had shrunk sufficiently to give a good air passage, the change in actual size was not as much in this case as had been hoped for. This result was probably due to the prolonged neglect, the influence of which will be commented on in connexion with another patient.

The cases of two brothers are the subjects of the next examples, as their symptoms were much the same, differing only in degree.

CASE II.

James N., aged 11, had tonsils projecting beyond the pillars of the fauces, and adenoids could be felt as scattered masses. The father stated that the boy was constantly getting colds and was becoming increasingly deafer, so much so that arrangements had already been made to send the child to a school for deaf children, but the parents decided to seek further advice before giving their consent to this proposal. The boy snored when asleep and kept his mouth open.

Very little change was noted at first, but an improvement could be observed about a week after the second dose. A letter of inquiry to the schoolmaster elicited a reply to the effect that both boys had improved considerably lately, James, if anything, more than Joseph (the other brother).

Six weeks after the initial treatment the boy showed marked improvement, though he was still snoring a little and breathing through his mouth.

After seven weeks the snoring had stopped and the mouth was kept shut. A further letter from the schoolmaster stated:

"Joseph—Improved. Can hear from the back of the class."

"James—Improved. Can only hear when sitting right in front of class."

Although the boy's hearing was far from perfect, as shown by the above report, his condition was immeasurably better than when it was considered advisable to send him to a school for the deaf.

On examination at this time the tonsils were found to be well within the pillars of the fauces, but they had not quite