axillary crutches, which are employed with the mistaken idea that they give the spine some support. The use of these crutches has been well deprecated by Vignard and Monod,⁴ whose observations on them may well be repeated here: "They constitute one of the gravest faults in the old-fashioned spinal jackets. Their object is, as has been already said, without doubt to support the spine by the aid of the shoulders, but raise the shoulders and you will appreciate their inefficiency. Movement of the shoulders in the vertical direction which this action causes the axillae to undergo shows that it is only when having pressed up the shoulders to the extreme limit of their possible excursion that the axillary crutches can have any effect. These movements of ascent are partly limited by the serratus magnus muscle which, attached to the scapula, acts on the ribs and not on the spine. Only the lower slips of the trapezius muscle can furnish the means of useful action, and this action on the spine is really only an apparent one. The raising of the axillae is accompanied by a swinging movement of the scapula in its lower angle. The muscles which resist this swinging movement are the rhomboidei which are attached to the lower cervical and upper dorsal vertebrae, which vertebrae are thus drawn from above downwards. The exaggerated raising of the shoulders by the axillary crutches produces a compression of the spine and not stretching, which stretching is, on the contrary, only possible when the shoulders occupy the lower position.

"Untenable from anatomical considerations, on physiological grounds still stronger objections may be made to the use of axillary crutches. Even light, long-applied axillary pressure causes a strain which is uncomfortable, and may, indeed, be painful. The action on the respiratory movements is still more important. The raising of the shoulders has an enormous influence on the respiration. It corresponds to forced inspiration. The shoulders remaining high, it is impossible for the patient to accomplish complete expiration. Indeed, it is as bad to keep the thorax in a position of inspiration as it is to impede thoracic movements by compression. For these reasons, then, the use of axillary crutches should be abandoned, and the shoulders should be allowed to remain free, but pressed backwards and somewhat downwards."

The Completion of the Celluloid Jacket.

The splint having been cut out in the manner indicated, the edges may be bound and stitched with leather and suitable hooks, eyelets and buckles inserted into the celluloid. The splint is then to all intents and purposes completed. Nothing more is really essential, but it may be found advantageous to strengthen it with light steel as figured, though this is not by any means essential. From hygienic reasons it is desirable to punch numerous holes in the jacket to allow sufficient ventilation of the subjacent skin.

Such a jacket will be found of the greatest value to patients being treated in the convalescent stages of spinal caries. Its weight is negligible, it affords no obstruction to respiration or digestion, the support and protection it affords is ample, and the comfort and security it confers upon the patient renders it of unequalled value in mechanical treatment at this stage of tuberculous disease of the spine.

REFERENCES.

- ¹ Calvé: Quelques considérations sur les appareils amovibles en celluloid dans le traitement de la Coxalgie et du Mal de Pott. *La Clinique*, April 28th, 1913.
- ² Batten, F. B.: The Use of Celluloid Splints in the Treatment of Poliomyelitis, *Lancet*, 1912, vol. ii, p. 80.
- ³ Gauvain: The Use of Plaster of Paris in the Mechanical Treatment of Tuberculous Disease of the Spine, *Practitioner*, January, 1913.
- ⁴ Vignard and Monod: Le corset orthopédique, *Lyons Chirurgical*, December, 1909.

A CASE OF COMPLETE AURICULO-VENTRICULAR HEART-BLOCK.

WITH A REPORT OF THE PATHOLOGICAL CONDITION OF THE HEART.

BY

T. WARDROP GRIFFITH, M.D., F.R.C.P.,
PROFESSOR OF MEDICINE, UNIVERSITY OF LEEDS; HONORARY
PHYSICIAN, LEEDS GENERAL INFIRMARY;

AND

ALEX. MILLS KENNEDY, M.D.,
ASSISTANT TO THE ST. MUNGO (NOTMAN) PROFESSOR OF PATHOLOGY,
UNIVERSITY OF GLASGOW; SENIOR ASSISTANT PATHOLOGIST TO
THE GLASGOW ROYAL INFIRMARY; PATHOLOGIST TO THE
GLASGOW MATERNITY AND WOMEN'S HOSPITAL.

[WITH SPECIAL PLATE.]

IN the Schorstein Lecture which was delivered by one of us (T. W. G.) at the London Hospital last October, and published in the BRITISH MEDICAL JOURNAL on

October 12th, mention was made of a case of complete auriculo-ventricular heart-block which presented some points of special interest. The patient died of pneumonia, and the examination of the heart was undertaken by one of us (A. M. K.). A preliminary note of the condition found was given in the lecture. The examination has now been extended and completed, and it is believed that the record will prove of greater interest if the paragraphs and figures in the lecture dealing with

the case are reproduced. These were as follows:

The tracing Fig. 38 [p. 1204] is from the case of a member of our own profession, and was taken when the patient was 81 years of age. The ventricles were beating at 40 and with great regularity; as all the pulse periods are seven and a half fifths of a second, except the few which are marked otherwise. The auricular beats are rather less than double* the number of those of the ventricles; they have no constant relationship to the ventricular beats, and we have here the additional evidence of the block being complete, in the fact that the auricular rhythm is markedly irregular, but that there is no reflection of this irregularity in the pulse.

There is another peculiarity in the tracing to which I desire to call attention. When a premature contraction of the auricle occurs it is followed by an interval which is seldom longer than those which separate the systoles of the auricle which occur about the same time. Now it is possible that this may represent nothing more than a sinus arrhythmia; but as it is clear that there is a profound interference with the conducting functions of the heart, I venture to suggest the possibility of another explanation, which I admit is highly problematical, and which I advance purely in the hope that others may meet with cases to confirm or to confute it. We have already seen that in complete auriculo-ventricular heart-block premature contractions of the ventricle are not followed by any compensatory pause. Let us apply the explanation of this to the possibility of a sino-auricular block, as suggested by the lower outline in the diagram I have already used (Fig. 32). When

Fig. 32.

the sinus rhythm is dominant, we have seen that a premature contraction of the auricle is followed by a pause which is more or less complete according as the stimulus does not or does pass back to disturb the rhythm of the sinus. If, as the latter part of the diagram shows, the auricle may be supposed, in

* In the report of the lecture an unfortunate slip occurred, the word *half* appearing instead of *double*.

consequence of a block, to be emancipated from the influence of the sinus, a premature beat on its part would be followed by an interval of the usual interauricular length.

The patient was a man of active habits, and up to the time of his death, which occurred two years afterwards, was in the

muscles show some degree of fibrosis. There is fibrosis of the endocardium of the left ventricle just below the attachment of the mitral valve, and this fibrosis is distinctly evident over the situation of the auriculo-ven-

tricular bundle—namely, below and behind the pars membranacea septi. In this situation the fibrosis is seen to be continuous with the atheromatous thickening of the anterior mitral cusp along its attached margin bordering the anterior edge of the pars membranacea septi.

There are no other valvular lesions.

A block of tissue was removed from the cardiac septum, which contained the pars membranacea septi, the adjacent portions of the auricular and ventricular muscular septa, the central fibrous body, and the attachments of the anterior cusp of the mitral valve and the septal cusp of the tricuspid valve. The block extended sufficiently far back to include the mouth of the coronary sinus, and it was cut out in such a manner that its long antero-posterior diameter was at right angles to the long axis of the heart. Such a block includes the whole of the auriculo-ventricular node and main stem of the auriculo-ventricular bundle and a portion of its two principal branches. It was cut in serial sections from above downwards in the horizontal plane, and a series at intervals of five were mounted and stained with haemalum and van Gieson's stain.

Examination of the sections shows enormous dense fibrosis at the base of the mitral valve, extending outwards as a thickened atheromatous plaque on the ventricular aspect of the anterior mitral cusp and inwards to the auriculo-ventricular node. The fibrosis extends downwards into the ventricular septum as a very thick and very dense layer; this layer extends from the endocardium of the left ventricle inwards to the auriculo-ventricular bundle, forming a thick mass along its left side, and extending underneath it, and appearing also along its right side, though here it is less abundant. The thick fibrous layer on the left side of the bundle can be easily followed from the mitral valve to the aortic valve, at the base of which it again increases in bulk. The normal space between the bundle and cardiac tissue proper is almost completely obliterated, and, particularly from the left side, strands of fibrosis can be seen extending into the bundle about its middle from the greatly thickened sheath, and so breaking the continuity of the main stem.* The left side of the node and the whole length of the main stem of the bundle are thus enclosed in a dense thick sheath, the thickening of which underneath the bundle raises the latter up, so that on leaving the node it takes a curved course upwards and then downwards to enter the ventricle; this gives the convexity towards the auricle, which has been described as normal, but which in this case is exaggerated and certainly abnormal.

In the dense fibrosis at the base of the mitral valve, along the left side of the bundle, and at the base of the aortic valve, there are areas of calcification. In the auricular tissue above the node there is probably some excess of fat, and a short distance below the node there is some fatty infiltration in the bundle visible as numerous large globules. The small arterioles in the node and bundle and cardiac tissue generally show well marked thickening and considerable periarterial fibrosis. The principal vein from the node and its tributaries show recent thrombus within them—a terminal formation, but the main trunk is not completely occluded.

* The preliminary note in the report of the lecture is not quite accurate; the continuity is completely interrupted.

Fig. 38.

habit of walking some five miles a day. His daughter told me that as long ago as the early seventies she remembers her father being subject to peculiar attacks. Sometimes when making up medicine in his surgery he would suddenly, after apologizing to those who were present, lie down on the floor for a few minutes as if overcome by a feeling of faintness, and then resume his work as if nothing had happened. Of these attacks the patient had no recollection during all the time, some two years or more, that he was under my immediate observation. I saw him about eighteen years ago, when he had influenza, and my notes show that during the moderate fever which prevailed his pulse varied from 104 to 112. What the pulse-rate was in the interval I cannot tell. He had often had attacks of giddiness, but of late years these had got much less frequent, and for some time they had ceased. There was some tendency to cyanosis; the arteries were very atheromatous; the systolic blood pressure was as high as 260, and about the end of May of this year, though he was still rather a wonderful man for his age, and continued his active habits, he appeared to me to be failing. I was asked to see him early in June by his medical attendant. He was clearly suffering from pneumonia of a low type, and he died at the end of the sixth day. A tracing of the radial pulse on the third day of his illness (Fig. 39

Fig. 39.

shows that the pulse-rate had risen to about 50, and it never rose above 60. The upper tracing shows a premature systole, and the lower two in succession, without in either case a compensatory pause.

The pathological examination of the heart shows that there is a generalized hypertrophy of the cardiac musculature, and dilatation of both auricles and of right ventricle; the left ventricle is contracted in systole, and there is an excessive deposition of fat on the surface of the heart. There is some superficial atheroma of the aorta, and, about 1½ in. above its valves, there is a large thick atheromatous patch which is softened in its centre; the coronary arteries are slightly fibrosed.

The mitral orifice is normal in width but the tricuspid is slightly dilated. The anterior mitral cusp on its ventricular aspect shows a uniform atheromatous thickening of its basal half, and extending outwards in strands to some of the chordae tendineae, and at its leftmost edge, where it meets the posterior mitral cusp, there is a small area of calcification about the size of a lentil. This atheromatous area on the anterior mitral cusp extends inwards and spreads on to and involves the adjacent half of the pars membranacea septi, so that it must at least pass over the auriculo-ventricular bundle. Some of the chordae tendineae are thickened—a chronic atheromatous thickening—and the tips of the papillary

There is fibrosis and slight calcareous infiltration in the ventricle, particularly on the left side.

A block of tissue, to include the sino-auricular node, was next removed from the junctional region of the right auricle and superior vena cava, and cut in serial sections, which were mounted and stained as above described.

Examination of the sections shows that the excessive deposit of fat, already noted on the surface of the heart, infiltrates into the right auricle in the sino-auricular nodal region, causing considerable alteration. The nerve endings in this region are all conspicuous, and a considerable number of the ganglion cells are visible, but there is apparently a scarcity of the true "nodal tissue" fibres and

abundant fatty infiltration instead; there is also periarterial fibrosis.

The main bundle of the auriculo-ventricular junctional system being the path by which the auricular contractions are transmitted to the ventricles, it may be inferred from the pathological findings alone in this case that a condition of complete auriculo-ventricular heart-block was present during life. Regarding the pathological changes in the sino-auricular region, it does not seem justifiable in the present state of our knowledge to make a definite statement as to what their effects on the cardiac rhythm would be. One must simply record the facts, clinical and pathological together, and await further confirmation.

THE POSITION AND FORM OF THE NORMAL HUMAN STOMACH.

By A. MELVILLE PATERSON, M.D. EDIN.,

F.R.C.S. ENG.,

PROFESSOR OF ANATOMY IN THE UNIVERSITY OF LIVERPOOL.

It may be generally admitted that the normal human stomach exhibits many variations of form, dependent on temporary physiological or permanent anatomical causes.

From personal observations I am of opinion that the normal stomach is, in the majority of cases, a four-

cardia and pyloric antrum may form a definite tubular canal (Fig. 5), or it may be a deep V-shaped notch in the greater curvature (Figs. 3, 6, and 9), producing often a typical hour-glass contraction. There is usually a considerable thickening of the muscular wall of the organ in relation to the pyloric vestibule and the pyloric canal. Out of twenty stomachs taken consecutively from subjects in my dissecting-room the majority showed this form (Figs. 1, 2, 3, 4). All these subjects were preserved by the same method—an injection of a mixture of equal parts of pure carbolic acid, pure glycerine, and water.

The variations from this type are due to repression of

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.

Fig. 6.

Fig. 7.

Fig. 8.

Fig. 9.

Fig. 10.

Fig. 11.

Fig. 12.

chambered organ (Figs. 1-12) composed of the following parts:

1. Cardia and fundus (Fig. 1, c).
2. Pyloric vestibule (P. V.).
3. Pyloric antrum (P. A.).
4. Pyloric canal (P. C.).

The constriction forming the pyloric vestibule between

one or more of the parts referred to or to confluence of one with another of certain parts of the organ.

The commonest variation is due to repression of the pyloric antrum, so that the pyloric vestibule and pyloric canal form one continuous tubular passage (Fig. 7). Less common is the form in which the pyloric vestibule is absent and the cardia and pyloric antrum are confluent (Fig. 8).

T. WARDROP GRIFFITH AND A. M. KENNEDY: AURICULO-VENTRICULAR HEART-BLOCK.

J. S. B. STOPFORD: SHAPE OF THE NORMAL
EMPTY STOMACH,

Photograph of specimen. The outline of the stomach is
indicated by the continuous black line.