An Address

ON

THE CLINICAL IMPORTANCE OF THE SYMPATHETIC NERVOUS SYSTEM.*

BY

S. A. KINNIER WILSON, M.D., B.Sc., M.R.C.P.,
ASSISTANT PHYSICIAN TO THE WESTMINSTER HOSPITAL AND TO THE
NATIONAL HOSPITAL FOR THE PARALYSED AND EPILEPTIC,
QUEEN SQUARE, LONDON.

THE knowledge we possess to-day of the finer anatomy and of the physiology of the nervous system has been attained only at the expense of years of controversy. If there is much land still to be possessed, that which is now our heritage has been systematically explored and surveyed till we would fain believe it is a country with familiar and unchanging landmarks. Yet this is far from being the case, and while it would be a simple matter to find illustrations of the truth of this statement from other sections of neurology, we need not look beyond the sympathetic nervous system, which forms the subject of my remarks to-day, for a demonstration that will serve our purpose.

During the greater part of last century the anatomy and physiology of the sympathetic system was a *Haupttummelplatz* of neurological discussion, and it was only with the magistral researches of Gaskell and of Langley that order took the place of confusion and experimental data of empiricism. These English physiologists laid a scientific foundation for the study of the question, which remains unshaken, and their terminology has been accepted by the schools of other countries. As junior students of physiology we become familiar with the complex lore of vaso-constrictors and vaso-dilators, and even now we may not have forgotten our quondam interest in the behaviour of the vessels of the rabbit's ear under varying experimental conditions; but as practitioners of medicine it may be questioned whether we concern ourselves much with the application of our hardy-gained knowledge to the common clinical phenomena of disease that must daily obtrude themselves upon us. A little thought will convince us that disturbances in the realm of the sympathetic nervous system are probably more frequent than any other class of morbid symptom, and even a slight acquaintance with their interpretation would stand us in good stead.

But our knowledge of the functions of the sympathetic system has been notably modified and enlarged by recent discoveries of importance. Advances that have been made in regard to the nature and physiology of the ductless glands have thrown a flood of light on the former, and this we cannot afford to ignore. We are compelled to regard the sympathetic nervous system from a fresh viewpoint: the main principles of its action must be restated, and it is no exaggeration to say the gain thereby to clinical medicine is bound to be far-reaching. This is one of the aspects of the question to which special attention will be directed.

Before entering on it, however, I may be allowed to remind you of the salient features of the sympathetic system, as established by the work of Langley and various other investigators.

ANATOMICAL AND PHYSIOLOGICAL CONSIDERATIONS.

The sympathetic system, embryologically, is directly derived from the cerebro-spinal nervous system. As early as the sixth week of fetal existence the general development of the sympathetic system is complete, in the sense that its essential features can be traced even at that stage. It may be said very briefly that the ganglion cells of the sympathetic issue from the mass of nerve elements forming the neural crest, that they migrate thence at differing times and rates, those of the thoracic region always being in advance of the cranial and lumbo-sacral regions, and that on reaching their approximate final positions they proliferate and clump themselves into ganglion formations. The processes of the cells elongate into fibrillae and so a communicating network of fibres is produced, extending centrally to the cerebro-spinal system and peripherally to viscera. A number of sympathetic

cells are left behind, however, in the cerebro-spinal axis, and later we shall see how the central connexions of the sympathetic will serve to explain the appearance of sympathetic phenomena in central lesions.

When full development is attained, the sympathetic system is found to consist of two divisions. The first is the sympathetic system proper, often called the vertebral sympathetic, consisting of the chain of ganglia of the cervical, dorsal, and lumbar regions. The cervical set is collected into three sympathetic ganglia, but in the case of the thoracic and lumbar there is a ganglion corresponding to each segment of the spinal cord. These ganglia are connected, on the one hand, with the spinal cord at their own levels, except that the central connexions of the cervical ganglia arise, not from the cervical cord, but by the first and second dorsal roots; and, on the other hand, from these ganglia fibres pass peripherally, as a rule through the medium of secondarily formed plexuses or ganglia, to all the viscera of the thoracic and abdominal cavities, to the vascular system, to the skin with its glands and other appendages, and to the pelvic viscera and genitalia.

The second division of the sympathetic is known as the parasympathetic, or, preferably, the autonomic system. At the cranial and caudal extremities of the cerebro-spinal axis are situated three sympathetic centres, two at the former and one at the latter. These centres are in the cerebro-spinal system, and from them sympathetic fibres pass out, again through the medium of intercalated ganglia, to all the viscera of the body, and to all the structures enumerated above as supplied by the vertebral sympathetic. In the case of the autonomic system there is only one set of ganglia, and they are to be found, as a rule, in the immediate vicinity of, or actually in, the organs innervated, whereas the vertebral sympathetic system, as has been said, possesses two sets—one the vertebral chain, and the other the plexuses, coeliac, splanchnic, and so on. In the case of the cervical sympathetic, however, there is only one set of ganglia on the path to the structures innervated by it.

The sympathetic nervous system, therefore, presents this remarkable feature—that it gives a double nerve supply to the skin and its appendages, the glands, vascular system, the alimentary system from mouth to rectum, and the respiratory system, and the importance of this fact will be the more appreciated when it is stated that there is much evidence forthcoming to show that the two divisions—the vertebral sympathetic and the autonomic—are mutually antagonistic in their function, as we shall see subsequently.

Let us look at the sympathetic and autonomic supply a little more closely. It should be premised that not all of the anatomical details about to be mentioned have been satisfactorily established, and there is difference of opinion on some not unimportant points. As far as is agreed on at present, it may be said that the arrangements are as follows:

The autonomic system has three centres, the mesencephalic, the bulbar, and the sacral, or lumbo-sacral. From the first of these pass autonomic fibres by the oculo-motor nerve and ciliary ganglion to the sphincter iridis, constricting the pupil in their action. The corresponding vertebral sympathetic supply, antagonistic in its function, is constituted by pupil-dilating fibres from the cervical sympathetic in the neck, reaching the eye by a circuitous route via the carotid plexus and the superior division of the fifth or trigeminal nerve.

The bulbar autonomic centre is formed by sympathetic cells in relation to the facial, glosso-pharyngeal, and vagus nerves. Connected with the first two of these are the fibres passing out to supply mucous and salivary glands, the blood vessels of the mucous membranes of mouth and nose, the lacrymal glands, and so on. These are antagonized by vertebral sympathetic fibres running in the cervical sympathetic and reaching the same distribution by the elaborate route already described. We may not have forgotten the differences between the autonomic "chorda saliva" and the "sympathetic saliva" of our student days. In the autonomic facial fibres, probably, are vaso-constrictors to vessels and sweat glands, in the cervical sympathetic are vaso-dilators, and so on. But I shall not particularize further, lest we should be unable to see the wood for the trees.

* Read at the quarterly meeting of the East Anglian Branch at Colchester and illustrated with lantern slides.

The vagus centre, however, deserves our attention because of its great clinical importance. There is reason to believe it is the dorsal nucleus of the vagus which forms the autonomic vagal centre, but we need not delay over this vexed question. The vagus supply is extraordinarily wide. It furnishes sympathetic fibres, belonging of course to the autonomic system, to the heart and blood vessels generally, to the alimentary tract from mouth to colon, to the liver, pancreas and kidneys, to larynx, trachea, bronchi and lungs; its branches go both to smooth, non-striped muscle fibres in these various vessels and viscera, and to glandular organs therein contained. The autonomic vagal action is antagonized in all these—again it must be remarked that this is a generalization the truth of which has not as yet been established in every instance—by vertebral sympathetic fibres supplying the same structures and arriving via the visceral plexuses. Thus, to take the best known instance, there are cardio-inhibitory fibres in the vagus and accelerating fibres in the sympathetic. Again, the vagus contracts and the vertebral sympathetic dilates the coronary vessels. The sympathetic contains the constrictors for the blood vessels, the vagus the dilators, if there are such. In the sympathetic are the fibres for the sweat glands of the body, probably stimulating and not inhibiting fibres. The sympathetic fibres relax the smooth muscle fibres of oesophagus, cardiac sphincter, and stomach; the vagal fibres contract them. The latter contract the gall bladder; the former relax it. And so on.

Finally, in the lumbo-sacral part of the spinal cord is the third autonomic centre, from which fibres pass to the vessels of the rectum, anus, and external genitalia—vaso-dilator in function. The vaso-constrictors to the same are in the vertebral sympathetic system. Again, the sacral autonomic supplies the muscles of the colon, rectum, anus, urethra, bladder, and external genitalia, being chiefly concerned with contraction, while the corresponding sympathetic fibres are chiefly concerned with relaxation.

THE CENTRAL CONNEXIONS OF THE SYMPATHETIC.

A further anatomical matter that ought not to be passed over is the question of the central connexions of the sympathetic. It has been shown that the vertebral sympathetic system is an outgrowth from the cerebro-spinal. The efferent fibres joining the two arise from a definite tract running the length of the spinal cord from the lowermost cervical region to the lumbar. This is known as the intermedio-lateral tract, and it is composed of cells, readily distinguishable from those of the anterior and posterior horns, which are situated at the lateral extremity of the grey matter, where anterior and posterior horns meet. The nerve cells of this tract rise and fall in number in a more or less regular way for each segment of the cord. From them pass out efferent sympathetic fibres to join the chain of sympathetic ganglia, travelling by the anterior roots, and leaving the latter by the white rami communicantes immediately after the formation of the mixed nerve.

Now, it is inconceivable that the spinal sympathetic tracts should have no connexion with higher centres, and although it has been doubted whether there can be cerebral centres for the functioning of vasomotor nerves, sweat glands, the alimentary tract, and so on, we do know as a matter of fact that the spinal tracts already referred to are in association with bulbar sympathetic centres—not the autonomic bulbar centre of which we have been speaking—whose existence has been demonstrated in a conclusive fashion. And why should the system not be represented in the cerebrum itself? We may admit that the sympathetic system is very little under the influence of the will, but it is notably under the influence of moods and emotions. Anger, fear, shame express themselves outwardly by obvious sympathetic disturbances—by pallor, blushing, perspiration, dryness of mucous membranes—to mention but a few of the symptoms which will occur to every one; and if we believe, as there is evidence to show we should believe, that the emotions are placed at least as high as the basal ganglia, then the bulbar sympathetic centres must be represented anatomically still further cerebralwards. I shall refer to clinical cases which appear to bear this out a little later.

The sympathetic system further resembles the cerebro-spinal system in that it possesses afferent—that is,

sensory—as well as motor, or efferent, connexions with the latter. There is excellent evidence forthcoming to prove that the viscera have a sensibility of their own, impulses underlying the appreciation of which are conveyed by afferent sympathetic fibres leaving the sympathetic chain and passing by the posterior roots to the spinal cord like any of the other sensory impulses. Normally we are quite unconscious of the functional activity of our organs, and this no doubt suggests that sympathetic afferent sensory impulses do not reach the brain and become conscious, or, rather, so that we become conscious of them. At the same time, we do well not to negative the possibility of such an occurrence.

It is of course well known, as I may venture to remind you, that in certain diseased states of viscera sensory impulses from them reaching the cord are ultimately interpreted by our consciousness as proceeding from the area of skin supplied by the ordinary spinal nerves running into the same segments of the cord as are reached by the abnormal sympathetic afferent impulses. Thus in angina pectoris the patient may have paraesthesiae apparently in the distribution over the skin of the inside of the left arm and hand. In gastric ulcer he may have hyperaesthesia over the skin of the upper abdomen on the left. Other instances will at once suggest themselves. These zones of referred pain have been carefully worked out by Head and others, and they are of localizing value. But that there is a possibility of actual sympathetic sensory impulses reaching consciousness in certain diseased states seems to me the only satisfactory explanation of some of the symptoms of neurasthenia and also of hysteria.

From what has been said it will appear that the sympathetic system is at once dependent on, and independent of, the cerebro-spinal system. Owing to the central connexions which the former possesses, there may develop obvious sympathetic symptoms in organic and functional disease of brain, spinal cord, or peripheral nerves. What of its possible independence?

We have seen that a very large part indeed of our somatic structure, our circulatory system, respiratory system, alimentary system, genito-urinary system, the skin, mucous membranes, glands, and so on, are as it were slung between the nerve supply of the autonomic system on the one side and the sympathetic on the other, and that these are mutually antagonistic. It may be concluded that the result is the constant maintenance of tone in these organs, a tone which is constantly varying, but which is not constituted by alternating periods of rest and unrest. There is always activity in function. Is it possible that sympathetic ganglia have an autonomous activity of their own, and that there is a peripheral independence in them not possessed by any of the organs reached by the cerebro-spinal system? It would appear to be not merely possible, but probable. There are short reflex arcs in certain hollow viscera the physiological activity of which appears to be independent of central connexions. The movements of the ureters, peristaltic action of the intestine, and the mechanism of the heart-beat, are illustrations in point. And while, as far as I know, a wider application of the principle has not been made, it seems possible that not a few of the phenomena seen in vasomotor neuroses, to take but one example, may be due to increase or decrease of peripheral sympathetic activity produced by morbid agents whose action is quite independent of the central representations of the sympathetic system.

SYMPATHETIC SYMPTOM-COMPLEXES.

Enough has been said to support my original contention, that disturbances in the realm of the sympathetic nervous system as a class of morbid symptom must confront us almost daily, and I should like now to sketch a few of the symptom-complexes, the significance of which can be understood at once by reference to the anatomical and physiological data we have been considering.

First of all, we shall look at the occurrence of obvious sympathetic symptoms in central cerebral lesions.

Reference has already been made to the familiar symptomatology of the expression of the emotions. Under the influence of an emotion we turn pale, we blush, we perspire, we shiver or shudder; under the influence of emotion the throat and buccal membranes of the public speaker become dry. Under emotion we experience a

"thrill" all over our body; our cheeks "tingle." All these clearly are somatic phenomena produced through the sympathetic system; the last is the pilomotor reflex, known no doubt to many of my audience.

Now it seems reasonable, if not indeed essential, to suppose that if these generalized somatic symptoms are produced, as they clearly are produced, through the agency of the autonomic and sympathetic system, then the vagus centre and the bulbar sympathetic centre must be capable of being acted on from higher levels. For the stimuli evoking the emotion pass in to the brain centres by such sense avenues as the visual or the auditory path, and the lowest point at which such sensory impulses could pass over from the afferent to the efferent side is in the mid-brain, which is definitely above the bulbar level of which we have been speaking. There must, therefore, be a higher representation of the sympathetic centres than the medulla, and there is evidence to show that this representation is at least as high as the basal ganglia, the optic thalamus and corpus striatum. I have seen an ordinary case of left hemiplegia of capsular origin in which there was a permanent "goose-skin" on the affected side of the body. I have seen another case of hemiplegia in which the affected limbs were the seat of a vasomotor paresis with oedema, and such cases are not really very uncommon. They are often met with in infantile cerebral hemiplegia. In such instances the lesion is certainly not lower than the pons, and is probably higher. It is patent, therefore, that both the autonomic and the vertebral sympathetic systems are centrally represented, and though this has been denied by one or two investigators it is difficult to set aside the evidence which has accumulated.

We are on surer ground when we come to the bulbar sympathetic centres, the centres in the medulla oblongata.

We may take the autonomic vagal centre first.

There is a definite symptom-complex associated with disturbance of the vagal centre to which it is well to direct special attention. The condition clinically is known to our American friends as a "nerve storm," and the description is not inapt. It is known also as pseudo-angina, cardiac neurasthenia, and under other designations, but the term which Sir William Gowers proposed, and which I much prefer to use, is "vaso-vagal attack." The attacks are usually periodic, often resembling epilepsy in their periodicity, and it is quite possible that in some instances, at least, they are of the nature of epileptiform discharges in these lower centres. Commencing as a rule with an epigastric aura, the attack rapidly develops into palpitation, forcible or racing cardiac action, sometimes almost cessation of the same, suffocation, and difficulty in respiration; with this the patient usually experiences an indescribably distressing feeling of impending death, or of impending misery of some sort; at the same time the extremities are as though of marble in their icy coldness, then to this may succeed clammy perspiration or burning flushes all over, and at the same time there is often trembling and quivering of the musculature. The attack may last a few minutes, or it may be drawn out over hours. It may be complete, after the description just given, or it may be very partial and incomplete. In either form the condition is far indeed from uncommon, and I have personally seen many instances of it. It is self-evident, I think, after what we have learnt, that the whole vaso-vagal attack is a pure symptom-complex of a functional disturbance in the vagal autonomic centre, and as such it ought to be known. Nothing is more desirable, more imperatively called for, than to pick such attacks out of the rubbish heap of neurasthenia, with which they are constantly confounded, for the treatment of the two entities differs widely. Of course such an attack may occur as an episodic syndrome in the course of another disease, as can be well understood, but in its pure and uncomplicated form it is frequently met with, even in young people, and has scarcely received the attention it deserves. Its treatment with nitro-glycerine, in any form, but especially the liquor trinitrini, is often strikingly successful.

Apart from the vagal autonomic centre in the medulla there is a sympathetic centre in its neighbourhood—almost certainly in the substantia reticularis—which is a bulbar representation of the vertebral sympathetic system, and anatomically in continuity with it, or at least with the cervical part of it. As a result of a lesion involving

this part of the medulla, an interesting series of sympathetic symptoms may develop of central origin. The region concerned is in the distribution of the posterior inferior cerebellar artery, which I may remind you is a branch of the vertebral and winds round the side of the medulla on its way to the cerebellum via the inferior cerebellar peduncle. As it winds round it sends in small terminal branches to the medulla, which penetrate its substance, and if the vessel is occluded from thrombosis—a not infrequent lesion—the area of the medulla supplied often suffers from ischaemia, and among the characteristic symptoms of this complex is paralysis of the cervical sympathetic nerve and loss of sweating on the face on one side.

When we come to the spinal cord, one of the common clinical pictures is that of irritation or of paralysis of the cervical sympathetic nerve, which arises from Budge's centre in the lateral horn of grey matter of the first and second dorsal segments, mainly, and passes out to follow that circuitous route in the neck and head to which reference has already been made. We have seen that there must be a continuation cerebralwards of this centre, for identically the same symptoms can be produced by a lesion of the posterior inferior cerebellar artery, as far as the cervical sympathetic is concerned, as by a lesion of the upper dorsal cord, or a lesion of the nerve trunk itself. The localizing value of paralysis of the cervical sympathetic is often precious. It reveals itself by a very clear-cut syndrome, consisting in ptosis, myosis, and enophthalmos, with or without vasomotor and sweat disturbances over the face and upper part of the trunk. A rare concomitant of cervical sympathetic palsy, which, however, has come under my notice at least three times, is heterochromia iridis, a pigmentary change taking place in the iris on the affected side. Stimulation or irritation of the cervical sympathetic naturally produces an opposite effect—namely, exophthalmos, widening of the palpebral aperture, and mydriasis. Sometimes almost the only sign of a commencing palsy from defect of this nerve path is failure of the pupil to dilate to shade. It is well worth remembering, also, that cocaine may be used as a test of the integrity of the cervical sympathetic; a drop instilled into the eye will cause the pupil to dilate only if the nerve be normal; atropin or homatropin is of no value in this connexion.

Many organic lesions of the spinal cord are accompanied by obvious vasomotor and trophic changes, which are to be regarded as due to implication of the intermedio-lateral tract. In compression paraplegia from whatever cause arising it is often possible to detect the level of the compression by the simple device of injecting pilocarpin, a sixth of a grain, and observing, adequate precautions having been taken to protect the skin surfaces with a cradle, whether the drug has failed to produce sweating in the lower limbs and up to a given level on the trunk. A valuable case of poliomyelitis has been recorded where after the onset there was complete anhidrosis of the lower limbs, and at the necropsy a haemorrhage was found in each intermedio-lateral tract. The vasomotor symptoms that often accompany transverse myelitis, spinal tumour, and other spinal cord conditions that need not further be specified, find a ready explanation in involvement of the important intermedio-lateral tracts.

The symptomatology of organic disease of the thoracic and abdominal sympathetic system, outside of the cord, is not at all well known, and yet evidence is steadily accumulating which will help us some day to differentiate more clearly between the various ganglia and plexuses than we can at present. Of course, it is recognized that many of the familiar visceral crises of tabes are occasioned by organic disease of the sympathetic plexuses supplying the viscera in question. Again, the abdominal symptoms of lead colic may be the result of irritation of the solar plexus. A syndrome of the solar plexus has been described, consisting in hyperaemia of the abdominal viscera, vomiting, diarrhoea, oliguria, and general loss of strength. Certain forms of visceral neuralgia, to which attention has again been drawn recently, may be associated with affections of the coeliac plexus. The relation of mucous colitis, of many of the symptoms of acute peritonitis, of certain forms of angina pectoris, and certain acute pulmonary conditions—to mention but a few—to disease of the thoracic and abdominal sympathetic

plexuses is a chapter of clinical pathology that has yet to be written.

THE PHARMACOLOGICAL ASPECT OF THE SYMPATHETIC SYSTEM; RELATION TO DUCTLESS GLANDS.

We must leave this part of our subject, however, only the fringe of which has been touched, to pass on to a view of the whole question which is eminently calculated, with the aid of further research, to lead to most fertile results. I refer to what may be called the "pharmacological" aspect of the sympathetic system, and the relation of the function of the latter to the function of the ductless gland system.

To appreciate this new standpoint we must begin by alluding to what is now called the chromophil or chromaffin system of gland cells. It has long been recognized that the cortex of the suprarenal gland is distinct from the medulla of the same. The medulla is constituted by cells which possess this peculiarity, that they stain deeply when soaked in the salts of chromic acid—for example, potassium bichromate—becoming a rich brown colour. So characteristic is this reaction that such cells can readily be picked out in this fashion whatever be their situation; hence the expression chromaffin, or better chromaphil, to signify them. Now these chromaphil cells are found in many other places beside the medulla of the suprarenals. From the embryological standpoint they are known to originate in the same situation or *Anlage* as the cells which eventually reach and form the sympathetic chain of ganglia. In fact, we may speak of sympatho-chromaphil tissue. In every ganglion of the chain, then, on each side, and mostly lying in depressions on the dorsal portions of these ganglia, are rounded masses of chromaphil tissue, of exactly the same nature as the medulla of the suprarenal. Further, there is an unpaired strand of this tissue lying on the front of the abdominal aorta, and known as the aortic body or paraganglion of Kohn. Then they are found not merely in the sympathetic ganglia, but as a constant occurrence in many of the sympathetic plexuses, and they have been traced in the kidney, ureter, accessory suprarenals, prostate, ovary and epididymis, and in the carotid bodies. In the chromaphil system, therefore, we have a widespread tissue, intimately allied to the sympathetic system in origin and localization, which not only presents the histological peculiarities already specified, but is endowed with equally important physiological properties. Chromaphil tissue secretes a substance which has a definite and characteristic action on non-striated muscle fibres—in other words, on structures which are innervated by the sympathetic system.

No modern drug is more familiar to the physician than adrenalin. The important point about it is that adrenalin, which is the product of the secretion of the suprarenal medulla—that is, is derived from the sympatho-chromaphil system—acts solely on tissues innervated by the sympathetic. This is a striking fact. What are the effects of injection of adrenalin, or preferably "adrenin," since adrenalin has commercial associations? It causes a rise of blood pressure; it dilates the pupil; it contracts the muscles of the hairs; it excites the salivary and lacrymal glands, the liver, the muscular tissue of the uterus and vagina, of the vas deferens, the vesiculæ seminales, the dartos muscle, and the muscular coat of the stomach. In no case does extract of adrenal correspond in its effect to what is produced by stimulation of cranial or sacral autonomic nerves; on the other hand, its effects are such as are produced by stimulation of some one or other sympathetic nerve.

There is, therefore, normally present in the body a secreting system which is normally producing a secretion with a definite pharmacological action on sympathetically-innervated tissue, and this tissue is thereby kept in a normal state of tone. If the secretion is defective, there is loss of sympathetic tone; if it is excessive, there is a pathological increase in sympathetic tone. In fact, our German *confrères* are already speaking of sympathicotonus and vagotonus, although, so far as I am aware, the expressions are not as yet employed in this country. Further, they speak of "adrenalin sensibility" and are using injections of adrenalin as a test for the functioning of the sympathetic system. The clinical and therapeutic importance of these facts must be at once apparent to us.

The question arises as to how adrenin acts, whether on

sympathetic centres or on sympathetic end-mechanisms. It has been shown that it acts mainly on end-mechanisms, and that, further, there are substances which are antagonistic in their action to adrenin. For instance, adrenin has no special action for the autonomous system, as has already been pointed out. The autonomous system, however, is stimulated by cholin and analogous poisons and paralysed by atropin. The sympathetic system is indifferent both to cholin and atropin. It would appear, therefore, that in the body are normally found certain "poisons"—for example, cholin and adrenin—which have autonomous-exciting and sympathetic-exciting actions respectively, which are mutually antagonistic and can compensate one another.

But the matter does not end here. We know that other internally secreting glands are controlled by the same system—to mention only one, the thyroid. The thyroid is supplied with vaso-constrictor and vaso-dilator nerves; on these, whichever is the sympathetic group (for there are differences of opinion on this matter), the internal secretion of the chromaphil system, it may be assumed, has a stimulating action. Again, between the suprarenals and the pancreas there is undoubtedly a very intimate relation. It has been shown, for instance, that neither puncture diabetes nor pancreatic diabetes can occur after removal of the suprarenal bodies. It has been shown, further, that in pancreatic diabetes adrenalin dropped on the conjunctiva will dilate the pupil, whereas it will not dilate it in this way in the normal eye, although these experiments require confirmation.

It is undesirable, however, though it would be easy, to amplify these interesting and difficult topics. I may recapitulate in a sentence and say that a substance normally secreted by the adrenal medulla passes into the general circulation and maintains the tone of all sympathetically-innervated tissue, including other ductless glands. The cells which secrete this substance are embryologically of sympathetic origin. Other substances are produced in the organism which have an antagonizing, neutralizing, or compensating effect when compared with that of the adrenal secretion, and they act rather on the autonomous system. Similarly, many commonly employed therapeutic agents have an action on one or other of these groups. Cocaine, atropine, caffeine, stimulate the sympathetic system and inhibit the autonomic system. Morphine, chloral, antipyrin inhibit the sympathetic system and stimulate the autonomic system.

APPLICATION OF PHYSIOLOGICAL FINDINGS TO CLINICAL MEDICINE.

The linking up of the sympathetic system and the ductless gland system is already proving to be a most important generalization in clinical medicine, and with increasing knowledge this importance will assuredly be intensified. There is a large group of diseases known as the angio-neuroses, included in which category are such conditions as Raynaud's disease in mild or severe variety, local pallor or "dead finger," local asphyxia, symmetrical gangrene, acroparaesthesia, scleroderma, angioneurotic oedema, and so on—diseases obviously associated with disturbances of the sympathetic innervation of the vascular system. Now the frequency with which affections of one or other of the ductless glands are noted in such cases is remarkable. Disease of the thyroid is practically constant in scleroderma. Myxoedematous patients constantly complain of feeling cold, and exhibit a malar flush; they often complain of acroparaesthesia. In Graves's disease vasomotor irritability is a cardinal feature. In fact vasomotor neuroses and disturbance of functions of other sympathetically-innervated tissue is an association that is no more than might be expected in view of what has been already said.

Again, many of the common symptoms of neurasthenia and hysteria are patently of sympathetic origin. Who of us has not seen the typical irregular red blotches appear on the skin of the neck and face as the neurasthenic subject "works himself up into a state"? The clammy hand, flushed or pallid features, dilated pupils, the innumerable paraesthesiae, the unwonted sensations in head or body, are surely of sympathetic parentage. In not a few cases of neurasthenia symptoms of this class are the chief or only manifestation of the disease. Here, then, is a condition of defective sympathicotonus; may it not have

much to do with impairment of function of the chromaphil system? Addison's disease, for all its continued obscurity, may be regarded as at least to a great extent attributable to disease of the suprarenals. One of its characteristic features is asthenia, which is sometimes profound. And there does not appear to me any tenable distinction between the asthenia of Addison's disease and the asthenia of neurasthenia. Cases of the former are not infrequently diagnosed as ordinary neurasthenia at first. It is difficult to avoid the conclusion that defect of glandular function is responsible for much in the clinical picture of neurasthenia.

In the familiar symptoms of the climacteric how prominent are those which are derived from the sympathetic system. Palpitation, noises in the ears, "rush of blood to the head," hot and cold feelings, mist before the eyes, sweating, trembling, paraesthesiae of all sorts, sinking feelings, vertigo—who does not know the whole gamut of climacteric symptomatology? Analysis indicates at once the sympathetic origin of practically all of it, but why should this branch of the nervous system be disturbed? The answer is ready to hand. At the climacteric a physiological involution of glandular activity occurs, ovarian function alters, and thereby the function of other glands is thrown out of gear. Sympathetic tone is dependent on adrenal support, and until the glandular equilibrium is once more attained sympathetic symptoms are likely to occur. Let it not be supposed that the male sex is free from such glandular upset. The age of 63 was regarded by those of another generation as the crucial year for the male; but it is not necessary to postulate an actual climacteric—it suffices that the physician should be able to interpret the symptoms complained of in the light of what has been said.

The protean phenomena of hysteria can in many instances be readily explained by reference to sympathetic function. Not only are vasomotor symptoms of very common occurrence in that disease, but a vasomotor theory has even been advanced to account for the whole symptomatology. It is impossible, obviously, to subscribe to such a view; at the same time the tendency to-day is to ignore the somatic symptoms of the disease, especially those of vasomotor origin—a tendency which should, in my opinion, be discountenanced. In hysterical paraplegia one often finds a vasomotor paresis; in hysterical anaesthesia failure to draw blood on pricking the skin is as easy to demonstrate to-day as it was 300 years ago. The results of such a failure, however, are not quite so serious from the patient's point of view as they were in the "good old days." In some cases of hysteria multiple cutaneous gangrene, so-called, is known to occur, and I have seen a case where the diagnosis of a somewhat similar skin condition was attended with the greatest difficulty. Dermatographia or urticaria scripta, no doubt, occurs as a more or less isolated phenomenon, yet in hysteria, above all other conditions, it is seen in marked degree.

The pathogenesis of the sympathetic impairment in hysteria is not easily elucidated. This is not the time to venture even on a brief attempt to sketch an adequate theory of hysteria, but I may remark that the occurrence of unmistakable sympathetic symptoms in that disease argues for the central representation of the sympathetic system, and the central control, in certain not yet clear conditions, of the ramifications of that system in viscera and blood vessels.

Many more conditions might be referred to in this connexion. I have said nothing of asthma, of migraine, even of transient symptoms which can scarcely be said to constitute a disease, such as blushing, local sweating, etc., although it may be remarked that in these and other morbid states of a like kind good results are being obtained by treatment on opotherapeutic lines. I have given only a brief account of some of the chief features of interest revealed by a study of the sympathetic system and its relations, but it may serve to show that the fresh light which is being thrown on sympathetic function by recent investigations exemplifies once more the mutual interdependence of scientific research and medical practice.

THE late Mr. Charles George Lee, Honorary Surgeon, Liverpool Eye and Ear Infirmary, left estate valued at £7,638.

VARIETIES AND TREATMENT OF CHOREA.*

BY

FREDERICK LANGMEAD, M.D., F.R.C.P.,

ASSISTANT PHYSICIAN TO THE HOSPITAL FOR SICK CHILDREN, GREAT ORMOND STREET, TO THE ROYAL FREE HOSPITAL, AND TO THE SEAMEN'S HOSPITAL, GREENWICH.

CHOREA is insidious in its onset. The child who becomes the subject of chorea is usually nervous, excitable, and imaginative. She is easily frightened. At school, though by no means a dunce, she lacks concentration of thought and perseverance. Her attention wanders from the work to the creations of her active mind. It is easily diverted by all manner of extraneous and trivial circumstances.

Chorea begins as an accentuation of this psychical unrest. Before the actual movements appear the patient becomes more excitable, more irritable, more emotional, and less reliable. The distinction between when the mental state should be called choreic and when it should be regarded merely as evidence of a neurotic disposition is subtle. There are some who look on the many symptoms of nervous instability to which choreic children are liable as evidence of a rheumatic toxæmia, whilst others speak of chorea as especially prone to attack neurotic children. A compromise between these opinions is probably nearer the truth, for, if we accept the former view, we must acknowledge a persistent rheumatic toxæmia dating from very early life, whilst if we ignore it we shut our eyes to the very obvious psychical disturbances which a definite attack of chorea brings with it.

As evidence of the ill-balanced and hyper-excitable mental state is the frequent occurrence in the history of the same child of such functional errors as night terrors or day terrors, somnambulism, enuresis, nervous diarrhoea, and tic.

Long before the chorea is recognizable by its erratic movements the child's school work deteriorates. It becomes inaccurate, slovenly, and uneven. Perhaps this is best shown in writing and arithmetic. The letters and figures are badly formed, irregular and wavy, crowded in some places, spaced in others, and sloping at various angles. This falling off is readily detected by the teacher, who metes out punishment which he believes to be condign, but which is, of course, unmerited, and may precipitate symptoms of a more obvious form. The attack is then not unnaturally laid by the parents at the teacher's door.

Forms.

In almost every case, in addition to the erratic pseudo-purposive movements and the psychical changes, there are also some loss of power and some sign of cardiac affection. The loss of power is denoted by the characteristic crouching attitude when the child sits up, due to weakness of the back muscles and those which hold the head erect. The grip is feeble, and there is atony of the muscles and laxity of joints. The signs found by examination of the heart in many cases are merely those which may accompany functional disturbance—rapidity, a short, sharp, and rather slapping first sound, approximation of these sounds, and sometimes irregularity. A slight systolic murmur and slight dilatation may be present, and probably indicate a local rheumatic toxæmia or infection, especially if a mid-diastolic murmur or so-called reduplicated second sound are heard at the apex. In many cases signs of previous valvular disease are found.

The mental state when the chorea is readily recognizable is usually ill-balanced and the emotions are beyond control. It has been aptly likened to April weather, sunshine and shower alternating in quick succession. Tears flow as part of real distress for which there is no adequate reason or perhaps no reason at all.

Each of these four factors which contribute to the clinical picture of chorea—namely, erratic movements, psychical disturbance, weakness, and cardiac changes—may be exaggerated, so that it assumes the dominant part, and in this way four chief forms occur: (1) Motor, (2) maniacal, (3) paralytic, (4) cardiac, to which two other forms may be added, (5) latent, and (6) residual.

In the *motor* form the movements take the chief place;

* Based on a lecture delivered at the Hospital for Sick Children, Great Ormond Street.