left-sided scrotal abscess and a good deal of oedema of both scrotum and penis developed. These were incised under local anaesthesia. The patient left the nursing home on June 16th, and I have since heard that his progress is satisfactory.

The signs of intraperitoneal laceration of the bladder are in nearly all cases immediately manifested, but here, as in rare instances, the patient passed a few hours after the accident in comparative freedom from urgent symptoms. The explanation of this, of course, lies in the fact that he was probably too anaesthetically intoxicated to be conscious of his condition.

In the treatment of intraperitoneal rupture, the adequate drainage of the peritoneum is obviously of prime importance, but the orthodox method of combining this with the second stage of the operation—namely, the discovery and accurate suture of the laceration—would have occupied time, the expenditure of which, in such a subject, would have been only to court disaster. Even so, during the shorter operation, the continuous administration of oxygen throughout the latter period of the anaesthesia, became imperative.

Yet again it may be urged that at the second operation the suprapubic wound might have been completely closed and a catheter tied in, or catheterization every six hours resorted to. It was found, however, after the preliminary operation, that the patient was particularly intolerant of the presence of the instrument. There was, in addition, some prostatic enlargement, and the bladder itself, friable with fatty infiltration of its walls, demanded very free drainage for the conservation of the intraperitoneal closure. With due regard to the varied complications, I am still inclined to the view that, on the whole, the plan of procedure ultimately adopted was the best.

A SIMPLE METHOD FOR DETERMINING THE AMOUNT OF GLUCOSE IN DIABETIC URINE AND OTHER LIQUIDS.

BY

A. F. DIMMOCK, M.D.,

HARROGATE.

THERE are many methods now in use for estimating the sugar in urine, and many others have been suggested from time to time. The ordinary method by Fehling's solution seems one of the best, but requires apparatus and needs a good deal of experience and skill to arrive at accurate results. The fermentation test is extremely accurate, but it takes a long time to obtain results. An objection to the picric acid test recommended by the late Sir G. Johnson is the fact that creatinin produces a colouring similar to the glucose. The following simple method can easily be used without much apparatus, and gives a quick result:

The urine is diluted twenty times, that is, 10 c.cm. of urine is measured and poured into a 200 c.cm. flask; this is made up to the 200 c.cm. mark, and the whole well shaken. A solution of potassium carbonate (K_2CO_3) 2 oz. to 6 oz. of distilled water, is filtered and made up to 8 oz. To 20 c.cm. of the diluted urine 10 c.cm. of the potassium carbonate solution is added in a small flask, and this is boiled carefully for three minutes, and when cool made up to a definite amount, say 50 or 100 c.cm., with distilled water. In order to estimate the amount of sugar present, a solution of pure glucose is prepared, 1 gram in 200 c.cm. of distilled water; 20 c.cm. of this and 10 c.cm. of the potassium carbonate solution are boiled together in a small flask for three minutes, and when cool made up to 50 c.cm. or 100 c.cm. The two solutions are then compared by holding the glass tubes over a piece of white paper at an angle of 45 degrees. By pouring the liquid from the known solution into a measure glass until the tints of both are alike, and observing the amount of the known glucose solution used, the percentage can be readily determined; for example, if 27 c.cm. of the pure glucose solution were required for the solution, then, multiplying by two, we obtain 5.4 as the percentage of glucose in the urine.

I am quite aware that it is a little troublesome to prepare two solutions, and attempts will be made to use some other method of comparison. Coloured glass seems to offer the best solution of the difficulty. Experiments have been made with various iron solutions (ferric acetate and ferric and ammonium citrate), but none have come up to expectations.

Allen says in his *Chemistry of Urine* that, using normal liquor potassae B.P., when equal parts of this and urine

are boiled in a flask or test tube, the test does not indicate with certainty the presence of a small amount—say less than 2 per cent.—of glucose. Albumin should be removed by acetic acid and heat. The colouring matters in urine do not seem to have any effect, but these could easily be removed by lead acetate or subacetate, taking care not to use excess, filtering or allowing to stand; any slight excess of lead is easily soluble in the potassium carbonate solution. The colouring matter of rhubarb and senna becomes reddish-brown with alkali before heating, and samples containing catechol (pyrocatechin) acquire a brown colour on exposure to air.

Memoranda: MEDICAL, SURGICAL, OBSTETRICAL.

A MODE OF RETRO-CATHETERISM.

SOME months ago a stockbroker's clerk came to me suffering from retention of urine. All the usual plans failed to relieve him; even that useful expedient of passing an endoscopic tube to the stricture, so as to widen out its face and thus find the aperture. I then used the following method, which I had thought out years ago:

The apparatus required consisted of a long, narrow, fully-curved trocar and cannula, scarcely 6 mm. in thickness, with a round and smooth proximal end, and a filiform 18 in. long, to pass through the cannula. After aspirating the bladder suprapubically by the trocar and cannula and filling the organ with boric acid solution, I reversed the trocar, so that its round blunt end protruded from the distal end of the cannula. This end was then depressed on the trigone: then lowering the hand whilst pushing gently, the instrument slid easily into and beyond the internal meatus. The trocar was withdrawn for the second time and the filiform substituted; aided by pulling the penis forward and upward the filiform passed out through the external meatus. Had it failed to do so, at worst it would have reached the posterior surface of the stricture, and any attempt to deal with the membranous urethra from the perineum would, on account of its presence, be more exact.

After twenty-four hours urethrotomy became easy; the urine in the interval trickled freely alongside the filiform as it always does. Thus a cure was established on cheap terms. I might have used and was prepared to use a larger trocar and cannula, but that would have entailed more risk.

I think the perineal mutilation one often sees is avoidable. If a Wheelhouse operation does not immediately succeed, it is better to open the membranous urethra behind the stricture or to do retro-catheterism; even better perhaps would it be to push a very fine trocar and cannula through the stricture, the direction being aided by the finger tip, a filiform being then passed through the cannula. In any case the puncture would be harmless.

The great secret in a Wheelhouse operation is not to obliterate the little pouch left at the front of the stricture until the orifice is found; if obliterated the chief landmark is gone.

London, E.C.

JAMES MACMUNN.

OCCCLUSION OF COLON IN A NEWBORN CHILD.

THE following case was of interest in that it showed an occlusion not only of the upper part of the rectum but of the whole colon. I attended a confinement in a primipara aged 28, which, though slow, was otherwise normal, on April 25th, 1913. The pregnancy was also normal, and the baby, a female, looked well nourished and healthy. The next day the child began to vomit at intervals, then the abdomen began to show signs of distension. Urine was passed, but the bowels had not moved. Examination of the anus showed that the sphincter was normal, also the rectum for $2\frac{1}{2}$ in. to 3 in., then it narrowed and became completely occluded. The vomiting became more frequent; there were evidences of pain such as crying and pulling up of the legs.

Post-mortem examination showed that the ileum was distended in its lower two-thirds to within 3 in. of the ileo-caecal valve, where it narrowed considerably. The caecum including the appendix was small but otherwise healthy and patent, but about 1 in. beyond the ileo-caecal