ON DIATHESIS IN INFANCY:

A PLEA FOR ITS CLOSER STUDY.

[WITH SPECIAL PLATE.]

BY HECTOR CHARLES CAMERON, M.D.,

PHYSICIAN IN CHARGE OF THE CHILDREN'S DEPARTMENT,
GUY'S HOSPITAL.

INTRODUCTORY.

PROFESSOR EMMETT HOLT, of New York, in an interesting article published recently, entitled "Infant Mortality, Ancient and Modern," referred to the prejudicial influence which was exerted in England by the growing tendency to encourage lay workers and members of the nursing profession to busy themselves with the doctrines of infant hygiene and the practice of infant dietetics. Professor Holt's words are:

If in England the measures referred to have been much less successful than in other countries, we must remember that in that country the care and feeding of the infant is generally believed to be the special province of the nurse. The physician is frequently not consulted at all. The effect of this is seen in the undue prominence given to nurses and social visitors.

While we must all admire the energy and devotion with which all over the country educated lay men and women of great intelligence have applied themselves to attack the problems of infantile mortality, it must, I think, be conceded that there is a strong element of truth in Professor Holt's remarks. It is significant that at the present time the National Association for the Prevention of Infant Mortality and for the Welfare of Infancy is petitioning the medical schools and examining bodies that the subject of Infant Hygiene should occupy a definite place in the medical curriculum. The high infant mortality among the poor of our great cities has aroused the compassion and interest of a large section of the educated public; it has left the medical profession, as a whole, comparatively unmoved. And yet it is to the medical profession alone that the country must look for improvement in our present conditions, because medical men alone can bring to the study of the problem the special knowledge, insight, and habit of mind which can conceive and isolate the true nature of the difficulties which confront us daily in dealing either with infantile mortality in general or with any one particular infantile complaint. Especially is this true in the study of infantile dietetic disorders. The lay mind is necessarily apt to start with the assumption that all digestive disturbance proceeds from faults in the composition of the diet, from gross excess, or from a want of cleanliness. No doubt among the very poor the proportion of cases directly attributable to these causes is much higher than among the better to do, and there is a wide field for the instruction of poor mothers in the elementary rules of health and infant hygiene. Yet even among the poor there is abundant evidence that malnutrition and dyspepsia, on breast or on bottle, is not always to be attributed to gross ignorance or culpable carelessness on the part of the mother. The doctor alone is able to detect other causes for disturbance.

Dyspepsia is very frequently the expression of an underlying infection. The infant at birth has but one function highly developed—that of absorbing an amount of food which, relatively to that of the adult, is enormous. A function so highly developed is very readily disturbed, and infections of all sorts and in the most various situations present themselves in the guise of a primary digestive disturbance. Still more interesting is the large group of cases in which dyspepsia is dependent on peculiarities of infantile diathesis, and upon inherited, inborn abnormality of digestion. Among the infants of the well-to-do classes, where the composition and amounts of the diet are well controlled and subject to the most careful scrutiny, and where the hygienic conditions are faultless, digestive disturbances are rarely *ex alimentatione*—from gross assaults upon the digestive powers by an absolutely irrational diet—but arise commonly *ex infectione* or *ex constitutione*.

The suggestion that it is possible to recognize types of individuals whose aspect and whose liability to suffer from a certain train of symptoms marks them off as belonging to a certain diathesis, is not new. Indeed, the

conception of diathesis is almost as old as the study of medicine itself. For long it held a foremost place in all medical literature. Later, in the second half of the last century especially, there came a time when it fell into disrepute. These years were marked by great advances in the study of the local disorders of the different parts of the body as the use of the stethoscope grew general, and the methods of physical examination became highly developed. Bright, Addison, Wilks, Virchow, and many others taught the importance of morbid anatomy, and the necessity of interpreting all clinical phenomena in the light of the examination of viscera *post mortem*. Later still the science of bacteriology sprang into being, and profoundly influenced our whole conception of the processes of disease. With more definite knowledge of these processes much of the old-time teaching upon diathesis was found to be no longer tenable, and the very name to a great extent fell into disuse. To-day it again stands in the forefront of interest, both in medicine as a whole, and especially in the study of the disorders of infancy. Although most of the types of diathesis formerly described have been discarded, it is still possible to recognize certain inborn and inherited constitutional anomalies marked by the constant association in the same individual of a wide variety of symptoms and local manifestations. The presence of one set of symptoms often enables us to predict the appearance of others at a later time.

Although we have no sufficient knowledge of the true nature of the constitutional anomaly, we believe that this anomaly accounts for the presence of the local disorder, and that treatment to be successful must be directed not only against any one local manifestation, but must combat the underlying cause—the diathesis itself.

Evidence for the Existence of Certain Diatheses in Infancy.

The evidence of these wide divergences of constitution is very marked in the study of the dietetic disorders of infants. It is universal experience that upon the same diet and under the same hygienic conditions we may meet with one child who thrives and develops perfectly, whilst another becomes rachitic, a third grows anaemic, and a fourth shows a lowering of tolerance to all infective catarrhal processes. So frequently are such variations encountered that we are driven to the conclusion that infants are born with considerable differences in the chemical structure of their several organs and tissues—differences which exert their influence in greater or less degree, according to the character of the diet.

Formerly when a breast-fed infant suffered from severe and persistent dyspepsia, the failure was commonly regarded as evidence of an idiosyncrasy on the part of the mother. It was thought that her milk was of unsuitable composition, or capable in some way of exercising directly a prejudicial effect upon a normal infant. To-day we know that the milk of every woman, not herself the subject of an infective disorder, is equally suitable for the normal infant, and that the composition as well as the amount of the milk is controlled to suit his needs by the varying stimulus of suction. If the milk of one woman causes digestive disturbances, the change to the milk of another is seldom of any avail. The same woman may suckle three or four infants, only one of whom is dyspeptic and fails to thrive. Although one infant may do very badly on the breast, the same breast may yet be safely used for other normal children. Of the three great groups into which infantile dyspepsia is divided, dyspepsia *ex alimentatione*, dyspepsia *ex infectione*, and dyspepsia *ex constitutione*, the last is not only of the greatest interest, it is perhaps also of the greatest frequency. No classification of infantile dyspepsia can be complete which ignores those fundamental distinctions between individual infants. That the same diet may produce a healthy infant in one case, and in another lead steadily to severe illness and even death, forces us to attempt the study of infantile diathesis. It is right that we should have clearly in mind the results of under-feeding (*Inanition*), of over-feeding, and a temporary loss of tolerance for all or some of the constituents of the food (*Dyspepsia*), of a similar but permanent loss of tolerance (*Marasmus*), and of the symptoms of the absorption into the blood stream of certain disintegration products of abnormal

digestion (*Intoxication*). At the same time we should endeavour to appreciate the usual result of imparting to the diet a one-sided character, so that either fat or carbohydrate is present in gross excess, and we should be able to distinguish between a primary digestive disturbance and a disturbance secondary to infection, whether of the bowel wall (*enteral*), or in other parts of the body (*parenteral*).

But we must also be prepared to recognize as a sufficient cause for disturbance the existence of certain constitutional abnormalities, the "exudative diathesis" of Czerny, for example, or extreme neuropathy. The study of the disorders of infancy must concede due prominence to the consideration of the child. The subject is not exhausted by an examination of a variety of diets, by enumerating rules for dilution and dosage, and by insisting on the necessity for cleanliness. If infants could always be fed by rule of thumb, if the whole matter could be dealt with in a few pages of general directions, infant feeding might be left as at present almost outside the medical curriculum, and be regarded as of greater interest to nurses than to medical men.

As knowledge increases it becomes more and more clearly recognized that dyspepsia in infancy is not always, or even usually, accounted for by gross excess in the amount of the diet or gross faults in its composition. It becomes more and more apparent that it is wrong to approach the study of the disorders of infantile digestion in the belief that the child is the constant and unvarying factor, and in the hope that we shall one day find a universal artificial diet applicable under all conditions and to all infants.

In spite of the confident statements of those who claim unvarying success with some particular method of feeding—whole milk, for example, or the so-called "humanized" milk—the digestive powers of individual infants are very various. We shall often find it our lot to study the abnormal reactions of particular children to food which may be said to be of perfectly suitable composition, and upon which in our own experience large numbers of infants of similar age have shown uniform and steady progress.

In the study of infantile disorders it is helpful to group together in our minds certain symptoms, certain peculiarities of aspect, certain tendencies and liabilities to particular disturbances, which experience teaches us are apt to occur in infants of a certain type and of a certain inheritance. The frequency with which these groups of symptoms occur in the same individual, either simultaneously or successively, at different times in the course of growth and development encourages the belief that they are only various expressions of an underlying constitutional idiosyncrasy.

The Exudative Diathesis of Czerny.

We owe to Czerny a clear statement of the clinical symptoms of a particular diathesis which is so common that its manifestations in greater or less degree are met with every day in practice and are familiar to all.

In the older medicine a group of symptoms, constantly recurring and of great frequency, was described under the name of "scrofula." Children who showed the scrofulous diathesis were characterized by a certain physical aspect—harsh, dry skin, with a pronounced tendency to certain scaly and lichenoid eruptions, fine silky hair, which grew irregularly beyond the true hairy scalp on to the forehead and over the zygomatic arch, long eyelashes, and a tendency to a further hairy growth on the back of the neck and between the scapulae; chronic catarrhs in various situations were almost constant—nasal catarrh, with a profuse secretion which irritated and infiltrated the upper lip, nasopharyngeal catarrh, bronchial catarrh, phlyctenular conjunctivitis, leucorrhoea, and mucous diarrhoea. As a result of the chronic irritation of the mucous membranes, a secondary glandular enlargement was the rule. Tonsillar overgrowth, adenoid vegetations, hypertrophy of the lymphatic glands and of the lymphoid tissue generally of the whole body were commonly found.

With the discovery of the true nature of the tuberculous infection it became recognized that a large number of scrofulous children were infected with tubercle. In many the enlarged glands became definitely caseous; in others the catarrhal and inflammatory symptoms were

combined with diseases of bone clearly of a tuberculous nature or with tuberculides of the skin.

Nevertheless, after the separation of those individuals in whom a tuberculous infection had declared itself, there are still left a very large number of children who would unhesitatingly have been described in former times as of a scrofulous or strumous diathesis. Those children who show the catarrhal and lymphatic changes of the old-time scrofula without evidence of tubercle, have been studied under many names. Virchow borrowed from Thomas White* (1784) the name "inflammatory diathesis"; Heubner, Escherich and others speak of "lymphatismus" and the "lymphatic constitution"; French authors write of "arthritismus" and "neuro-arthritismus," the English authors of "mucous disease" and "status lymphaticus." Czerny, who has traced the diathesis backwards to earliest infancy, where it forms one of the commonest causes of digestive disturbance, impressed by the frequency with which exudative processes occur upon the skin and mucous membrane, has given it the name of the "exudative diathesis." He has emphasized especially that the increased vulnerability of skin and mucous membranes is primary, the lymphatic enlargement secondary.

Symptoms of the Exudative Diathesis in Later Childhood.

The manifestations of the diathesis in infancy are those with which we are immediately concerned. The developments later in childhood and in adult life, though not dissimilar in nature, are more varied in type and must first be described. The subjects of the diathesis are marked by a characteristic appearance (see plate). The child's face is usually plump and rounded and marked by a high colour on the cheeks and on the front of the chin. In these situations a dry, scaly eczema is usual. Seborrhoea of the scalp is common. The hair is often sparse, but growing in wavy curls rather than straight. The frequent catarrhs of nose and pharynx are accompanied by tonsillar and pharyngeal overgrowth and by enlargement of the cervical glands. Otitis media is common and the children are often in consequence deaf. The so-called "geographical tongue" is frequently seen. Bronchial catarrhs and asthma-like attacks are frequent. In some cases the symptoms in childhood persist and a permanent asthmatic tendency may be established. In such cases a history of asthma in the parents or among relatives is frequently obtained. Many asthmatics will be found to have suffered in infancy and childhood from seborrhoea of face and scalp.

In some cases the intestinal mucous membrane shares in the general tendency to catarrhal and exudative inflammation, with the production of chronic enteritis and secondary overgrowth of the intestinal lymphatic tissue. This condition is well described by English authors (Eustace Smith, Gee, etc.) under the name of "mucous disease" or "coeliac disease." In general, restlessness, irritability, and nervousness are marked symptoms, and a violent and passionate fit of crying will often make the examination of a young child difficult.

A further group of symptoms has a close association with the phenomena of anaphylaxis, urticaria, sometimes giant urticaria, transitory oedema or exudation into joints, as well as spasmodic asthma. In other cases difficulties in the digestion and metabolism of fat stand in the foreground. The "cyclical" vomiting of childhood is frequently accompanied by evidence which marks the child out as of this particular diathesis.

Perhaps the most characteristic of all the symptoms is seen in the profound constitutional disturbance which is apt to accompany those apparently slight catarrhs to which, at longer or shorter intervals, the child is usually liable. In some children hardly a week passes during which they do not sicken of an acute feverish attack. At nightfall the temperature may be 104° or 105° F., and the child may present all the appearance of serious disease; and yet the morning finds every symptom gone. If, on the other hand, the infection is more sustained, weeks of high fever may elapse, yet repeated and careful physical examination does not disclose any physical signs of disease. In the severer degrees of the condition numerous

* *A Treatise on Scrofula, commonly called the King's Evil.* By Thomas White, Surgeon to the London Dispensary; 1784.

infections are fatal which would have caused but slight disturbance in a normal child. Pneumonia has a higher mortality; a trivial laryngitis may lead directly to death; in an acute attack of diarrhoea drowsiness may gradually deepen into coma, until death ensues; sometimes sudden heart failure, from fear or other emotion, the subcutaneous injection of antidipltheritic serum, anaesthesia, or other trivial causes, may seem to be the direct cause of death. Acute infections may cause death with great rapidity, and very early in the course of the disease. A few weeks ago I performed an autopsy on a girl 17 years of age who had fallen down dead in one of the streets of the City. She had not complained of feeling ill. Marked hyperplasia of the thymus and thyroid glands and of all the lymphatic glands of the body was at once apparent. Naked-eye examination discovered no other cause for death, yet careful histological examination by Dr. Johnson showed a very early miliary tuberculosis of the lung. Lastly—as the French authors especially have insisted—arthritis, chorea, endocarditis, and other rheumatic manifestations are most commonly and most typically seen in children who exhibit marked signs of the diathesis.

The symptoms so described embrace a very large number of disorders, not usually regarded as akin. The hypothesis that all are dependent upon an underlying diathesis is founded only upon the observation that groups of such symptoms tend to occur in the same individual. As a rule, in any one individual a particular group of symptoms recurs again and again. In other cases the type of the manifestations changes at different stages in the development of the child. In one child the tendency to pyrexial attacks, nasal or naso-pharyngeal catarrhs, otitis media, tonsillar and adenoid hypertrophy, and enlargement of the cervical glands is most marked. In another digestive disturbances and the antipathy to a diet rich in fat is prominent. In a third urticaria and bronchial catarrh, or true spasmodic asthma, predominates; in a fourth seborrhoea or eczema of skin and scalp. The diathesis is hereditary. Often it occurs in all members of a family, and it will then usually be found that one or other parent has suffered severely in youth in the same way.

The therapeutic measures by which we may hope to control the manifestations of the diathesis are both local and general. In the latter group are a number of well-ascertained dietetic and hygienic regulations.

For my present purpose the chief interest lies in the extension by Czerny and others of the recognition of the diathesis back to the earliest days of life, and the emphasis more recently laid on the part which it plays in the production of digestive disturbances both on breast and on bottle.

Symptoms of the Exudative Diathesis in Infancy.

In early infancy the most common way in which the inheritance shows itself is the complete or relative failure of breast-feeding. The children often show none of the strength, vigour, and firmness of the normal breast-fed baby. From the first they may be meagre, small, and complaining, and persistently dyspeptic. The mother's breasts may be well formed and full of milk, and yet for weeks and even months no improvement takes place. Dyspepsia, with the passage of green stools and vomiting is frequent, and in a breast-fed child should always suggest the existence of this diathesis. In other cases, however, a different picture presents itself. The infant achieves a rapid rise in weight, yet the more rapid the growth the more marked do the other symptoms of the exudative diathesis become. Finkelstein especially has emphasized the tendency for infants who show the diathesis to belong to one or other of these two types—the thin and wasted type, or the fat, pasty and eczematous type.

Cutaneous Symptoms.—Seborrhoea of the scalp is extremely common in the form of a dirty greyish-brown fatty secretion appearing irregularly upon the hairy scalp, which resists all the mother's attempts to remove it. This symptom appears in breast-fed infants suffering from dyspepsia with very great frequency.

A dry eczema of the cheeks and chin is common, especially in thin infants. The skin in these situations is reddened, dry, scaly and rough. In the fat and pasty type an acute irritating eczema is more frequent.

Urticaria papulata is often seen. The minute papules are scattered on the scalp, body, and extremities, but not on the face. They are often present on the soles of the feet. Each crop tends to last two or three days.

Intertrigo occurs readily, in spite of the greatest cleanliness, in the groin and axilla, in the genital folds, or behind the ear.

Symptoms in the Mucous Membranes.—The same vulnerability which appeared in the skin shows itself in the mucous membranes. Naso-pharyngeal catarrh, with accumulation of secretion in the posterior nares and pharyngeal wall, and slight enlargement of the cervical glands, is common. Because the infant lies constantly on his back the anterior nares often remain free from secretion, so that the appearance does not suggest nasal catarrh. The interference with nasal breathing is so great that suckling is rendered difficult. Often this obstruction is the direct cause of weaning, a course to which the mother is inclined because of the coincident dyspepsia. The catarrh often sets in with high fever, loss of appetite, diarrhoea and vomiting, and prostration; a diagnosis of bronchopneumonia is often wrongly made. Examination of the naso-pharynx and the cervical glands makes the diagnosis clear. Laryngitis, bronchitis, and otitis media are frequent complications. The lingua geographica, conjunctivitis, eczematous or phlyctenular, and circular caries of the teeth are seen in later infancy. The mucous membrane of the intestine is also affected. In earliest infancy constipation or dyspeptic green stools are common symptoms. Later follicular colitis, mucous diarrhoea, and threadworms are frequent.

Symptoms of Overgrowth of Lymphoid Tissue.—The spleen, thymus, tonsils, lymphatic glands, and intestinal follicles all show a tendency to an enlargement, which is probably secondary to the chronic irritation in the areas which they drain.

Treatment.—The symptoms of the diathesis are encouraged or controlled largely by the composition of the diet. Especially marked, as a rule, is the ill effect of a fat rich diet, such as cow's milk. Even the breast milk may disagree for this reason. Limitation of the amount of milk and the early substitution in part of carbohydrate food is usually the most successful line of treatment. Without dietetic regulations the local treatment of the various manifestations, the repeated catarrhs, or the lymphoid overgrowth—as, for example, the adenoid enlargement—is often without avail. Over-feeding is especially harmful, and the food should be reduced so that it just covers the physiological needs. As might be expected, the subjects of the diathesis are apt to show a very violent reaction to all infections in which catarrhs are common. Measles and whooping cough may endanger life.

Illustrative Cases.—The diathesis so described is of such common occurrence that numerous cases might be quoted from among the patients seen upon a single day in the out-patient department of a large hospital. The following case illustrates the association of infantile dyspepsia, seborrhoea, adenoid overgrowth, and asthma.

CASE I.

An infant, 3 months old, suffering from severe seborrhoea of the scalp, was brought to me on December 3rd, 1911. Although nursed on the breast he had suffered severely from dyspepsia almost from the first. At 5 weeks old he had been weaned. He was always sick when cow's milk even in considerable dilution was given. He had had much less dyspepsia upon sweetened condensed milk. The seborrhoea had appeared at 3 weeks. The scalp when seen was completely covered with masses of secretion.

A fat-low carbohydrate-rich diet was ordered, and recovery followed in about six weeks' time. On April 20th, 1914, he was again brought to me. I was told that he was well except for a tendency to suffer from slight scaly eczema of the cheeks and chin, repeated nasal catarrhs, and severe spasmodic attacks of dyspnoea. Two maternal aunts suffered from asthma. He had some tonsillar enlargement and a mass of adenoid vegetations.

Dietetic rules were given. He had been taking large amounts of cow's milk, and a diet predominatingly vegetable and cereal was recommended. The adenoids may have to be removed later. The association between them and the asthma cannot, however, be regarded as cause and effect.

CASE II.

A girl aged 8 was brought to the hospital for growing pains. She had been weaned for failure to thrive at three months, although the mother had plenty of milk. She walked and cut her teeth somewhat late. In her third, fifth, and sixth year

she was admitted to hospital for bronchopneumonia, but recovery was always rapid, and pyrexia short-lived. The mother thought that the attacks did not differ from many others which the child had had, and which she had almost grown to neglect. They set in suddenly with burning heat, flush, and dyspnoea. At such times the glands in the neck always enlarged. (In another case with similar history the mother told me that a diagnosis of mumps had been made in the hospital surgery on four separate occasions, and on each she had attempted to obey instructions and carry out isolation of the child.) At 2 years old she had had eczema of the scalp. She had then been fat, but had got thin later. On examination she was seen to be a small child, with sparse, wavy, curly, reddish hair, and dark eyelashes. Fine downy hair grew low down on the forehead, beyond the true hairy scalp and over the cheeks almost like a whisker. There was marked enlargement of tonsils and adenoid overgrowth. The nostrils were pinched and atrophied. The cervical glands were large and palpable, the axillary and inguinal glands smaller, but easily palpable and visible under the thin skin. The face was reddened with rough, dry skin on the cheeks. The tongue showed the condition *lingua geographica*. The chest was narrow, with little subcutaneous fat. The skin was very white over the chest, and changed to a deeply pigmented colour over the distended abdomen. She complained of abdominal pain at intervals, often had diarrhoea and blood-stained stools. There had been frequent prolapse of the rectum. At one time threadworms had been seen. Lately she had complained much of joint pains, and a diagnosis of acute rheumatism had been made.

CASE III.

The portrait of C. E., aged 5, is selected to show the sparse hair with indeterminate anterior margin, dry eczema of cheeks and scalp, adenoid overgrowth, and enlarged pigmented abdomen. She was brought to the hospital for repeated attacks of diarrhoea and abdominal pain. She had been weaned at 3 months, and had never thrived on the breast, although the attempt had been persisted in. The cervical, inguinal, and axillary glands were palpable. She was subject to repeated pyrexial attacks of short duration, during which she vomited, became drowsy and listless, and felt "burning hot."

CASE IV.

A male infant, aged 3 months, admitted to Guy's Hospital under my care. In the surgery the temperature was 105°, and there was marked dyspnoea. Examination showed that the lungs were free from physical signs. The cervical glands were enlarged, and the fauces and pharyngeal wall were reddened and inflamed. The temperature remained high for two days, when recovery took place rapidly. The child had suffered continuously since birth from dyspepsia, and the weight—7 lb.—was less than at birth. Weaning had taken place at 6 weeks, and the diet had since consisted of equal parts of cow's milk and water, with added cane sugar. There was a slight roughness of the skin of the cheeks and forehead. Upon a diet of citrated whole milk the weight rose rapidly, but as the child grew fat the eczema became acute, and spread actively over the cheeks. After four weeks the diet was changed to sweetened condensed milk, when the eczema immediately improved, and the face became quite clear. On three occasions there have been acute paroxysmal attacks of dyspnoea, without pyrexia, which suggest the presence of asthma.

Such cases might be multiplied indefinitely. Czerny has called the disorder the commonest ailment of childhood. I am well aware of the danger of laying too great stress upon such hypothetical conceptions as that of a diathesis the true nature of which is still a subject for speculation, or of allowing explanation by assumed causes to take the place of inquiry into anatomical and clinical facts. I believe, however, that in ignoring this aspect of disease altogether, and in concentrating all attention upon the local phenomena, we commit a serious error. In the large new textbook of children's disease, edited by Batton, Garrod, and Thursfield, I find little mention of the question of diathesis at all, although perhaps no subject has been dealt with so minutely in the recent literature of diseases in children or debated so earnestly at meetings of societies and congresses on the Continent. Speaking of the spasmophilic diathesis, so good an observer as Dr. Leonard Guthrie dismisses the conception somewhat contemptuously as savouring of an extinct medicine.

Neuropathy in Infancy.

A second type of infantile diathesis which is worthy of close study, and which is also the direct cause of much infantile dyspepsia, both in breast-feeding and in artificial feeding, is found in severe neuropathy in infancy. The neuropathic inheritance may show itself from the earliest days of life in an altogether abnormal restlessness and sensitiveness to external stimuli. The neurotic infant is usually meagre and spare. The limbs often show a considerable degree of hypertonus. The abdomen tends to be strongly contracted, so

that the recti muscles stand out prominently. Such children sleep badly from the first. When the healthy child sleeps he can be moved from his cot without awakening, and ordinary conversation in the room in which he lies has no power to disturb him. The neuropathic infant sleeps only to wake at the slightest sound. If any one approaches his cot, with a start he is wide awake again. No doubt, when mother as well as child are of neurotic temperament, the condition of the child is aggravated by the atmosphere of his surroundings. His will and his temper suffer no opposition. If food, for example, is given to him on a single occasion to quiet his midnight cry, there is no rest for his mother for many subsequent nights. More than other children, the neuropathic child, in the natural desire of mother and nurse to quiet him, suffers the risk of overfeeding, just as more than other children he tends to receive excessive psychological stimulation. Of all infants, he is in most need of rest; yet, in the hope of coaxing him to quietness, mother and nurse exhaust themselves and him in devising new psychical stimuli. The constant appeal to his senses produces a precocious mental development, but it aggravates the nervous instability and exhaustion. Finally, he can hardly be left alone for a moment, so accustomed has he become to the efforts of those who minister to his sensations. A little tyrant in the house, he may only rest on his daily excursions abroad, when at last he may lie still looking at the clouds, the traffic, or the trees.

Among such infants with a strong neuropathic inheritance dyspepsia is very frequent, even apart from the likelihood that their fretfulness and clamour will lead to overfeeding. In a few cases vomiting may be so marked that all food for a time seems to act as a veritable emetic. In other cases oesophageal or pyloric spasm develops, and it is almost exclusively in infants of this type that we find the development of hypertrophic pyloric stenosis.

CONCLUSION.

Many other examples of congenital inborn abnormality of constitution might be quoted, but enough has been said to make clear the contention that the problems of infantile dietetic disorders require for their solution all the trained faculties of medical men, and that a close supervision of the amount, composition, and quality of the diet will not avail in a great number of cases to detect the cause or ensure recovery. In infantile disorders, as elsewhere in curative and preventive medicine, the help given by trained, competent, and intelligent nurses and voluntary workers can only be subsidiary to the work of experienced medical men. At the present time, because of the relatively small number of medical men who are interested in the subject or who study it in a rational and broad-minded way, there is some tendency for their place to be usurped by lay workers, who are perhaps more interested but whose lack of training renders them unable to appreciate the complex nature of the problems concerned. The scientific study of the ailments of very young infants is of supreme importance to the State. Progress in this country of the control of infant mortality is hindered not by the apathy of the public or the State, but by the comparative neglect of the profession. To that extent the criticism of Professor Holt is true.

AN International Congress on School Hygiene will be held in Brussels in 1915. The following subjects are proposed for discussion: The school building and its equipment; medical inspection in urban and rural schools; the prevention of contagious diseases in school; the teaching of hygiene to teachers, scholars, and parents; school hygiene in its relation to physical education at the different ages of the children; teaching methods, syllabuses, and school equipment in relation to hygiene; school hygiene in its relation to exceptional children; school hygiene with regard to adolescents. The Congress will be held under the patronage of H.M. Albert, King of the Belgians, and under the auspices of the Belgian National Institute of Paedology and of the Belgian Paedo-technical Society. M. J. Corman, Director-General of Primary Studies at the Department of Science and Art, and Dr. J. Demoor, Rector of the University of Brussels, are Presidents of the Organizing Committee. All communications and inquiries should be addressed to Dr. H. Rulot, secretary-general, 66, rue des Rentiers, Brussels.

FIG. 2.

FIG. 1.

DR. H. CHARLES CAMERON : DIATHESIS IN INFANCY.