Jackson's pericolic membrane, on the other hand, may give rise to symptoms simulating chronic appendicitis. This occasional membrane is a congenital abnormality, and takes origin from the primitive right lateral process of the great omentum. When only a limited band persists as a relic of the membrane, constriction of the ascending colon may result, though often the symptoms resembling chronic appendicitis are due to an associated abnormal mobility of the proximal colon.

In the condition of mobile proximal colon, which is a reversion to the primitive pronograde arrangement, the patient is predisposed from birth to constipation and the various ills resulting from it. Symptoms resembling chronic appendicitis may result, or actual inflammation of the appendix, but in either case the right surgical procedure is not a simple appendicectomy, but fixation of the colon back into the right loin.

The commonest cause of an erroneous diagnosis of chronic appendicitis is right tubo-ovarian disease, often of gonorrhoeal or septic origin. In such cases a spasmodic contraction of the right psoas muscle sometimes occurs on palpation, which may be mistaken for an abscess or a neoplasm.

Many cases of chronic pain in the right iliac fossa cannot, with our present knowledge, be diagnosed with precision before operation. All that can be said is that they need exploration. In exploring, the incision should be adequate for an inspection of all the possible sources of the mischief, and such source when found should be dealt with by appropriate measures. The practice of a blind appendicectomy in such cases through a 2 in. gridiron incision is unscientific, and often disastrous to the patient's prospects of a cure.

AN UNUSUAL CASE OF APPENDICITIS.

BY

E. W. M. HUBERT PHILLIPS, M.A., M.B., B.S. OXON.,
SURGEON TO MISS TALBOT'S HOSPITAL, MARGAM, GLAMORGANSHIRE.

A COLLIER was suddenly seized with abdominal pain of such severity as to necessitate the administration of chloroform by his medical attendant, Dr. Cooper of Kenfig Hill, Glamorgan. On the following day I was sent for in consultation and advised removal to Miss Talbot's Hospital at Port Talbot. The patient complained of sharp continuous pain situated under the heart and in the upper part of the abdomen, of obstinate constipation, and of vomiting of clear green fluid. In 1887 he had a similar pain which lasted five days; it came on after a fall from a tree. On five occasions since then he had suffered from the same pain, of greater or less severity, the attacks lasting five or six days.

His face was pale and typically Hippocratic and covered with beads of perspiration. The tongue was clean, the pulse was 78, and the temperature 96.8° F. The abdomen was not distended. The respiratory movements were good, but better on the right side than on the left. Superficial and deep epigastric tenderness were present, and there was a zone of hyperaesthesia in the left hypochondrium. There was deep tenderness on pressure over the left kidney region posteriorly. The whole of the left side was tympanitic. The liver dullness was normal. The apex of the left lung showed some apical retraction, Krönig's sign being present, and there was some impairment of breath sounds. There was a slight systolic murmur at the apex of the heart, conducted out to the axilla. The urine was normal. There was slight leucocytosis. Loewi's adrenalin mydriasis test was applied with a view to determine the possible presence of pancreatitis. The result was negative. There was nothing definitely indicative of perforation of a peptic ulcer. Repeated enemata were unavailing.

The patient was seen by my partner, Dr. J. H. Davies, and it was decided to wait twelve hours before proceeding to laparotomy in view of the patient's history. The patient appeared to improve slightly, but in about eight hours he suddenly became much worse; the temperature fell to 96°; the pulse went up to 102, and became very thin. He was now evidently moribund. Operation was out of the question, and in a few hours the patient died, evidently of shock.

Post-mortem Examination.

The caecum and ascending colon were enormously distended and hyperaemic. This part of the gut was freely movable, and had a large mesentery. It lay looped between the hepatic and splenic flexures, below the transverse colon and great omentum, and suspended on the left to the cardiac end of a dilated stomach, by the tip, middle, and mesentery of a long recurved appendix.

The appendix, when unravelled, was 9½ in. long, was full of hard concretions, and was gangrenous at its tip. The tip was densely adherent to the greater curvature of the stomach on the left side, and by fine adhesions to the spleen. The right iliac fossa was occupied by coils of small intestine.

There was a small calcareous old focus of phthisis at the apex of the left lung. There was some thickening of the mitral valve.

A CASE OF RUPTURE OF THE KIDNEY FROM SLIGHT TRAUMA.

By J. LEE ATKINSON, M.R.C.S., L.R.C.P.,
VICTORIA, AUSTRALIA.

RUPTURE of the kidney is a comparatively rare result of trauma, and more particularly so when the actual kidney tissue itself is lacerated. Keen quotes Küster's statistics of injuries under his observation in the clinics of Berlin and Basle, and these show only 10 cases in 7,741 injuries, or a little over 1 in 1,000.¹ Israel's, with a large experience of renal surgery, had only 1 operative case.²

But in practically all cases met with either in actual practice or in the records, the condition has always been caused by a severe accident, such as falling from a height, crushing between buffers of a train, or by the wheel of a heavy cart passing over the abdomen. In view of these facts the following case is of interest, as the injury was caused by an accident of the most trivial nature, and the subsequent history also contains points of interest.

R. A., a girl of 14, above the average height and well made in proportion, had never had a previous illness of any consequence. At 9 p.m. on August 28th, 1913, she ran along the verandah of the house after a dog. Between the two corner posts of the verandah some light wire netting had been fixed to a height of 2 ft. and fastened on each side to the post by a light staple. The floor of the verandah was 11 in. from the ground, and immediately beneath it at that point was a large flower bed which the day previously had been dug over, and in consequence was in a soft condition. The end of the verandah was only 15 ft. from the door, and so she could not have gained much momentum in the short distance. She ran into the netting, having forgotten it was there, and it caught her in the middle, gave way, and she fell on to the flower bed. She immediately complained of intense abdominal pain with nausea, and was carried into the house. A quarter of an hour later she wanted to micturate, and passed a quantity of bright arterial blood. Dr. Parker of Kingston was sent for and he telephoned for me.

I saw the girl at 1 a.m., and she was lying in bed in a condition of profound shock. The abdomen was rigid, and she complained of great pain, not confined to any particular region. Another marked feature was thirst. The temperature was 97° F.; the pulse 124, thin and thready. The face was pallid and the facies Hippocratica well marked. The right flank was dull on percussion, but the liver dullness was hard to make out; it was diminished in area. She had passed blood by the urethra three times since the accident—about a pint in all. Owing to the fact that the parents were away from home, operation was delayed until their return at 10 a.m. on the following day.

Operation.

A median subumbilical exploratory incision was first made, and immediately the peritoneum was opened about a pint of blood-stained serum gushed out, together with a large quantity of faecal-smelling gas. With sponges the abdominal cavity was dried as far as possible and the bladder examined. There was no sign of any injury to that organ, but it was of bluish colour, due to the blood it contained.

The ascending colon was pushed towards the middle line and was congested and dilated at the caecal end. About midway between the caecum and the splenic flexure there was a kink, formed by the lower end having a long mesocolon allowing a partial volvulus. Just above this kink, and about on the level of the junction of the lower and middle third of the kidney, was a transverse tear, about an inch long, into the lumen of the gut. On replacing the lower end of the colon in its normal position and fixing it there by shortening stitches in the mesentery, a quantity of flatus escaped from the rent in the bowel wall and the whole of the ascending colon collapsed.