he got a considerable number of children with definite physical signs in the lungs, and was accustomed to inquire very carefully into their history in order to discover whether there were night sweating, inability to eat food, constant tiredness, and lessened weight, because, if these existed, the case must be treated. It was safer, for some of them at least, to rely far more on symptoms—although this might be heterodox—than on physical signs, the full meaning of which many of them were unable to appreciate.

Dr. D. BARTY KING (London) said that some years ago, while examining a number of London school children, he thoroughly tested for true phthisis, with the result that out of 1,500 examinations only about 4 per 1,000 children revealed this condition on ordinary clinical examination. The presence of glands in the neck was sometimes put down as a sign of pulmonary tuberculosis, but was quite fallacious. It was quite a frequent occurrence to have, as he had had the previous week, a case which had been treated for six months as one of pulmonary tuberculosis, with marked dullness at both apices, and to find, on taking the case into the observation wards, that nothing of this nature was discoverable in the lung at all, and that the *x*-ray report and the cutaneous tuberculin test were negative. In one case recently brought to his notice the *x*-ray report stated that there was peribronchial mottling, and inferred definite pulmonary tuberculosis of the left lower lobe of the lung. Clinically, from the posterior aspect, there were a few crepitations. The patient died of another affection, and at the necropsy he took care to section the lung from the roots towards the periphery, finding thickened bronchi, but no trace whatever of tuberculosis. On staining for tubercle there was no trace. He would suggest that skiagraphers should report, in the first place, exactly what they saw, and on the other side, quite separately, what interpretation they placed on it.

Dr. F. J. SMITH (President of the Section of Medicine), after summing up the discussion, said he preferred to trust more to a patient's feelings than to any physical signs for his view as to taking therapeutic measures. Bronchopneumonia slew as many children as tubercle, though in a different manner.

Dr. D. B. LEES, in reply, desired to say (1) that the dull areas, though found very frequently, were not universal, even in children. In many cases, both in adults and in children, none of the dull areas characteristic of pulmonary tuberculosis could be detected; (2) the frequency with which these dull areas were found in children was roughly approximate to the results obtained by the tuberculin test. If percussion were perfectly carried out, the parallelism between clinical, experimental, and pathological results would become very much closer than it had been in the past; (3) as had been brought to their notice already, and as he himself had emphasized elsewhere, the discovery of these dull areas did not prove an active condition. One judged partly by auscultatory signs and partly by symptoms, but in all doubtful cases it was best to advise a short course of treatment by rest and antiseptic inhalation for two or three weeks. If at the end of that course the dull areas had distinctly diminished in size, there was good reason for believing that there was a certain amount of activity about them. The younger the patient in whom such dull areas were detected, the greater was the probability that they indicated active disease. An incipient pulmonary tuberculosis was often insidious in onset, and might slowly extend, with very little auscultatory evidence, and with few or no general symptoms. (4) Several speakers had expressed very definitely a belief that the two conditions of "apical phthisis" and what had been called "hilus tuberculosis" were different in nature. One saw a similar difference in cases of tuberculosis in adults. In children the difference was more marked; there was greater tendency to spread, and less tendency to break down. They were different types of the same disease, pulmonary tuberculosis. (5) Until physicians would undertake to improve their percussion they would not make progress in this matter, and this failure in accurate diagnosis might easily cause disaster in the practice of those who were treating cases by means of tuberculin.

Dr. CLIVE RIVIERE, in his reply, pointed out that what he and his colleagues in opening the discussion had endeavoured to do was to draw attention to points in diagnosis which they considered the most helpful; they were quite alive to other points. The speaker, for instance, fully recognized the distinction between tuberculous infection and tuberculous disease, and they agreed with Dr. F. J. Smith and Dr. Prest on the importance of considering symptoms, and disagreed with Dr. McNeil as to the value of cutaneous tuberculin tests, since these could only discover tuberculous infection. The fact that they were discussing chronic disease practically excluded infants, and the cutaneous tuberculin tests were not of much value after infancy was past. He agreed with Dr. Eve that cases of tuberculous chest glands hardly merited the title of "pulmonary tuberculosis," although the lung was often slightly involved. The speaker had introduced them into the discussion because they gave definite lung signs, and because these signs were commonly mistaken for those of true lung infiltration. Both *post-mortem* experience and skiagrams upheld the view of a spread of tuberculosis outwards from the glands against the lymph stream; the work of Ghon, quoted by Dr. Barty King, dealt only with primary lung infection in infants, and not with the later stage in older children.

Dr. IRONSIDE BRUCE, in reply to a question, said that as to the diaphragmatic excursion in children, the distinction between the right and left diaphragm was less considerable than in adults. The right was a little higher in children, but not in such a marked degree as in cases occurring later in life. In children it was not possible to distinguish between a serous and a purulent fluid. Dr. Langmead had stated that by *x*-ray examinations active and inactive conditions could not be distinguished, but this was not invariably true. Of some cases it was possible to say that they were active, and of others that they were inactive, although, of course, many cases were indistinguishable. He had purposely omitted any mention of striations extending from the root, as they were seen in perfectly healthy children, so he did not rely upon them; but he did rely upon opacities which were arranged in groups, for they seemed to represent catarrhal infiltration of the lung.

ON SOME CONDITIONS SIMULATING CHRONIC APPENDICITIS.*

By JOHN MORLEY, M.B., Ch.M., F.R.C.S.,

SURGEON TO ANCOATS HOSPITAL, MANCHESTER; LECTURER IN CLINICAL ANATOMY, UNIVERSITY OF MANCHESTER, ETC.

THE undoubted fact that in past years many unnecessary operations for appendicitis have been performed where the appendix was not the cause of the symptoms should challenge us to a far more critical attitude of mind. It is particularly in that variety of chronic appendicitis characterized by chronic pain and tenderness in the right iliac fossa, without any history of acute attacks, that mistakes occur, and only those other causes of similar symptoms that lie in the immediate neighbourhood of the appendix are here considered. Lane and his followers hold that symptoms simulating chronic appendicitis may be caused by a kink of the lower ileum produced by a peritoneal band occasionally present on the left or inferior aspect of the ileum some three inches from the caecum, which tethers down the gut near the pelvic brim. Lane's view of the origin of this band—that it was acquired as the result of a tendency to enteroptosis—has been quite disproved, as has the peritonitic theory, though inflammatory bands may occur in the same situation. The band is congenital in origin and a relic of the descent of the testis or ovary. There is no evidence that the band had caused any intestinal stasis in my own cases, either from radiographic findings, clinical histories, or appearances at operation, and it is argued that, apart from the very rare accident of inflammatory contraction of the band, it probably never gives rise to trouble, so that the many ingenious operations devised for dealing with it are the outcome of misdirected enthusiasm.

* Abstract of a paper read before the annual meeting of the Lancashire and Cheshire Branch of the British Medical Association.

Jackson's pericolic membrane, on the other hand, may give rise to symptoms simulating chronic appendicitis. This occasional membrane is a congenital abnormality, and takes origin from the primitive right lateral process of the great omentum. When only a limited band persists as a relic of the membrane, constriction of the ascending colon may result, though often the symptoms resembling chronic appendicitis are due to an associated abnormal mobility of the proximal colon.

In the condition of mobile proximal colon, which is a reversion to the primitive pronograde arrangement, the patient is predisposed from birth to constipation and the various ills resulting from it. Symptoms resembling chronic appendicitis may result, or actual inflammation of the appendix, but in either case the right surgical procedure is not a simple appendicectomy, but fixation of the colon back into the right loin.

The commonest cause of an erroneous diagnosis of chronic appendicitis is right tubo-ovarian disease, often of gonorrhoeal or septic origin. In such cases a spasmodic contraction of the right psoas muscle sometimes occurs on palpation, which may be mistaken for an abscess or a neoplasm.

Many cases of chronic pain in the right iliac fossa cannot, with our present knowledge, be diagnosed with precision before operation. All that can be said is that they need exploration. In exploring, the incision should be adequate for an inspection of all the possible sources of the mischief, and such source when found should be dealt with by appropriate measures. The practice of a blind appendicectomy in such cases through a 2 in. gridiron incision is unscientific, and often disastrous to the patient's prospects of a cure.

AN UNUSUAL CASE OF APPENDICITIS.

BY

E. W. M. HUBERT PHILLIPS, M.A., M.B., B.S. OXON.,
SURGEON TO MISS TALBOT'S HOSPITAL, MARGAM, GLAMORGANSHIRE.

A COLLIER was suddenly seized with abdominal pain of such severity as to necessitate the administration of chloroform by his medical attendant, Dr. Cooper of Kenfig Hill, Glamorgan. On the following day I was sent for in consultation and advised removal to Miss Talbot's Hospital at Port Talbot. The patient complained of sharp continuous pain situated under the heart and in the upper part of the abdomen, of obstinate constipation, and of vomiting of clear green fluid. In 1887 he had a similar pain which lasted five days; it came on after a fall from a tree. On five occasions since then he had suffered from the same pain, of greater or less severity, the attacks lasting five or six days.

His face was pale and typically Hippocratic and covered with beads of perspiration. The tongue was clean, the pulse was 78, and the temperature 96.8° F. The abdomen was not distended. The respiratory movements were good, but better on the right side than on the left. Superficial and deep epigastric tenderness were present, and there was a zone of hyperaesthesia in the left hypochondrium. There was deep tenderness on pressure over the left kidney region posteriorly. The whole of the left side was tympanitic. The liver dullness was normal. The apex of the left lung showed some apical retraction, Krönig's sign being present, and there was some impairment of breath sounds. There was a slight systolic murmur at the apex of the heart, conducted out to the axilla. The urine was normal. There was slight leucocytosis. Loewi's adrenalin mydriasis test was applied with a view to determine the possible presence of pancreatitis. The result was negative. There was nothing definitely indicative of perforation of a peptic ulcer. Repeated enemata were unavailing.

The patient was seen by my partner, Dr. J. H. Davies, and it was decided to wait twelve hours before proceeding to laparotomy in view of the patient's history. The patient appeared to improve slightly, but in about eight hours he suddenly became much worse; the temperature fell to 96°; the pulse went up to 102, and became very thin. He was now evidently moribund. Operation was out of the question, and in a few hours the patient died, evidently of shock.

Post-mortem Examination.

The caecum and ascending colon were enormously distended and hyperaemic. This part of the gut was freely movable, and had a large mesentery. It lay looped between the hepatic and splenic flexures, below the transverse colon and great omentum, and suspended on the left to the cardiac end of a dilated stomach, by the tip, middle, and mesentery of a long recurved appendix.

The appendix, when unravelled, was 9½ in. long, was full of hard concretions, and was gangrenous at its tip. The tip was densely adherent to the greater curvature of the stomach on the left side, and by fine adhesions to the spleen. The right iliac fossa was occupied by coils of small intestine.

There was a small calcareous old focus of phthisis at the apex of the left lung. There was some thickening of the mitral valve.

A CASE OF RUPTURE OF THE KIDNEY FROM SLIGHT TRAUMA.

By J. LEE ATKINSON, M.R.C.S., L.R.C.P.,
VICTORIA, AUSTRALIA.

RUPTURE of the kidney is a comparatively rare result of trauma, and more particularly so when the actual kidney tissue itself is lacerated. Keen quotes Küster's statistics of injuries under his observation in the clinics of Berlin and Basle, and these show only 10 cases in 7,741 injuries, or a little over 1 in 1,000.¹ Israel's, with a large experience of renal surgery, had only 1 operative case.²

But in practically all cases met with either in actual practice or in the records, the condition has always been caused by a severe accident, such as falling from a height, crushing between buffers of a train, or by the wheel of a heavy cart passing over the abdomen. In view of these facts the following case is of interest, as the injury was caused by an accident of the most trivial nature, and the subsequent history also contains points of interest.

R. A., a girl of 14, above the average height and well made in proportion, had never had a previous illness of any consequence. At 9 p.m. on August 28th, 1913, she ran along the verandah of the house after a dog. Between the two corner posts of the verandah some light wire netting had been fixed to a height of 2 ft. and fastened on each side to the post by a light staple. The floor of the verandah was 11 in. from the ground, and immediately beneath it at that point was a large flower bed which the day previously had been dug over, and in consequence was in a soft condition. The end of the verandah was only 15 ft. from the door, and so she could not have gained much momentum in the short distance. She ran into the netting, having forgotten it was there, and it caught her in the middle, gave way, and she fell on to the flower bed. She immediately complained of intense abdominal pain with nausea, and was carried into the house. A quarter of an hour later she wanted to micturate, and passed a quantity of bright arterial blood. Dr. Parker of Kingston was sent for and he telephoned for me.

I saw the girl at 1 a.m., and she was lying in bed in a condition of profound shock. The abdomen was rigid, and she complained of great pain, not confined to any particular region. Another marked feature was thirst. The temperature was 97° F.; the pulse 124, thin and thready. The face was pallid and the facies Hippocratica well marked. The right flank was dull on percussion, but the liver dullness was hard to make out; it was diminished in area. She had passed blood by the urethra three times since the accident—about a pint in all. Owing to the fact that the parents were away from home, operation was delayed until their return at 10 a.m. on the following day.

Operation.

A median subumbilical exploratory incision was first made, and immediately the peritoneum was opened about a pint of blood-stained serum gushed out, together with a large quantity of faecal-smelling gas. With sponges the abdominal cavity was dried as far as possible and the bladder examined. There was no sign of any injury to that organ, but it was of bluish colour, due to the blood it contained.

The ascending colon was pushed towards the middle line and was congested and dilated at the caecal end. About midway between the caecum and the splenic flexure there was a kink, formed by the lower end having a long mesocolon allowing a partial volvulus. Just above this kink, and about on the level of the junction of the lower and middle third of the kidney, was a transverse tear, about an inch long, into the lumen of the gut. On replacing the lower end of the colon in its normal position and fixing it there by shortening stitches in the mesentery, a quantity of flatus escaped from the rent in the bowel wall and the whole of the ascending colon collapsed.