ARTIFICIAL PNEUMOTHORAX.

BY

H. DE CARLE WOODCOCK, M.D.,

LEEDS,

AND

J. A. M. CLARK, M.D.,

LEEDS.

THE suggestions we have to offer as to the value and dangers of artificial pneumothorax are based upon experience gained in over 2,000 operations. Many were done in nursing homes, but by far the greater number in the hospitals at Armley and Killingbeck, to which we are attached. Our cases are of three types:

1. Early cases with localized unilateral disease, characterized by little pleuritic thickening and few constitutional symptoms.
2. Medium cases with more extensive disease, generally bilateral, with more evidence of pleuritic thickening, and with (a) slight or (b) marked signs of constitutional disturbance.
3. Severe bilateral cases with much constitutional disturbance, offering no hope of cure by ordinary methods of treatment.

Preparation of Patient.

The patient should spend one day in bed before the first operation; more is not needed, except for cases in Class 3. A hypodermic dose of urea hydrochloride (grains 2) and quinine sulphate is given overnight, together with $\frac{1}{100}$ grain of morphine and $\frac{1}{100}$ grain of atropine. Just before the operation the spot selected for puncture is painted with tincture of iodine, and novocain and adrenalin are injected into the site of puncture, the injection being so managed that both the superficial and the deep parts are anaesthetized. Again urea hydrochloride and quinine are administered subcutaneously, since deep injections sometimes cause violent spasm in the intercostal muscles, which should be avoided.

Site of Operation.

With us the site of election for operation is the junction of a mid-axillary line with a line running straight out from the nipple, but this point may have to be abandoned, especially if there is much basal pleuritic thickening. The thickened pleura has to be regarded as a continent shelving into an open pleural sea. We seek, therefore, a resonant spot, preferably in the axillary line, and push the needle in until the manometer shows that it is in the free pleura. Often it may be necessary to forego the local anaesthetic, for novocain must not be given beyond the dose of safety, and therefore cannot be frequently injected at the same operation.

1. Early Unilateral Cases.

The operation is easy, though the shock is always severe. Embolism is unlikely, and has never, we think, occurred with us. Patients of this type are not very ill; they are in no urgent dread of a fatal termination to their illness, and they rarely see any imperative need for operation. Extreme nervousness is often shown by patients who have, it may be, received exaggerated accounts of the pain of puncture. At the first introduction of the patient to the operation couch we go through all the steps of preparation, not going beyond the injection of the local anaesthetic. When after four days the patient again presents himself, the process is carried farther, and the special needle (we use Saugman's) is pushed in until oscillation of the manometer shows that it is in the pleural cavity. At the third meeting a small amount of gas is introduced. By this method nitrogen may be injected forthwith; it is not necessary to use air or oxygen. The fear is not of embolism but of shock.

If the oscillation of the manometer is 1 in., the dose given may be 300 c.cm. Should there be any doubt as to the position of the needle, or should the manometer not oscillate freely, then no more than 30 c.cm. should be introduced. The entry of this small amount of gas will not cause embolism, while if the needle is slightly blocked it will push aside the obstruction, on which the oscillations will jump to view. The manometer is the guide.

After the gas has been injected, the patient is made to lie still for half an hour; then, if all is well, he may be sent by ambulance back to the ward.

Some patients suffer from shock even with this careful method. The effects are not necessarily shown on the instant. The operator may be injecting gas into the pleura, encouraged by the smiling comfort of the patient, when suddenly it is seen that all is not right. The drugs used may so have deadened the patient's sensation that he is not mentally aware of his physical distress until a dangerous point is reached. The pulse may quicken and intermit, the patient may become distinctly blue, the respiration may quicken, and pain may suddenly occur and rapidly become excessive. It may be necessary to give morphine to ease the pain; it is at any rate advisable to stop the supply of gas, and also to open the tap of the needle and thus allow the gas to regurgitate, so to speak, from the pleura. The escape of a little gas may at once give ease to the patient and relief to the operator. The severity of the shock in these cases which are unilateral and limited in extent is probably due to the healthy condition of the pleura. It is not thickened; it is not adherent; its nervous mechanism is sensitive, and there is no bar to immediate displacement of the heart. Movement on the part of the patient seems to precipitate the distress. Well do we remember completing a first operation on a girl who thought the whole affair a triviality until we had finished; then she showed acute distress and immediate fear. For a few minutes she was in considerable danger; we had let the gas run in too quickly and had given her too much, and all at once cardiac and respiratory resentment showed itself.

2. Medium Cases.

In medium cases, often with bilateral disease and a considerable amount of pleuritic thickening, the shock of needle puncture is not marked. The difficulty is to find the pleural space and the danger is embolism. It may be necessary to try several times before the manometer oscillates; the oscillation is often slight and may even be due to the needle being in loose tissue immediately outside the parietal pleura. Even when the needle reaches the pleural space, it may be in a pocket, and there may be only the possibility of inducing a very partial inflation. The shock, apart from fear, is not great, and the heart is not violently displaced. The thing to be feared is gas embolism.

If an attempt is made to force in gas at any pressure above 3 in. of water, there may be severe local pain, and in an hour surgical emphysema may be recognizable. The occurrence may be guessed at if there is a feeling of constriction in the neck. The necessary precautions are to limit the first introduction of gas to 50 c.cm., and to watch respiratory rather than cardiac movements. If oxygen is to be given at all it should be given in these cases.

Another point is worth mentioning: after four or five injections it is necessary to alter the point of puncture. We take it that a local patch of pleurisy has resulted from the frequency of the trauma. Another trouble in these chronic cases arises from a change in the pleural lining: it becomes haemorrhagic and is apparently turned into something resembling granulation tissue, and the needle may again and again draw blood, as is shown by the reflux into the bulb, which in our instrument is inserted in the line of india-rubber tubing leading from the gas bottle to the needle. There may then be great anxiety lest a vein has been tapped. If the manometer does not show its initial rise, followed by the welcome oscillations, the operation must be stopped. Often when filling the pleural space slowly the gas at a moment rushes in rapidly; this is due to peeling off the softly adherent pleura; the pressure must be lowered or the gas even shut off for a time, or there may be severe pain or even shock.

Haemoptysis of the severest kind allows any risks; nothing else does. In one patient we introduced 1,800 c.cm. of gas at one time; we did it slowly and it took an hour. There was no shock—indeed the patient said he was quite comfortable, and the haemoptysis, which had threatened to be fatal, terminated. There was just an ounce of blood on the morrow, and then no more.

In all cases of severe haemoptysis threatening life, artificial pneumothorax should be thought of. The only contraindication would be such a massive amount of disease that the pressure of a large quantity of gas in the

pleura might lessen the available breathing space until the minimum necessary for life had been passed.

3. Cases of Very Severe Disease.

In severe cases with bilateral cavities and much of both lungs affected, the danger of artificial pneumothorax is obvious. The patient has little lung left; he may therefore be suffocated if any of that little is taken away. Still, this is not the last word. In these cases we have many times given the patient the only chance left to him, and in a few instances the result has been good.

The apparatus made at Dr. Woodcock's suggestion by Messrs. Reynolds and Branstons, Limited, of Leeds, seems to offer some advantages over the instruments used by most workers, and the plan of it has been approved by a great Continental authority. It is well made and accurate, easily kept clean, and simple in its working. In the latest form of this apparatus a regulator is used, which controls all the pipes by manipulating one indicating needle. This regulator is an invention of Dr. T. D. Lister of London.

EIGHTY-SECOND ANNUAL MEETING

OF THE

British Medical Association.

Held in Aberdeen on July 29th, 30th, and 31st.

PROCEEDINGS OF SECTIONS.

SECTION OF LARYNGOLOGY, RHINOLOGY, AND OTOLOGY.

H. LAMBERT LACK, M.D. Lond., F.R.C.S., President.

ABSTRACT REPORT.*

PRESIDENT'S INTRODUCTORY REMARKS.

THE President, after welcoming the members, went on to consider the significance of the two subjects which had been selected for general discussion—the treatment of inoperable growths of the nose and throat, and oto-sclerosis in its various aspects. The fact that the former had been chosen was itself an indication that some progress had been made in the direction of alleviation, if not of cure.

Marvellous progress had marked the last year, as was shown by the introduction by Killian of suspension laryngoscopy and by Nagelschmidt of diathermy, which had been found useful not only in malignant disease, but also in intractable lupus of the upper air passages. In conclusion, the President alluded to the use of the intra-tracheal administration of ether as a great step in advance, and to the efforts which were being made by Freeman to provide a rational and successful method of treating hay fever.

DISCUSSION ON THE TREATMENT OF INOPERABLE GROWTHS OF THE THROAT AND NOSE.

OPENING PAPERS.

I.—W. D. HARMER, M.C. Cantab., F.R.C.S.,
Surgeon, Throat Department, St. Bartholomew's Hospital, London.

DIATHERMY.

THE subject of treatment by diathermy had been discussed, Mr. Harmer said, by Nagelschmidt in his *Lehrbuch der Diathermie* (Berlin, 1913). A paper on the treatment was also read by him before the Electro-therapeutical Section of the Royal Society of Medicine in 1910.

The apparatus produced a high frequency current of great power. It had two electrodes, one large which was wrapped in wet cloths and applied to the patient's chest, and the other small which was applied to the growth. The passage of the current produced intense heat at the

small electrode only. The instrument must be controlled by a skilled person, and the operator should wear rubber gloves to avoid shocks; the current was not turned on until the small electrode was buried in the growth, and was turned off before the electrode was removed from the tissues; this was necessary in order to avoid sparking; strict aseptic precautions were necessary. A general anaesthetic was required for large growths. Atropin internally and frequent sponging locally were necessary to keep the parts dry. The passage of the current blanched the tissues around the electrode in a few seconds. After a series of punctures had been made the cauterized areas might be removed by forceps or curette and the deeper parts of the growth attacked. Although bleeding did not result from the burning of even vascular growths it was necessary not to remove sloughs too boldly. In large tumours dyspnoea during or subsequent to the operation might require tracheotomy. Too rapid burning was to be avoided as it might lead to dangerous swelling of the neck. More than one sitting was necessary, as a rule.

Mr. Harmer had treated by this method 21 cases of cancer of the tonsil, tongue, palate, pharynx, and larynx. Naevi of the nose and tongue and naso-pharyngeal fibroma had also been successfully dealt with. The subsequent pain was slight, there was rarely any oedema or inflammation, and the patient was able to swallow about forty-eight hours after the operation. Sepsis was less frequent than after cutting operations, and there was no shock. The wounds left healed rapidly, with but trivial scarring. Nearly all the cases he had treated were enormously improved, and in one instance (cancer of the tonsil and tongue) life was prolonged in comfort for about two years. These were striking results in what were cases of very advanced disease. Cases in which the growth had reached the neck or had invaded bone were unsuitable for this treatment.

The advantages over the knife were the rapidity and ease of operation, the absence of haemorrhage, and the sealing up of the vessels and lymphatics by the thermal agent. Mr. Harmer believed that diathermy retarded the subsequent progress of the disease, both in its original site and in the cervical glands. Local recurrence seemed to be less frequent than after a cutting operation. For simple vascular tumours, such as naevi, the results were all that could be desired.

II.—WILLIAM HILL, M.D.,

Surgeon for Diseases of the Throat, Nose, and Ear, St. Mary's Hospital, London.

RADIUM.

RADIUM, Dr. Hill said, was worth trying in inoperable cancer in these regions if an adequate amount were available, if the primary growth were accessible and not too extensive, if metastases were absent, and if the general health was fairly good. When extensive growths were irradiated massive doses of toxins were liberated and might lead to serious complications. In pharyngo-laryngeal cancer radium should not be tried if the growth had invaded the lower air passages or if there was marked glandular involvement.

Sarcomata and endotheliomata reacted better to radium than cancers, but nevertheless some epitheliomas of the nose, throat, and gullet did react well. So well did sarcomata react that accessible sarcomata of the jaw and naso-pharynx might be treated in the first instance with radium and operation postponed until the effect of the radium was manifest. This applied particularly to sarcoma of the tonsil.

As regards the quantity of radium to be used, Finzi held that the equivalent of at least 50 mg. of radium bromide—that is, about 26 mg. of the metal, should be employed. Success might follow the use of smaller quantities, but the necessary increase in the exposure time was inconvenient and even dangerous in cavities like the throat and gullet. Doses as large as 100 to 200 mg., suitably screened, were advantageous in that they permitted a decrease in the time of exposure. There was no reason to believe that the salt itself was more deleterious than the emanation. The great advantage of emanation was that special tubes to suit regions might be employed. Dr. Hill believed that efficient screening was of importance in the case of oesophageal cancer. The needles used by Mr. Pinch,

* A full report has appeared in the *Journal of Laryngology*.