LEECHES.

By A. E. SHIPLEY, Sc.D., F.R.S.,
MASTER OF CHRIST'S COLLEGE, CAMBRIDGE.

Hardly anything real in the shop but the leeches and they're second-hand. —BOB SAWYER.

As Mr. W. A. Harding has pointed out, eleven species of fresh-water leeches occur in these islands. But one of these, the *Hirudo medicinalis*, seems to be vanishing, and yet it is just the one we should cherish and preserve.

There are people who don't like leeches. This is shown by the agitation amongst the travellers in an omnibus, as depicted in *Punch* by Leech, years and years ago, when an old gentleman had upset a bottle of them in their midst. But the medical leech, which is our theme, is really the friend of man and of the soldier, and is a beneficial and not a harmful animal. There are, of course, other leeches in our rivers and in our seas, but of the latter our knowledge is scanty and it is difficult to increase it at present, at any rate in the Channel or in the North Sea, and in any case they have no human interest except the influence they exercise on our fish-food supply, and this is practically negligible.

Zoologically speaking leeches are undoubtedly degenerate Earthworms (*Oligochaeta*), and some very interesting "Zwischenformen"—in spite of the war we may as well employ a useful expression captured from the enemy—have been found in Russia and Siberia, forms which combine many of the characters of the *Oligochaeta* and the *Hirudinea*. Possibly the degeneracy which leeches are said to exhibit is associated with a semi-parasitic habit of life. But a semi-parasitic habit does not apply to all leeches, in fact it applies but to few genera; there are many others, equally degenerate—if degenerate they be—who have no trace of semi-parasitism.

A curious thing about leeches is that all the varying genera have the same number of somites or segments, and though some of these segments or somites are masked and fused, when analysed by the number of segments in the embryo, by the number of the nerve ganglia, and so on, leeches seem always to have thirty-four such segments. These do not correspond with the rings or annulations so visible on the outside, but a certain number of these annulations, varying in each species, "go" to each somite, and so constant are these numbers that it would not be very difficult to represent any given species of leech by a mathematical formula.

The known species readily fall into two sub-orders: (1) The *Rhynchobdellae*, which are marine and fresh-water leeches with colourless blood, with no jaws, and with an extensile proboscis; and (2) the *Arhynchobdellae*, which are all fresh-water or terrestrial, with red blood, and generally with jaws. There is no extensile proboscis, and the anterior sucker has a ventral aspect, and is in no way distinct from the body. There are always in this group seventeen pairs of nephridia. We shall have mostly to do with the latter sub-order.

Hirudo medicinalis, the medicinal leech, is found in

stagnant waters throughout Europe and the western parts of Asia. It is rather commoner in the southern parts of Europe than in the north. It used to be common enough in England, where it was bred, but already 100 years ago its numbers were diminishing.

In a treatise on the Medicinal Leech, published by J. R. Johnson in the year 1816, he records "formerly the species was very abundant in our island, but from their present scarcity owing to their being more in request among medical men, and to the rapid improvements which have of late years taken place in agriculture, particularly in the draining and cultivation of waste lands, we are obliged to receive a supply from the Continent, chiefly from Bordeaux and Lisbon." In his time he considered that for every native leech employed at least a hundred foreigners were used.

The same scarcity was very apparent to the poet Wordsworth, whose insatiate curiosity is recorded in the following lines in 1802—Wordsworth was always a curious man:

My question eagerly did I renew,
"How is it that you live, and what is it you do?"
He with a smile did then his words repeat:
And said that, gathering leeches, far and wide
He travelled; stirring thus about his feet
The waters of the pools where they abide.
"Once I could meet with them on every side;
But they have dwindled long by slow decay;
Yet still I persevere, and find them where I may."

In Europe, where the leech was once very abundant, it is now chiefly confined to the south and east, and in Germany it is still found in the island of Borkum and in Thuringia—but just now we need not trouble ourselves very much about their distribution in Germany.

In 1842 leeches were occasionally found in the neighbourhood of Norwich, and there are villagers still living in Heacham in Norfolk who remember the artificial leech-ponds. In the middle of the last century the medicinal leeches "of late years . . . have become scarce." At about the same time it is recorded that they were becoming scarce, though still to be found, in Ireland. Apparently the species is now practically extinct in England, although I know of a naturalist who can still find them in the New Forest, but he won't tell anybody where. If they were getting scarce in the beginning of the nineteenth century they are far scarcer now. For there is no leech in London. At least there are only a dozen or two, and they, like those of the firm "Sawyer late Nockem'orf," are second-hand. I have just heard that there is a similar shortage in North America. And yet leeches are wanted by doctors. But Harding tells us that:

Hirudo medicinalis is not the only leech which has been used in phlebotomy. *Hirudo troctina* (Johnson, 1816), occurring in North Africa and in Southern Europe, where it is perhaps an

Fig. 1.—*Hirudo medicinalis*; about life size. 1, Mouth; 2, posterior sucker; 3, sensory papillae on the anterior annulus of each segment. The remaining four annuli which make up each true segment are indicated by the markings on the dorsal surface.

Fig. 2.—View of the internal organs of *Hirudo medicinalis*. On the left side the alimentary canal is shown, but the right half of this organ has been removed to show the excretory and reproductive organs. 1, Head, with eye spots; 2, muscular pharynx; 3, first diverticulum of the crop; 4, eleventh diverticulum of the crop; 5, stomach; 6, rectum; 7, anus; 8, cerebral ganglia; 9, ventral nerve cord; 10, nephridium; 11, lateral blood vessel; 12, testis; 13, vas deferens; 14, prostate; 15, penis; 16, ovary; 17, uterus, a dilatation formed by the conjoined oviducts.

introduced species, was largely imported at one time for medical uses.

Several other species have been used for blood-letting in different countries. *Limnatis (Pezomachus) granulosa* in India, *Hirudo officinalis* in Mexico, *Hirudo nipponia* in Japan (Whitman), and *Macrobdella decora* in the United States (Verrill), are or have been used in phlebotomy.

Our chief hope seems to lie in India.

Leeches are still used much more than the public are aware. One pharmaceutical chemist in the West End of London tells me he sells between one and two thousand a year, and as they are bought wholesale at about one penny each and sold retail at about sixpence, there is some small profit.

Leeches were well known to the ancients, and it would be easy to quote case after case from the classical medical authorities of their use in fevers and headaches and for many ill-defined swellings. They were frequently used for blood-letting where a cupping-glass was out of the question. With his curious uncritical instinct, Pliny records that the ashes of a leech sprinkled over a hirsute area or formed into a paste with vinegar and applied to the part will remove hair from any region of the body. Leeches were also used by the Greek and Roman physicians in angina accompanied by dyspnoea.

Probably the traffic in leeches reached its height in the first half of the nineteenth century. Harding reminds us that in the year 1832 Ébrard records that 57½ million of these annelids were imported into France, and by this time the artificial cultivation of leeches had become a very profitable industry. Although in a small way leeches may have been cultivated in special ponds in Great Britain, the English never undertook the industry on a large scale. In Ireland the natives used to gather the leeches in Lough Mask by sitting on

Fig. 3.—Head of a leech, *Hirudo medicinalis*, opened ventrally to show the three teeth and the pharynx *p*, with its muscles; *s*, a nephridium.

the edge of the pool dangling their legs in the water until the leeches had fastened on them. But the native supply was totally inadequate, and the great majority of leeches used in this country were then imported. In 1842 Brightwell mentions a dealer in Norwich who kept a stock of 5,000 of these annelids in two large tanks. The traffic, as we have seen, was very considerable.

The French merchants recognized five classes as follows:

1. Les filets ou petites Sangsues, qui ont de un à cinq ans;
2. Les petites moyennes, qui ont de cinq à huit ans;
3. Les grosses moyennes, qui ont de huit à douze ans;
4. Les mères Sangsues ou les grosses, qui sont tout à fait adultes;
5. Les Sangsues vaches, dont la taille est enorme.

They also recognized many colour-varieties, of which we need only mention the speckled, or German-leech "*Sangsues grise medicinalis*," with a greenish-yellow ventral surface spotted with black, and the green Hungarian leech, with olive-green spotted ventral surface. Both are merely colour-varieties of *Hirudo medicinalis*, which shows great variation in colour, and often forms colour races when bred artificially.

The varying sizes of the five categories mentioned above may be seen by the fact that one thousand of "les filets" weigh from 325 to 500 grams, one thousand of "les petites moyennes" weigh 500 to 700, one thousand of the

"grosses moyennes" weigh 700 to 1,300 grams, and one thousand of the "grosses" 1,300 to 2,500 or even to 3,000 grams. Whereas one thousand of "les vaches" weigh up to 10 kilograms, and sometimes even more. To increase their weight the dishonest dealer sometimes gives them a heavy meal just before selling them. They are transported from place to place in casks, half filled with clay and water, or in stone vases full of water. Sometimes they travelled in sacks of strong linen, or even of leather, and these have to be watered from time to time.

One of the modes of moving them was to place them in baskets full of moss or grass soaked in water, but care had to be taken lest they escape. These baskets could not be packed one on another, or the leeches were crushed. In the old days each sack often weighed 20 to 25 kilograms, and travelling thus, suspended in a kind of hammock, *dans une voiture ou fourgon*, from Palota near Pesth, reached Paris in from twelve to fifteen days. They generally travelled via Vienna to Strassburg, where twelve great reservoirs were appropriately placed near the hospital, and here they rested for awhile. Others collected in Syria and Egypt came by ship to Trieste, whence they are sent to Bologna, to Milan, and to Turin, or by water to Marseilles. Marseilles also received directly by sea the leeches from the Levant and Africa, and expedited them to Montpellier, Toulouse, and many other towns in the south.

The best time of year for their journey was found to be the spring and autumn. They were more difficult to manage in the summer, and they were all the better for having a rest every now and then, as they used to do at Strassburg. There were

Fig. 4.—*Hirudo medicinalis*. *a*, Anterior sucker covering triradiate mouth; *e* points to an annulus midway between the male and female openings, *s* to a nephridium, *u* to the bladder of the latter; *a*, anus. Four testes and four lateral diverticula of the crop are also shown.

days when consignments of from 60,000 to 80,000 used to leave Strassburg for Paris. As in England, however, for the most part the artificial cultivation of leeches has now ceased in France, though half a century ago leech farms were common in Finistère and in the marshes in the neighbourhood of Nantes. There are some years when, if the season was favourable, the peasants carried to market 60,000 a day. Spain and Portugal also furnished them for a long time, but by the middle of last century the Peninsula had become very depleted. But some leeches were still at that period being received from Tuscany and Piedmont. Perhaps the richest fields which still exist are the marshy regions in Hungary. In 1806 a thousand leeches in France fetched 12 to 15 francs, but in 1821 the price had risen to 150 to 200 and even 283 francs. In the latter year they were retailed at 20 to 50 for 4 to 10 sous.

It will be observed that, probably without the knowledge of General Joffre, Field Marshal French, the Grand Duke Nicholas, and even without the knowledge of the German Emperor, the Allies are fighting the Germans on some of the best leech areas in Europe—a point to which we shall return when dealing with the leeches of the Orient. One wonders what the leeches think of it all.

There is no doubt that the medicinal leech is one of the most beautiful of animals. Many of its cousins are uniform and dull in colour—"self-coloured," as the drapers would call them—but the coloration of the medicinal leech could not be improved upon. It is a delicious harmony of browns and greens and blacks and yellows, a beautiful soft symphony of velvety browns and greens and blacks, the markings being repeated on each segment, but not to the extent of a tedious repetition. So beautiful are they that the fastidious ladies who adorned the salons at the height of the leech mania during the beginning of the eighteenth century used to deck their dresses with embroidered leeches, and by repeating the design one after the other constructed a chain of leeches which, as a ribbon, was inserted around the confines of their vesture.

Harding tells us that the dorsal surface of *H. medicinalis* is "usually of a green, richly variegated colour, with orange and black spots, exhibiting an extremely variable pattern, based generally upon three pairs of reddish-brown or yellowish more or less longitudinal stripes, often interrupted by black or sessile spots occurring on the rim of each somite. The ventral surface is more or less green, more or less spotted with black, with a pair of black marginal stripes."

The shape of the medicinal leech, and indeed of other leeches, is difficult to put into figures, as its body is as extensible as the conscience of a politician. The length of *H. medicinalis* in extreme extension ranges from some 100 to 125 mm.; in extreme constriction 30 to 35 mm. The width in the former state would be 8 to 10 mm., and in the latter 15 to 18 mm.

The movements of the medicinal leech are as graceful as its colour is tasteful. When in the water they move like looper-caterpillars (Geometrids), stretching out their anterior sucker, attaching it to some object, and then releasing the posterior sucker they draw the body up towards the mouth. Or, casting loose from all attachment, the leech elongates and at the same time flattens its body until it assumes the shape of a band or short piece of tape, and by a series of the most seductive undulations swims through the water. Kept in an aquarium they are rather apt at times to leave the water and take up a position on the sides of their home an inch or two above the aqueous surface. When outside the water they keep their bodies moist by the excretion of their nephridia and this fluid plays the same part in a leech as the coelomic fluid of an earthworm, which escapes by the earthworm's dorsal pores. There is very little doubt that both these fluids contain some bactericidal toxin which prevents epizootic protozoa and bacteria from settling on their skin, as they do on so many fresh-water crustacea, such as Cyclops, which is a floating aquarium of Ciliata. In fact, leeches, like earth-worms, have a self-respecting, well-groomed external appearance. Like our dear soldiers, they are, so to speak, always clean shaved.

There has been a very widely spread tradition that in their comings and goings in and out of the water, leeches act as weather prophets. The poet Cowper, who throughout his chequered career had always shown but an imperfect sympathy with science, tells us that "leeches in point of the earliest intelligences are worth all the barometers in the world," and Dr. J. Foster mentions that leeches "confined in a glass of water by their motions foretell rain and wind, before which they seem much agitated, particularly before thunder and lightning." Modern opinion, however, prefers the barometer.

The great Chancellor, Lord Erskine, kept a couple of tame leeches, and Sir Samuel Romilly records in one of his decorous letters:

He told us how that he had got two favourite leeches. He had been blooded by them last autumn when he had been taken dangerously ill at Portsmouth; they had saved his life, and he had brought them with him to town, had ever since kept them in a glass, had himself every day given them fresh water, and had formed a friendship with them. He said he was sure they both knew him, and were grateful to him. He had given them different names, Home and Cline (the names of two celebrated surgeons), their dispositions being quite different. After a good deal of conversation about them, he went himself, brought them out of his library, and placed them in their glass upon the table. It is impossible, however, without the vivacity, the tones, the details, and the gestures of Lord Erskine, to give an adequate idea of this singular scene. He would produce his leeches at

consultation under the name of "bottle conjurors," and argue the result of the cause according to the manner in which they swam or crawled.

The medicinal leech lives on the blood of vertebrates or invertebrates, and H. O. Latter records that "cattle, birds, frogs and tadpoles, snails, insects, small soft-bodied crustacea, and worms are all attacked by various species," but the true food of *Hirudo medicinalis* is the blood of vertebrates. The three teeth, which cause the well-known triradiate mark on the skin, are serrated and sharp. The strong sucking pharynx has its wall attached by numerous muscles to the under side of the skin of the leech. By the contraction of these muscles its lumen is enlarged, and by thus creating a vacuum the blood of the host flows in. In the walls of the pharynx and the neighbouring parts are numerous large unicellular glands which secrete an anti-coagulative fluid which prevents the blood of the host clotting, so that even when the leech moves its mouth to another point the triradiate puncture continues to ooze. The same anti-coagulative secretion no doubt prevents the blood coagulating in the enormous crop of the leech in which this meal of blood is stored. Opportunities for a meal presumably occur but seldom, and the leech is the "boa-constrictor" of the invertebrate world. Its interior economy is laid out on the basis of a large and capacious storage and of a very restricted and very slow digestion. The blood sucked into the sucking pharynx passes on to the thin walled crop, which occupies almost all of the space in the animal. This crop is sacculated, having eleven large lateral diverticula on each side. In a fed leech the whole of this crop is swollen with blood, which, as we have said above, is prevented from coagulating by the secretion of numerous unicellular glands in the pharynx and neighbouring parts. The actual area where the digestion takes place is ludicrously small, as shown in

Fig. 5.—Cocoon of the medicinal leech, and longitudinal and transverse views of the same cut open.

Fig. 2. The rectum, which runs from the real seat of assimilation to the opening of the posterior sucker, transmits the undigested food, and there's not much of it.

An active leech will draw from one to two drachms of blood, and as much more will flow from the wound when the leech moves, because the blood does not begin to coagulate owing to the secretion of the anti-coagulative. No scab or clot is formed. The flow of blood can, if necessary, be stimulated by hot fomentations. Sometimes the bleeding is so great that artificial means have to be taken to check it. When leeches are applied to the human integument they are generally first dried in a cloth, and if they will not bite the part required the part should be moistened

with sweetened milk or a drop of blood. To remove leeches when replete salt, sugar or snuff sprinkled over the back are used. They may be made to disgorge by placing them in a salt solution of 16 parts salt and 100 of water at 100° F. A full meal is said to last leeches nine months.

Leeches are hermaphrodite, and in some genera the acting male inserts spermatophores or little cases containing spermatozoa anywhere in the skin of the leech that is being fertilized, and the spermatozoa then make their way through the tissues of the body till they arrive at the ovary and there unite with the ova. In the medicinal leech the mating is said to be encouraged

by adding fresh water to the vessels in which the leeches are living.

The eggs are laid in capsules or cocoons attached to some water plant or buried in the mud, about twenty-four hours after the leeches have mated. The cocoon is formed as it is in an earthworm, by the secretion of certain glands in the skin forming a secretion which hardens and takes the form of a broad ring, as it were, round the body of the leech. Through this broad ring the body of the leech is with-

Fig. 6.—A *Nephelis* forming its cocoon and withdrawing from it.

drawn and the fertilized eggs are deposited in it, and the two ends close up, but not entirely, for the young leeches eventually make their way into the outer water through one of the remaining pores. Within the cocoon are 6 to 20 ova and these gradually mature and the young hatch out. When they leave the cocoon they are minute, and of the thickness of a pack-thread. More than one cocoon is deposited by each leech, and unless the cocoons are anchored to some submerged object they often rise to the surface of the water and float half submerged, and are apt to be destroyed by water rats, voles, and other enemies of leeches. At times the leeches themselves destroy the cocoons.

The exact time of the emergence from the cocoon does not seem to be very definitely known, but leeches are long-lived annelids. It is not till their third year that they are of any use for medical purposes, and they are said not to pair until they are six or seven years old. They certainly live twelve or fifteen years. But the fact that they grow up so slowly and live so long shows that it will be difficult to replace the shortage of leeches in Great Britain and Ireland during the present war. This could hardly be done by home culture, for, even if the war lasts three or four years, we have lost the cocoons of the summer of 1914, even if we ever had them.

Leeches have many enemies—water rats, voles, the larvae of the *Dytiscus* beetle, the larvae of *Hydrophilus*, the *Nepa* or water scorpion, the larvae of the dragon-fly, and the adult *Dytiscus* all feed upon them. Many birds also eat leeches, and it is recorded that at one artificial leech farm, where there were 20,000 leeches, they were all eaten up in twenty-four hours by an invasion of ducks. Frogs and newts also devour them, and they are not above eating their own brothers. *Aulostoma* will devour its own species as readily as it will an earthworm.

Those artificially reared, as is usually the case with animals reared in captivity and probably against their will, are peculiarly liable to disease of various sorts. They not only get disease themselves, but they act as carriers of disease and play the same part to fish which biting insects play to man and other terrestrial animals. They convey to fishes protozoal diseases similar to those that insects convey to man and other warm-blooded vertebrates.

Leech farming used to be a profitable undertaking, but now it has almost fallen into desuetude. Leeches are, however, still cultivated in some parts of the world, and in America, Latour describes a farm, situated at Newton in Long Island, where there is, or was, a leech farm some thirteen acres in extent. The farm consists of oblong ponds of about $1\frac{1}{2}$ acres, each 3 feet deep. The bottom of each pond is covered with clay, and the banks are made of peat. The French writers recommend, as a rule, the use of clay for the banks. The "eggs" (cocoons) are deposited in the peat from June onwards till the weather gets chilly. The adult leeches are fed every six months with fresh blood placed in stout linen bags suspended

in the water. A more cruel method of feeding these domesticated leeches is that of driving horses, asses or cattle into the ponds, and this was adopted in the leech farms of France.

Some leeches show a considerable amount of maternal affection. *Glossosiphonia heteroclita*, for instance, carries its eggs about with it, and *Helobdella stagnalis* has its little young larvae attached by their tiny suckers to the mother's back, which they are loath to leave.

Aulostoma gulo, the horse-leech, is, as we have said above, a very ferocious feeder. Exactly why this species is called a horse-leech is a matter of speculation, but "horse" used as an adjective seems to imply something large and something rather coarse—for instance, horse-chestnuts, horse-play, horse-sense, and horse-laugh.

The rapacity of the daughters of the horse-leech is dwelt on in the Bible.¹ I can make no claim to be an authority on exegesis, but I am never quite sure whether these two ladies were not the offspring of the local veterinary surgeon. But *Aulostoma* occurs in Palestine, and its voracity may very well have been known to the Hebrews.

I entirely reject the idea that the word indicates some ghost or phantom; that explanation is due to the craven policy of taking refuge in the unknown.

I conclude with a couple of sentences taken from Dr. Phillips's *Materia Medica* on the present use of leeches:

The special value of leeching is shown in the early stage of local congestion and inflammations, such as arise from injuries, and in orchitis, laryngitis, haemorrhoids, and inflammations of the ear and eye,

cerebral congestions, and congestive fixed headache.

Leeches are also of service in a manner less easy to understand in inflammations of deep-seated parts without direct vascular connexion with the surface—for example, in hepatitis, pleuritis, and pericarditis, as well as in pneumonia, peritonitis, and, according to some observers, in meningitis. In all these disorders, however, they are very much less used than formerly—in the larger hospitals, for instance, when at one time they cost many hundred pounds annually, a few dozens in the year would represent the total employed.²

REFERENCES.

- ¹ Campbell's *Lives of the Chancellors*, vol. vi. ² Proverbs xxx 15. ³ *Materia Medica and Therapeutics*. By Charles D. F. Phillips, p. 1015.

M. EMILE COMBES, formerly Prime Minister of France, and still a member of the French Senate, has founded a hospital for the wounded in Pons, of which he is Mayor. M. Combes, who is a doctor of medicine and has had a large experience as a practitioner, is acting as chief physician to the hospital and spends all his time in ministering to the patients.

Fig. 7.—Cocoons of *Nepheleis*, showing the growth of the eggs and the issuing larvae, which, in the lower figure, are leaving the cocoons.

Fig. 8.—*Helobdella stagnalis*, with adhering young. Ventral view, magnified. (From Harding.)

Fig. 9.—*Glossosiphonia heteroclita*, with eggs and emerging embryos. Ventral view, $\times 4$. (From Harding.)

Fig. 10.—A leech farm in the south of France.