Case 22.

No. 1165. Coal-miner (26). Whilst standing astride, a pony backed and struck the inner side of his left knee, and he fell, twisting round to the right with the left knee inwards. Felt severe pain and a "click," as if something slipped into its place on rising. A week later, whilst hewing coal, with the knees slightly bent and rotating the left femur inwards as he used the pick, he suddenly felt "something go out" in his left knee. As he straightened the leg with difficulty something slipped in again. The cartilage has been out five times altogether. The slightest twist now displaces it.

Left meniscus torn near its convex border from near the capsule = bucket-handle tear.

II. TORN EXTERNAL CARTILAGES.

Case 1.

No. 831. Coal-miner (22). Whilst lifting a tub (small wagon) with his hands behind his back and the knees flexed, he slipped and twisted round to the left on his left leg, which was fixed. He felt acute pain in the left knee, and something "gave" in it, and then he doubled up. The knee swelled, and he could not straighten the limb. Later he got the same knee jammed between a tub and a prop, when the knee was bent in, and it swelled up again. Knee can be fully extended, but not completely flexed.

Left outer cartilage was split across transversely at its middle and the posterior half projected into the joint. Note that rotation of the thigh *outwards* caused tearing of the *outer* cartilage, whilst rotation *inwards* causes tearing of the *internal* meniscus, and that displacement of the posterior half interfered with flexion.

Case 2.

No. 934. Ship's stoker (39). Whilst right foot was fixed and knee bent he was forcibly twisted around to the left. Pain over outer cartilage on full extension.

Right meniscus torn across about its middle. Anterior half removed. (This case is against the rule, but my cases of torn external cartilages are too few to reason from as to the exact causation.)

SUMMARY OF CASES.

Torn inner cartilages	...	...	22
Torn outer cartilages	...	...	2-24
Right knee	...	...	14
Left knee...	...	...	10

Ages.—18 to 48 years.

Sex.—Males, 24; females, 0.

Occupations.—Coal-miners, 15; football accidents, 2. The chief industries in this district are coal-mining and ship-building.

NATURE OF TEAR.

A. In Inner Cartilages (Total 22).

1. In posterior half only	...	...	3
2. Torn anterior attachments only	...	...	2
3. Torn anterior attachments and from anterior part of capsule	...	...	5
4. Bucket-handle tears—that is, from the capsule only:			
(a) In whole length	...	...	3
(b) In middle half	...	...	2
(c) In anterior third	...	...	1-6
5. Longitudinal splints in substance of cartilage	...	...	2
6. Transverse tears in anterior half	...	...	4

B. In Outer Cartilages (Total 2).

Both had transverse splits	...	...	2
----------------------------	-----	-----	---

THE *Bulletin administratif du Ministère de l'Instruction publique* gives statistics as to the students in different faculties in France on January 15th, 1913. At that time the total number of students in the faculties of medicine was 8,247, consisting of 7,400 men, of whom 803 were foreigners, and 847 women, of whom 489 were foreigners. The total number of students in the schools of medicine was 1,497; of that number 1,287 were men, of whom 29 were foreigners, and 210 women, of whom 9 were foreign. These numbers are almost exactly the same as those for the previous year. For the several faculties the numbers were as follows: Paris, 4,211; Lyons, 1,055; Bordeaux, 751; Montpellier, 699; Toulouse, 346; Nancy, 306; Lille, 285; Algiers, 156. Of the various nationalities represented, the list was headed by the Russians, who numbered 831, of whom Paris had 585 and Montpellier 132. Next came the Turks with a total of 85, of whom 68 were studying in Paris and 11 at Montpellier. There were 67 Bulgarians—20 at Lyons, 28 at Montpellier, and 12 at Nancy. The Roumanians numbered 59, of whom 54 were studying in Paris and 5 at Montpellier. Of other nationalities there were 13 English and 24 Greeks in Paris, while the South American Republics were represented by 22, all in Paris.

A POSSIBLE FALLACY IN THE DIAGNOSIS
OF INTERNAL DERANGEMENTS OF
THE KNEE-JOINT.

By SPENCER MORT, M.D., CH.B.GLASG., F.R.C.S. EDIN.,
MEDICAL SUPERINTENDENT, EDMONTON INFIRMARY.

THE interesting abstract in the *EPITOME* of the *BRITISH MEDICAL JOURNAL* (November 15th, 1913, par. 255) of a recent article by Bahr¹ on spurs of the patella in relation to inflammation of the knee-joint, recalls some examples of a similar deformity which I have recently studied.

In few cases of displaced, loose, or fractured semilunar cartilages are all the prominent signs found. Allingham² recounted the classical signs as follows: Tenderness over the upper part of the tibia, fixation of knee in semiflexed position, inability to extend the knee though flexion is free and painless, cracking of the joint on movement, weakness on walking or supporting the body on the limb, rotation of the foot, and synovitis. In a summary of his cases he describes three varieties:

1. The internal or external semilunar cartilages may be torn away from their—

- (a) Anterior attachment.
- (b) Posterior attachment.
- (c) Coronary ligaments.
- (d) Coronary ligaments and anterior attachment.
- (e) Coronary ligaments and posterior attachment (rare).

The cartilage at its injured part is completely movable.

2. All or some of the attachments of the semilunar cartilages may have been stretched or partially ruptured.

3. Split cartilages.

Recent operative measures have shown the frequency of actual fracture of the cartilages, or in many cases of the existence of rheumatoid arthritis causing irregular projecting synovial fringes with fibrosis and irregular destruction of the cartilages, which easily become displaced over the typical lipping or moulding of bone of a "mushroom"-like form accompanying these changes.

In many cases pain in the knee is the chief symptom, drawing attention to deformity elicited on palpation from projection of a cartilage, or to rheumatoid change in the bones and joint with recurrent synovial effusion. In elderly patients it is always wise to look for rheumatoid conditions before diagnosing traumatic displacement, even when an injury first leads to the examination of the painful joint affection.

In three such cases seen recently operation showed little or no change in the interior of the joint, but there were periarticular rheumatoid affections of a spur-like character simulating it. The symptoms were pain, swelling, and clicking sensations on movement, and the condition was one resembling that got in *hanche* or *genou à ressort*, thickened bands of tendon or fascia gliding suddenly and precipitately over bony prominences.

The site was internal in all three cases, the tendon of the sartorius muscle riding upon a distinctly lipped portion of the inner border of the femoral condyle. Examination of the borders of the patella, tibia, and outer surface of the femur should in every case be made, and inquiry made into the condition of other joints. In the three cases—two males and one female—injury was adduced as the supposed cause; incipient rheumatoid arthritis was the real cause, and operation was beneficial in allowing the deformed spur of bone to be burred off, though signs of typical rheumatoid change were developed afterwards.

I believe such cases with *external* joint trouble, and others with typical *internal* mischief, often referred to internal traumatic damage, to be much more commonly the resultant of rheumatoid disease than is usually diagnosed. Sometimes they are due to exostoses, in other cases tuberculosis may be a factor. The importance of their clear recognition, especially in cases of compensation proceedings for injury, is obvious.

REFERENCES.

¹ *Deut. med. Woch.*, July, 1913. ² Allingham: *Internal Derangements of the Knee-joint*.