At the hospital it was found that there was blood in the patient's motions and also in the urine.

Widal's test being negative, he was given calcium lactate, gr. x t.d.s., and a provisional diagnosis of Henoch's purpura was made.

As there was no improvement at the end of a week, his blood was sent to the Clinical Research Society to be tested against the paratyphoid group of organisms. The report, dated February 11th, 1913, was:

The serum agglutinates paratyphoid B in 1 in 10, 1 in 25, 1 in 50, partial with 1 in 100, but none with 1 in 200. Negative to paratyphoid A. No reaction is obtained in controls of either organism made with normal serum.

The dose of calcium lactate was doubled, still without benefit. At the end of a fortnight his condition was very much as on admission, though the spasmodic attacks of abdominal pain were rather less frequent. The purpura continued to break out over his body, fading in one place and appearing in another, and there was still blood in the faeces and urine. The temperature from February 16th till just before his discharge is shown in the chart.

It was my custom to visit him about once a week, and on February 27th Dr. Rattrie and I again examined the

patient carefully. Our opinion was that sepsis somewhere was keeping up the condition, and on looking into his mouth we found it full of septic and carious stumps.

On March 1st, under an anaesthetic, all the decayed teeth were removed. The result on the temperature is seen in the chart. From that day all his symptoms disappeared, and he left the hospital practically well. In a few weeks he was back at work, and he has remained well since.

FURTHER NOTE UPON A CASE OF HYSTERICAL MONOPLÉGIA FOLLOWING ELECTRIC SHOCK.

By PURVES STEWART, M.D., F.R.C.P.,
PHYSICIAN TO WESTMINSTER HOSPITAL.

THIS case was recorded in a previous number of the BRITISH MEDICAL JOURNAL (December 20th, 1913, p. 1580). It is unnecessary to repeat the history in detail. Briefly, the patient, a young man of 21, after an electric shock to the left hand, developed sudden paralysis of the left upper limb, at first spastic in type, later with complete flaccidity and with total anaesthesia of the limb to all forms of stimuli up to the level of the shoulder-joint (see Fig. 1). It will be remembered that pins could be thrust through the anaesthetic skin without causing bleeding. Nevertheless, the skin of the limb showed normal reaction to a fly-blister; also it sweated normally after pilocarpin administered hypodermically.

Nine months after the accident he was admitted to Westminster Hospital, where, for a further period of over two and a half months various forms of suggestive treatment were tried, including strong electrical stimulation to the paralysed limb, the application of cutaneous irritants—such as *Mucuna pruriens*—to the anaesthetic area of the skin, and so on. Despite these measures, however, the clinical phenomena remained at a standstill, and we therefore decided to send the patient for suggestive treatment to a medical colleague who was

a skilled hypnotist of high reputation. For over two months repeated and assiduous attempts at curative suggestion were made at the psycho-therapeutic clinic, but without producing the slightest effect upon the patient's symptoms. In fact, if anything, he became worse, inasmuch as the limit of cutaneous anaesthesia and analgesia advanced in a proximal direction on to the trunk, so as to include the outer part of the pectoral region in front and a similar area over the scapula behind. By the end of January, 1914, the medical hypnotist wrote that neither he nor his assistant could do anything more for the patient.

Fig. 1.—Hysterical monoplegia of left upper limb. Photograph taken when symptoms had lasted nine months. The black line indicates the upper limit of the anaesthesia. The patient is attempting to elevate both shoulders.

Fig. 2.—Hysterical monoplegia of left upper limb of eleven months' duration. The same patient after the administration of a general anaesthetic. The anaesthesia has completely disappeared, and the motor functions are normal.

Accordingly, the patient having been over two and a half months in hospital, and the paralysis having now lasted eleven months from the time of the original electric shock, we decided, before discharging the patient, to try the effect of giving him a general anaesthetic, in the hope of loosening his obsession during the stage of delirium produced by the inhalation of ether freely mixed with air. This was accordingly done on January 20th. The patient was placed on a couch with chest and arms exposed. He was then given ether and nitrous oxide, with plenty of air,

so as to prolong the stage of excitement. The normal right arm was fixed to the trunk by a bandage; the paralysed left arm was allowed to hang free. Within a few seconds, whilst the anaesthetic was being administered, and without any suggestion on the part of the physician, the patient began to make slight movements of the paralysed arm, and in a short time these amounted to violent fighting movements, in which the patient tried to snatch the mask out of the hands of the anaesthetist. After a

few minutes the general anaesthetic was discontinued, and the patient passed into a hypnotic state, during which voluntary movements of the previously paralysed limb were normally executed in response to verbal commands, reinforced by occasional painful stimuli, such as twisting the ear, pressing the supraorbital notch, pricking the previously anaesthetic limb, etc. During this stage analgesia of the opposite limb was temporarily suggested, and pins could be passed through the skin without pain. Lastly, it was suggested that all his sensory and motor disabilities should disappear, and, as the effects of the ether passed off, this was found to be actually the case.

Ever since his séance with ether and nitrous oxide, the patient has remained able to move the limb normally at all joints (see Fig. 2), whilst all varieties of cutaneous and other stimuli are promptly and normally recognized.

The Milroy Lectures

ON

THE HYGIENIC ASPECT OF THE COAL-MINING INDUSTRY IN THE UNITED KINGDOM.

DELIVERED BEFORE THE ROYAL COLLEGE OF PHYSICIANS OF LONDON.

By FRANK SHUFFLEBOTHAM, M.A.,

M.D.CANTAB., J.P.,

MEDICAL REFEREE UNDER THE WORKMEN'S COMPENSATION ACT FOR THE NORTH STAFFORDSHIRE DISTRICT.

LECTURE II.

CAUSES OF EXPLOSIONS.

UP to a comparatively short time ago colliery explosions were attributed to the inflammability of fire-damp; but in a modern well-ventilated mine, while the explosion of fire-damp may be the initial agent, it plays practically no part in the actual explosion in the workings. An explosion can be produced readily by firing a mixture of coal dust and air, and the proportions necessary to form an inflammable mixture are, roughly speaking, 1 lb. of coal dust to 160 cubic feet of air.

COAL DUST WATERING.

The finer the coal dust the more explosive. While fire-damp explosions begin at the working face, coal dust explosions usually take place in the main haulage roads. The coal dust may be ignited either by a blown-out shot (a shot that had been put into the hole in such a manner as to be blown backwards without blasting the rock or coal), or by the exposed flame of a broken safety lamp, by a fused electric cable, or by sparks emitted from coal-cutting machines. In other cases, however, there has been an initial explosion of fire-damp at the working face, carried through the workings by the ignition of coal dust, and a large number of explosions have been caused through the ignition of fire-damp by naked lights in pits where safety lamps are not employed.

Fire-damp explosions can be and are prevented by improved methods of ventilation and by the early detection of the fire-damp in small quantities by means of the cap over the flame of the safety lamp. In considering the prevention of coal dust explosions, the first problem is how to prevent the accumulation of coal dust. The presence of coal dust is inevitable in a mine. The removal of dust by periodical sweeping of the roadways, sides, and roof has been advocated, but a more effective method is by watering the roadways, a method practised for some years in Germany and Belgium, and successfully adopted in many mines in this country. It is necessary to water the sides and roof as well as the roadways. It is, however, the general opinion that watering can never, for various reasons, be universally adopted, and spraying might under certain conditions have an injurious effect upon the miners' health. Again, falls of ground are more numerous in wet mines than in dry. Dust removal and spraying may be of some use, but they do not entirely remove the danger, though it may be hoped that

as a result of the Government experiments at Eskmeals a possible remedy is near at hand.

POISONOUS GASES.

The poisonous gases found naturally in mines are black-damp, fire-damp, carbonic acid, and, in small quantities, sulphuretted hydrogen and sulphurous acid; as a result of explosives we may also find oxides of nitrogen. Carbon monoxide is invariably generated as the result of an explosion, whether of fire-damp or coal dust.

Black-damp.

This is the residual gas after oxidation of materials present in the coal, oxygen of the air being taken up with the formation of carbonic acid. It is found more or less in all mines, and always in the return air of the upcast ventilating shaft.

The composition of black-damp varies under different conditions, but, speaking generally, it is a mixture containing about 87 per cent. of nitrogen and 13 per cent. of carbon dioxide. It is probably formed in the process of oxidation of iron pyrites which is present in the coal. This sulphide is oxidized to sulphuric acid and sulphate of iron, and as calcium carbonate in the form of calcite is also present in the coal, calcium sulphate is formed with the liberation of carbonic acid, and, according to Haldane, the process may be represented by the following equation:

Black-damp as a rule has a greater specific gravity than air, and consequently is found on the floor in low-lying workings, but there are cases in which black-damp is lighter than air, cases in which the black-damp contains less than 5 per cent. of carbonic acid (Cadman), or when it is mixed with other gases, such as fire-damp. When there is an accumulation of black-damp lighter than air it is most important that it should not be examined except with a safety-lamp, on account of the possibility of the presence of fire-damp. With a falling barometer the percentage of black-damp may be very greatly increased, and to such an extent in some workings that it is necessary for the miners to suspend their operations.

This gas is recognized by the safety lamp from the fact that when present in small quantities the flame will only burn dimly, and when in larger quantities the flame may be extinguished altogether. This is due to the lack of oxygen, and not to the presence of carbonic acid.

The effects of black-damp on the human subject are due to the presence of carbonic acid and the deficiency of oxygen. The initial symptoms are shortness of breath, quickened pulse, and increased respiration-rate, followed by headache and giddiness. If the victim is unable to get away he may be overcome with drowsiness, and if the black-damp be of sufficiently high percentage he may become insensible and eventually die. Before death there may be marked cyanosis, and the heart may continue to beat for some little time after the breathing has stopped. Death is always due to suffocation or to injuries produced as the result of a fall when the miner is overcome by the gas.

Treatment.—It is most important to get the victim into pure air as soon as possible, and if breathing has stopped, artificial respiration should be resorted to without delay. If the heart has not stopped and oxygen be administered at the same time there is every chance of recovery. If oxygen be available in the pit the rescuer should administer the gas to the sufferer if he be unconscious without removing him, and artificial respiration should be carried out by a colleague.

The *post-mortem* appearances due to black-damp poisoning are similar to those of asphyxia.

Fire-damp.

The constitution of fire-damp is at the present time a matter of controversy. Dr. Haldane holds that it is pure methane, and he has found that the fire-damp met with in lead mines, shale mines, and the Cleveland ironstone mines has also reacted as pure methane. On the other hand, Le Chatelier finds that fire-damp in French coal mines apparently consists of two constituents, one of which is combustible and the other non-combustible, the former constituting in some cases about 95 per cent. of the