Then, further, that no aperient medicine should be given by the mouth or by any other route.

That all food should be withheld, and rectal salines may be given after the lower bowel has been cleared by an enema. Whenever an acute abdominal condition is suspected the patient should be kept in a semi-recumbent position, in order that any septic products may gravitate to the pelvis instead of tending to become diffused throughout the upper abdomen. This point needs emphasis, as it is by no means systematically attended to.

That under no circumstances should opium be given to a child whose symptoms in any way suggest the possibility of an acute abdomen being present, although in cases where a definite diagnosis has been made a single dose of opium sufficient to relieve pain is of the greatest advantage to a patient who has to travel some distance to hospital.

In this connexion we suggest that it would be of advantage to the community if cottage hospitals throughout the country could more fully utilize the services of experienced operators for the treatment of acute cases occurring in their immediate neighbourhood.

Lastly, that it will be of advantage to the medical attendant himself, and therefore to his patients, if, where at all possible, he makes a point of being present at operation on a patient whom he has had under his observation, and whom he has sent into hospital for treatment.

REFERENCES.

¹ Mitchell, Contribution to the Study of Appendicitis in Children, *British Journal of Diseases of Children*, 1912. ² Keen's *System of Surgery*, 1908. The Surgery of Appendix Vermiformis, by John B. Murphy, M.D.

SOME OBSERVATIONS ON THE OPERATIVE TREATMENT OF FRACTURES BY METAL PLATES AND SCREWS.*

By WILLIAM SHEEN, M.S., F.R.C.S.,
SURGEON, KING EDWARD VII'S HOSPITAL, CARDIFF.

THESE observations do not profess to be complete, but raise certain points which seem worthy of consideration and discussion. They are based on 25 cases of operation on the long bones in adults by Lane's plates and screws, and on a survey of some of the recent literature of the subject.

Preliminary Remarks on Non-operative Treatment.

It ought to be an axiom that every paper on the operative treatment of fractures should start by emphasizing the necessity and importance of the frequent study and practice of non-operative treatment, and such necessity I desire to emphasize. Particularly does the necessity arise in hospitals with attached medical schools, whence the students go to all parts of the world to practise under all manner of circumstances—it may be far from hospitals, from x-ray apparatus, from special environment, and from specially skilled hands. Many fractures do not need, many cannot get, operative treatment. Fractures are common, occur anywhere, and often have to be treated where they occur. Just as the x-ray should not supplant other methods of diagnosis, so the operative should not cause to be ignored other methods of treatment. Non-operative reduction, often only possible under full general anaesthesia, the intelligent use of suitable fixation apparatus, and the use of massage and movements may none of them get full attention in hospitals where operative treatment is entirely the vogue. There is an obligation upon surgeons, particularly those engaged in teaching, to see that the non-operative treatment of fractures is never neglected.

Selection of Cases for Operation.

The cases for operation can be divided broadly into two classes: First, that in which operation is a necessity to avoid or remedy a deformed, disabled, more or less useless limb; secondly, that in which operation is more likely to give a better result—a nearer perfect limb—than no operation, although skilled non-operative treatment may be

followed by good functional recovery. Cases of the second class should never be operated on except by practised hands, with special tools and in the very special aseptic environment necessary. Given such circumstances, there is undoubtedly this large second class of cases in which better results can be obtained with operative than with non-operative treatment—results which can often be described as perfect—and the operation risks, in cases selected with due regard to the general health and the local condition, are small. Many surgical advances have been brought temporarily into disrepute, not only by the failures resulting from inexperience, but also by the selection of only difficult or late cases for operation. So with fractures, operators at first had many disappointing results; but with increasing knowledge of the imperfect end-results obtained in many cases not operated on came a widening of the operative field, so as to include easy and early cases, and this widening, coupled with improved technique, led to greater operative success, giving patients better limbs, and enabling them to live more useful lives.

Early and Late Cases.

These are two distinct classes. A case is early when the bone can be brought into perfect alignment and the broken ends and pieces imbricated. This may be possible for as long as two months after the injury. A case is late when callus and bone ends have to be removed to obtain alignment. After a week or ten days it becomes increasingly difficult, as time goes on, to obtain perfect imbrication and alignment. The late cases are much less satisfactory to operate on than the early ones. In the late cases the trauma is greater, and there is therefore more risk of sepsis and of shock; the bone ends cannot usually be got into absolutely perfect apposition; it is difficult to avoid some "winding-up" of one fragment on to the other; from removal of bone limb-shortening results. Each late case operated on is a reproach to its early treatment.

Should Fresh Compound Fractures be Operated on?

The answer to this question should not be a universal "No." When the bone has just punctured the skin and then gone back, it is best to wait until the wound is healed, retaining the broken bone in the best possible position meanwhile. Where the wound is a wide one with exposed projecting fragments, plates and screws should be used, and although there will almost certainly be more or less sepsis and they will have to be removed later, a good position is maintained, and often union in a good position results. If such a fracture is left to heal the healing may take many weeks; the operation will be difficult, through much scar tissue, to encounter perhaps irremediable deformity.

The Value of X Rays.

Much of the fascination of this class of operation centres round the beautiful x-ray pictures before and after operation. An x-ray photograph before operation, in whatever plane taken, must have the anticathode of the tube plumb over the seat of fracture; otherwise, in these "shadow pictures," the "deformity will be distorted—either disguised or exaggerated. An x-ray photograph after operation which shows perfect alignment and perfect position of plates and screws may not mean a perfect functional result. Such photographs may be obtained from fractures where sepsis is present, and where later in the case screws loosen and plates become displaced with accompanying deformity; or they may be obtained where there is excess of painful callus or crippling stiffness of a neighbouring joint. The conscientious operator must record his results not only in terms of x-ray photographs, but also having regard to all points which concern the functional capacity of the limb.

Avoidance of Shock.

In operations on fractures of the femur, particularly late operations in debilitated middle-aged patients, there is risk of death from shock. Two such late cases on which I operated died from shock. In another late case I sent the patient back from the operating table because of his weak condition, and operated later. I now operate on femur cases under spinal anaesthesia; stimulants are given before operation, food is not withheld for long; if the patient is particularly weak, subcutaneous infusion of saline is given during the operation; the operation is

* Read before the South Wales and Monmouthshire Branch of the British Medical Association, December 4th, 1913. A number of illustrative skiagrams were shown at the meeting.

expedited as much as possible; this is helped by the present practice of not suturing the deeper wound structures. With these precautions I have had no further trouble.

Some Points in Technique.

Lane's minutely aseptic technique should be followed. To avoid putting even the gloved hands into the wound is a counsel of perfection. With those less experienced and dextrous, and less well equipped with special instruments than Lane, the hands must sometimes enter the wound. There is no objection to the gloves being thick, and they are then less likely to be pricked or torn.

After the usual long incision, the skin should be turned back a little before the cloths are attached to it. In some situations making skin flaps gives best access. The mops of sterile gauze are wrung out of boiled saline; there is usually no ligaturing or suturing; if, because of tension, some silkworm-gut sutures are required in the skin in addition to Michel's clips, each suture should carry two needles, so that the skin may be penetrated from within outwards on each side of the wound; thus there is no risk of germs being carried in from the deeper layers of the skin to the subcutaneous tissues.

The Plates and Screws.

The drill and screw must be appropriately related in size, the drill being the same size as the shank of the screw—that is, the screw without its thread.

Good strong ordinary steel should be used for the plates, and thin plates, the strength of which must be especially cared for, for smaller, lighter bones, such as the radius. Screws should be threaded to the head, and there should be a proper relation between screw heads and holes in plates. Screws must be just long enough to penetrate the compact tissue of the bone. These are points, small but important, to some of which attention may not be given. A plate must have at least two holes on each side of the fracture line; otherwise it may swing. Plates of special material and screws of special design seem to be unnecessary.

Do Screws Loosen apart from Sepsis?

This is a disputed point, and merits further inquiry. To me it seems that they may loosen and that they may not. They will loosen if the drill hole is too big for the screw; if they are not tight and firm in the bone at the end of the operation; if there is much strain on the plate after operation, such strain being either due to the plate not having been fixed accurately (by bending if necessary) in contact with the bone along the whole length of the plate, or to early movements of the limb when it is not adequately supported at the fracture seat. Apart from sepsis screws will, if of proper size for the drill holes, if fixed firmly in healthy bone and not subjected to undue strain, remain tight.

Sampson,¹ detailing Lane's cases in children, concludes that "rarefying osteitis," causing loosening of screws, probably does not occur except as the result of organisms introduced from without. Hey Groves (as I heard him at the recent International Medical Congress, and I trust I am not misquoting) says that screws frequently loosen apart from sepsis, and the proper thing to do is to use screws which perforate the bone right through, and are fixed on the other side by nuts. (This procedure must sometimes add to the technical difficulties of the operation.) My experience is that occasionally the screws loosen apart from sepsis; it occurred in the femur, due, I think, to too great strain on the screws from too early movement.

The Management of the Periosteum.

I have been accustomed to strip the periosteum back; in my early cases (usually not with complete success) to resuture it; in my later cases to ignore it. Subperiosteally the exposure of the bones is easier and more bloodless, and the plates and screws can be fixed more accurately. Macewen² has shown, contrary to the usual ideas, that the function of periosteum is not to produce bone but to limit its production. Wide stripping of the periosteum has in my experience been followed by the formation of excessive painful callus in the case of the femur, but not with other bones. To introduce the fingers into the wound for suturing is a departure from Lane's technique; introduced suture material also adds to the risk of sepsis; it is often impossible to get the periosteum together. Bearing these

facts in mind, I am inclined in future femur cases to try wrapping a piece of detached fascia lata round the plated bone to limit callus formation.

Sampson³ (he is referring to children) writes: "The periosteum receives no special consideration. It is stripped off the bone . . . and no steps taken to restore it." Lane⁴ writes: "The ends of the broken bone are freely defined, the periosteum being divided and separated if necessary." On the other hand, we have Beckman⁵ stating that "The periosteum should be saved wherever possible . . . a plate . . . should be placed on top of the periosteum, and not underneath it." Lane Joynt⁶ writes: "I never disturb the periosteum if it can be avoided."

Callus Formation after Operation.

There are two aspects of callus formation to be considered: too little callus, and too much. As slight mobility favours callus formation, firm fixation of the fragments by plates and screws may delay it. Too little callus is adverse to union only in unhealthy subjects—for example, with syphilis, or in operations on old ununited or mal-united fractures, where the reparative power of the bone is lessened. In the former the condition causing bad general health must be dealt with; in the latter some form of congestion helps union, either by letting the plated limb hang suitably supported or by the use of a congestion bandage. I recommend that in all cases of non-union after plating with no obvious cause a Wassermann reaction should be taken.

Too much callus has only troubled me with the femur, where with perfect asepsis and a perfect x-ray picture I have seen callus formation to such excess as to resemble a sarcoma of the bone. Probably this resulted from wide stripping of the periosteum. It is in such a case that in future I should try fascia lata wrapping.

Stiffness of Neighbouring Joints following Operation.

This has been with me an almost constant trouble with the knee after plating the femur in adults—equally when I have splinted the parts and when I have practised joint movements from the first. Sampson⁷ (cases in children) says: "Early passive movement seems to cause a secondary stiffness" (presumably of neighbouring joints). This stiffness (like too much callus) may occur where there is perfect asepsis and perfect x-ray pictures. It requires long and persevering treatment for its removal.

After-Treatment.

No plating is satisfactory unless the bone junction is perfectly firm so that the weight of the limb can be borne by it. Bearing this in mind, is any external support necessary, and can the patient get about any earlier than with non operative treatment? Sampson⁸ says: "No curtailment of the period usually allowed with conservative treatment is permitted." My experience is that the femur requires fixation (but not continuous extension) as with conservative treatment. With other bones a sling or quite light support is all that is necessary. Careful massage and passive movements are commenced at the end of a week or ten days. Support must be continued for as long as with conservative treatment. "Congestion" treatment is of value after the plating of old fractures.

Occurrence of Sepsis.

Apart from plating of compound fractures *per primam*, I have only had one septic case. This was a femur in which the lower end was broken into many pieces; a plate used had to be removed; union eventually resulted. In 21 cases—some between 3 and 4 years old—the plates and screws remain in and give no trouble.

Comparison with Other Operative Measures.

I have used wire, Lambotte's apparatus and bone collars. Nothing permits such perfect alignment and imbrication as plates and screws. Wire is obviously ineffective. It is difficult consistently to avoid sepsis with apparatus which comes from the bone to the exterior like Lambotte's. Perfect fixation may be impossible with intramedullary splints made from the patient's tibia. It may be that in some cases such an intramedullary splint coupled with the use of a plate or plates will prove the best treatment.

REFERENCES.

- ¹ *Lancet*, August 17th, 1912, p. 433. ² Macewen: *The Growth of Bone*, Glasgow, 1911. ³ *Loc. cit.* ⁴ *BRITISH MEDICAL JOURNAL*, November 30th, 1912, p. 1532. ⁵ *Mayo Clinics*, 1912, p. 675. ⁶ *Lancet*, March 1st, 1913, p. 624. ⁷ *Loc. cit.* ⁸ *Loc. cit.*