ON SOME CLINICAL MANIFESTATIONS OF
CONGENITAL SYPHILIS.*

BY

HUBERT ARMSTRONG, M.D. VICT. AND LIVERP.,

HONORARY ASSISTANT PHYSICIAN, LIVERPOOL INFIRMARY FOR
CHILDREN.

THE appointment of a Royal Commission on Venereal Diseases having recently directed our attention, amongst other matters, to congenital syphilis, and congenital syphilis being unfortunately so prevalent that no practitioner can go very long without encountering it in his work, a consideration of some of its clinical manifestations may be useful at the present time. Some of the cases used as the basis of this paper are taken from the earlier years of my connexion with the Infirmary for Children, and some of them are not typical, but rather of the class into which an element of doubt has entered, but they are none the worse illustrations on that account.

My own experience is confirmatory of the observation that, while no reduction is taking place—rather the reverse—in the number of cases of congenital syphilis seen, the severity of the disease appears to be decreasing. I am sure that there are far fewer of the appalling cases of infants practically rotting away under our eyes than there used to be. Such cases as the following are now almost rare. It occurred five years ago.

A female infant, aged 4 months, had been under treatment in the out-patient department since the age of two months for typical congenital syphilis. She was a first, and I believe an illegitimate, child. The rash had disappeared, and the baby was doing fairly well, though only putting on weight very slowly, when a prolapse of the rectum occurred, which the mother found impossible to keep back. The child lost condition rapidly, and four days later was admitted an in-patient. Half a grain of hyd. *creta* had been taken three times a day continuously. Three days after admission there was a sudden recrudescence of the nasal discharge, and an apyrexial bullous dermatitis developed. Mercurial inunction was immediately started, but the rapid spread of the skin lesions left the whole body one almost continuous raw area, looking, except for the underlying copper colour, as if it had been badly burnt all over. The nasal and aural cartilages sloughed, the nails were shed; at the same time emaciation was very rapid, and in four days what was left of the child mercifully died.

I had another case in the out-patient department a few months ago, where in the four days between the first visit and the second a similar tragedy took place. But these are the only two recent cases I can recall which have succumbed in this fulminating way, despite energetic mercurial treatment. When the next one comes, if I have time, I shall treat it with salvarsan, though whether the result will be different is doubtful. The sudden slaughter of so many spirochaetes and the throwing of their endotoxins into the system of an infant whose life already hangs on a thread are too likely to prove additional weights which will effectually snap the slender link.

Cases with bone necrosis also appear to me to be fewer, and though this might be explained by altered hospital arrangements preventing the subjects of them from coming under my view, still Mr. R. W. Murray and other surgeons tell me that it is also their experience. On the other hand, parasyphilitic disease in childhood (juvenile tabes and general paralysis) and also spastic plegias (cerebral syphilis) seem to be on the increase—sequels which may very naturally be expected to follow if the early case mortality be lessened, though there are probably other factors also in their causation. Improved methods of diagnosis, too, are adding to the cases found to be suffering with congenital syphilis, whether that disease be the cause of their symptoms or not. Personally I am awaiting with impatience the perfection of the cuti-reaction test for syphilis, and the placing on the market of Noguchi's reagent "luetin," at present only obtainable by favour. As the result of 134 trials made with it at the Babies Hospital, New York (reported by A. Brown¹), 3 unsuspected cases gave a skin reaction which was not confirmed by the Wassermann test, and 4 known syphilitics did not respond—all cases of severe infection; this is in line with our experience of the von Pirquet cuti-reaction for tuber-

culosis. In the remaining 127 the result obtained, whether positive or negative, corresponded with that given by the Wassermann method. In cases of late syphilis, in some of which the Wassermann reaction is obtained from the cerebro-spinal fluid but not from the blood serum, the luetin test is the more reliable. But in early cases I am told by Dr. Frederick Wilson, surgeon to the Lock Department of the Liverpool Royal Infirmary, to whom I am indebted for several valued hints incorporated in this paper, it is less so, and cannot therefore entirely supersede the Wassermann.

The original Wassermann reaction is so complicated and expensive, both in blood, time, and money, that we are unable to make use of it with anything like the frequency that is desirable. Where the necessity is pressing, with the mother's consent, her blood may be examined and the result taken as applying to her offspring. In certain cases this method has an advantage over that of examining the child's blood, for children of syphilitic parentage do not always give the reaction until, or just before, symptoms show themselves.² On the other hand, in some cases the newborn baby may give the reaction and yet never develop symptoms, having presumably absorbed reacting substances from the placental circulation without actual infection.³ In these cases, though the reaction would gradually die out in the child, it would be present in the mother. It has been found that some children of syphilitic origin presenting marasmic symptoms never give a positive Wassermann reaction; here a toxic influence upon the ovum must be held accountable. Such cases are not examples of true congenital syphilis; they are to be looked for when the disease in the parent is long-standing, for the more recent the syphilis in the parent the more obvious are its results upon the offspring. All the textbooks on children's diseases warn the practitioner to suspect inherited syphilis in marasmic babies who do not improve under the influence of rational feeding and hygienic surroundings, but I have seldom, if ever, obtained any beneficial result that I could trace to the exhibition of mercury in this class of case. I have been disappointed so often that I was particularly interested to read a few months ago an abridged paper by Dr. Emmett Holt of New York,⁴ in which he states that of 62 infants suffering with malnutrition only 5 gave a positive Wassermann reaction (Noguchi modification), and that these 5 are included in 11, of whom 7 were, upon fuller investigation, considered to show clinical signs of syphilis. Two San Francisco physicians⁵ report 8 triple positive Wassermann reactions in 100 marasmic infants without physical signs of syphilis, and although they claim from this that "many so-called cases of marasmus are due to latent lues," to my mind these and Holt's figures confirm the opinion I had formed for some time—that in congenital syphilis of the very young diagnostic signs, when looked for, are generally not wanting, and that if syphilis play any great part in the production of marasmus of otherwise unassignable origin it is by its toxic influence upon the ovum in the manner already indicated.

It is said that not more than 5 per cent., or at the most 10 per cent., of the conceptions of syphilitic mothers survive at the end of the first year of life. The slaughter of the innocents begins in the abortions and miscarriages which occur in the first years after infection of the progenitors. At times a series of miscarriages is followed by the birth of a monstrosity, such as was under my care at the Liverpool Infirmary for Children recently. In this infant, which lived for eleven days (Fig. 1), there was an absence of ossification in the bones of the calvarium, so that any pressure caused the brain to bulge in the opposite direction, and upon the vertex was a meningo-encephalocele as large as a Tangerine orange. There was a widely gaping hare-lip on the right side and complete cleft palate; the left eye was absent or represented externally only by a furrow in the skin, while the right eyelids were missing, and the eye appeared like a little black island in a sea of pink inflamed conjunctiva. The fingers and the toes of the left foot were webbed, while the right foot, nearly severed from the leg by a deep constriction, was a shapeless mass terminating in rudimentary digits.

Some cases like this being undoubtedly syphilitic and other deformities (such as hare-lip and congenital heart disease) having been shown by the Wassermann reaction

* Adapted from papers read before the Manchester Clinical Society and the Liverpool Medical Institution, January 20th and February 5th, 1914.

also to occur in congenital syphilis,⁶ there is a little tendency on the part of some to jump to the extreme of declaring that all cases of congenital malformation owe their origin to syphilis, a position which I think is untenable. Hochsinger⁷

Fig. 1.

found congenital heart disease only 7 times in 500 cases of (clinical) congenital syphilis, while Holt⁴ says that in 56 consecutive cases of congenital deformities he failed to get a single positive Wassermann. The usual syphilitic rash is not difficult to diagnose, but there are two rashes occurring in babies which are occasionally wrongly diagnosed as syphilitic. The first of these is the eruption known as "pemphigus neonatorum," which is not a true pemphigus, but the result of a streptococcal infection of the skin, and therefore allied to impetigo contagiosa. A bullous syphilide is sometimes found in newborn babes, though not very frequently. It is almost always present at birth, and though it may appear elsewhere, always affects the palms and soles, the skin of which rapidly assumes the typical raw ham hue. The pemphigoid eruption arises in all cases after birth; and, though it may affect the hands and feet, only does so incidentally, and as often as not the dorsal surface rather than the palmar. The syphilitic baby is sure to be wasted, and more than likely to die, this rash being symptomatic of a very severe infection; the other may be—very likely is—well nourished, and the lesions rapidly respond to local antiseptic treatment.

The other non-syphilitic rash I have in mind is the disease known as eczema seborrhoicum, which, like pemphigus neonatorum, is probably a skin infection (due to the *Staphylococcus epidermidis albus*). In the frequent form of a scaly, greasy, yellowish eruption upon the scalp and on the face—chiefly in the folds between the nose and lips, and behind the ears—there is little if anything in common between eczema seborrhoicum and a rash of congenital syphilis. But at times we are confronted by a case in which the thighs and abdomen—in fact, the whole of the napkin area—is covered by an extensive eruption from whose surface the scales have been removed by contact with the evacuant discharges and the napkin itself. There is left an inflamed, moist, oozing surface, red rather than yellowish, not perhaps so dusky as, but by no means easily distinguishable from, the more serious disease. Points in the diagnosis are again the state of nutrition of the infant, and the presence, or absence, of lesions on the hands or feet, and other evidence of syphilis; if the case be eczema seborrhoicum circumscribed patches of the rash will within a short time be found beyond the napkin area, and covered with the diagnostic scales.

In the January number of the *British Journal of Children's Diseases* two cases are epitomized from a German paper by Bosányi,⁸ in which Reynaud's disease, affecting the right hand and both feet in one and both hands and feet in the other, was a symptom of congenital syphilis, but the association of these two diseases is uncommon. The occurrence, as a sequel, of such exten-

sive gangrene as is seen in Fig. 2 must be very infrequent. This patient, a boy of 8 months, was under the care of Mr. R. W. Murray at the Liverpool Children's Infirmary. He had had previously frequent attacks of oedema of the feet, but otherwise was said to have been well till a fortnight before admission, when the "foot turned black." The gangrene was superficial, involving the skin of the right foot, except that of the sole, and there was a definite line of demarcation. The diagnosis of syphilis was confirmed by a bad attack of laryngitis, while traumatism was excluded, and frostbite, although the case occurred in January, was highly improbable. Under antisiphilitic treatment internally, and antiseptic fomentations locally, the child recovered with the loss of all its toes.

During the first few years after infancy the commonest symptoms are condylomata and laryngitis. The former are easily recognized and as easily cured. The laryngitis is chronic in type, with acute exacerbations, and in the absence of other indications of syphilis its nature is apt to be missed. Thus, as in the case of William M., aged 15 months, the result of treatment would sometimes seem to point to syphilis, although there be no other reason to suspect it. This child had been ill for a fortnight, and under treatment as an out-patient for ten days, with a laryngeal cough and hoarse voice and cry. The only recorded rise of temperature was 104.8° on admission. Steam inhalations and a mixture containing ipecacuanha and ammonium acetate effected little improvement, but upon a change to potassium iodide he made a rapid recovery. Potassium iodide, apart altogether from syphilis, is a very useful drug in early laryngeal and bronchial catarrh, and it is not claimed that in this particular case the result of the treatment is conclusive proof of the syphilitic origin of the laryngitis, but the possibility in such a case is always worth bearing in mind.

In other cases, which are distinctly rare in infancy, but perhaps commoner towards puberty, the laryngitis proceeds to ulceration of the vocal cords. In a case under the care of the late Dr. Carpenter⁹ this was associated, in a female child aged 13 months, with interstitial pneumonia—white hepatization—another of the manifestations of syphilis liable to escape diagnosis or to be confounded with the ordinary bronchopneumonia which so often terminates the existence of weakly infants. It generally extends from the apex downwards, and if truly syphilitic is amenable to treatment. It is, however, difficult to be certain in any given case whether the consolidation is syphilitic, and not either tuberculous or due

Fig. 2.

to delayed resolution of a pneumococcal pneumonia in a syphilitic subject. The older the child the more likely is it to be the latter. I read in the *BRITISH MEDICAL JOURNAL*¹⁰ recently that in Glasgow the health authorities now supply the general practitioner with tubes for the collection of blood and perform Wassermann reactions free for him. When other towns have followed this example we shall have all these obscure cases accurately diagnosed before they reach the hospitals.

An American physician recently reported a case of paroxysmal haemoglobinuria in a girl,¹¹ whose first attacks began at 2 years. He found a positive Wassermann reaction in this patient, and collected sixteen others occurring in congenital syphilis, as well as a further similar number in acquired syphilis. Congenital syphilis must also be held responsible for a

specific nephritis—interstitial and parenchymatous, either combined or separately. The first case to be recognized clinically and treated successfully was a patient of a Manchester physician, Dr. S. Messenger Bradley, and was recorded in 1871.¹² This is one of the earlier manifestations, and is to be looked for in the first few months, or even weeks, the lesion in fact being sometimes present at birth. Oedema is generally, but not always, a prominent symptom, and probably a greater number of cases without oedema would be found were it not for the difficulty in obtaining specimens of the urine from infants. If the case come to *post-mortem* examination it is important to remember that, as Carpenter pointed out,¹³ the naked-eye appearances of the kidneys are frequently normal, and sections must be cut before they can be declared free from disease. In some instances an infantile syphilitic nephritis may have been the starting-point of some of the chronic cases encountered later in life; but it would be difficult, if not impossible, to prove the connexion. Its frequent presence is a factor to be considered in salvarsan therapy.

Dr. Dean¹¹ classifies the cases in which congenital syphilis is the cause of disease of the nervous system into three groups:

1. Infants born with marked evidence of syphilis. In these cases the brain, together with other organs of the body, is the seat of marked changes. These children as a rule die early.

2. Children who appear to be healthy at birth but develop mental defects at the time of the second dentition or puberty. To this class belong the cases of general paralysis.

3. Those cases of mental deficiency, imbecility, and idiocy in which the Wassermann reaction constitutes the sole evidence of syphilitic origin.

I shall say nothing about the last class, except to note the extraordinary differences in the findings of various observers, which range from 0 per cent. to 100 per cent. of the cases examined. On the whole the inference seems incontestable, especially as a positive reaction is always found in greater percentage in the earlier years, that inherited syphilis plays a much greater part in the production of mental degeneration than was supposed. I think it may also be presumed to have some bearing on the fact, revealed by the statistics of the 1911 census, that the largest increase of insanity is among children under 15 years of age.

I shall only offer one observation, too, on the first class, those born with lesions such as hydrocephalus and gross cerebral deformities, who as a rule die early. It is to remark that whatever is meant by "early," these children do not invariably die in the first few weeks of life. Some of the cases of spastic diplegia which bring chronic grief to the parents, grist to the orthopaedic surgeon's mill, and kudos to nobody, are of ante-natal and syphilitic origin. They are always accompanied by mental defect, often of severe grade, and by epileptic fits.

Dr. Dean's classification does not appear to me quite comprehensive, as it makes no preparation for those—an important and, I think, increasing section—who owing to blood vascular changes develop meningitis, secondary hydrocephalus, and plegias (in other words, "cerebral syphilis") at any time during childhood.

The earliest of these symptoms to present itself is meningitis. This is generally basal and the clinical picture that of posterior basic (or meningococcal) meningitis. As the treatment of the two is very different it is essential that a lumbar puncture should be made early and the cerebro-spinal fluid examined by a competent pathologist. If the meningitis be syphilitic the fluid will be clear, or at any rate less turbid if unmixed with blood than in meningococcal meningitis; it will contain an excess of lymphocytes instead of polymorphonuclears; be sterile instead of giving a culture growth on nasgar; and, though spirochaetes may possibly be found with luck, the characteristic intracellular diplococcus of Weichselbaum will not be seen. A good deal is being written just now about the tests for and presence of globulin in the syphilitic fluid, and the precipitation of colloidal gold¹⁵ (Lange's test). If the fluid does not give the Wassermann reaction the blood may. The immediate prognosis under treatment is better than that of meningococcal meningitis, but blindness, deafness, and mental impairment are likely to be more permanent in the event of recovery. The following is a typical case:

Edith C., aged 3 months, under my care at the Liverpool Children's Hospital from April 8th to June 1st, 1910. The

mother gave a history of two stillbirths and one child dead after convulsions. The patient weighed 4 lb. at birth and was breast fed for the first month; afterwards fed on milk and water. Has been vomiting and constipated, and seems ill since a fall last night.

Condition on Admission.—Puny infant. Pulse 104, regular; temperature 99.2°; no rash; chest and abdomen normal. Slight cervical opisthotonos. Fontanelle normal tension. Kernig's sign present. Knee-jerks obtainable and equal. Superficial reflexes normal. Pupils equal and reacting to light.

Slight or even well-marked cervical rigidity is very common in gastro-intestinal troubles in infants, but the next day this symptom was increasing, and the day after, although the vomiting on a suitably adjusted milk diet had decreased, was marked. So I performed lumbar puncture, and removed 40 c.cm. of slightly turbid fluid, and injected 15 c.cm. of Flexner's serum. On examination of the fluid, however, it was found to reduce Fehling, to show lymphocytosis, and gave a negative cultural reaction. At the time this was taken to represent a meningococcal meningitis in the stage of recovery. The symptoms abated after the operation, but the child lost weight. On April 15th the cervical rigidity began to increase again, and a copper-coloured rash made its appearance on the buttocks and thighs, circinate on the latter; while raw fissures were noted about the angle of the mouth. One of the papules on the thigh was scraped, and a *Treponema pallidum* found in the serum which exuded. Lumbar puncture was also done again, but none discovered in the cerebro-spinal fluid, which was similar to the first specimen withdrawn. By April 18th there was more rash, on the nose and face. Up till May 1st the child slowly lost weight, although the meningeal symptoms improved, and between that date and its discharge all symptoms, both of meningitis and of syphilis, disappeared, except that the child was left with nystagmus and apparently blind. I could not see its discs. It was, of course, treated with mercury from the time of the appearance of the syphilitic rash, in increasing doses till the symptoms began to ameliorate.

Apart from this form of basal meningitis, I believe syphilitic pachymeningitis to be less common than in adults with the acquired disease.

Writing of inherited syphilis, W. P. Lucas, of Boston, says:¹⁶

That the syphilitic virus often attacks with unusual severity the heart and blood vessels is only lately being fully recognized. The arteries of the arch of Willis are most often attacked of the small arteries. An acute gummatous endarteritis of some of the large arteries in the brain . . . is now positively and not infrequently recognized; obliterative endarteritis of the small blood vessels of the brain . . . is met with.

It is evident that we have here all that is necessary for the production of attacks of hemiplegia, such as are so frequent in acquired syphilis, and when hemiplegia is met with in children, apart from anything that may be gleaned from the previous or family history, the following features will be found to indicate a syphilitic origin:¹⁷

1. The hemiplegia is often preceded by convulsions at a longer or shorter interval before the attack.

2. It is often recurrent—one or more attacks following the first, after each of which the prognosis of functional recovery decreases.

3. Headache and vomiting may lead up to the paralytic seizure.

4. According to Box, the silent development of acute hemiplegia is very suspicious.

In using these axioms as a guide to diagnosis we must first eliminate other causes of hemiplegia. For instance, convulsions in a non-syphilitic infant may occasionally cause cortical haemorrhage, with subsequent compression hemiplegia. Recurrent attacks may occur from embolism in infective endocarditis, while headache and vomiting are the commonest prodromal symptoms of tuberculous meningitis, in which hemiplegia may develop, especially in association with a tuberculous cerebral tumour. Let me quote two illustrative cases, not of the exceptions, but of the rule:

X., a girl aged 11, seen in consultation with Dr. Kidd in Birkenhead. No definite syphilitic history obtainable, but there had been one miscarriage and several of the dozen brothers and sisters had died with "wasting" in their first year. Five months ago she had vomited, gone into convulsions, and the next day had right hemiplegia with aphasia. From this a good recovery was being made, and speech was being learnt. The complete use of the lower limb had been obtained, but the upper remained spastic. Fourteen days before I saw her she began to vomit again, complained of headache, and soon developed the typical clinical picture of meningitis. She had optic neuritis; no tubercles in the choroid could be seen. The pathologist's report on the fluid obtained by lumbar puncture read "Alb. +, Fehling very deficient, enormous excess of lymphocytes, no organisms seen." The child was far gone when I saw her, and died a few days later. I regarded the case as one of syphilitic endarteritis with

thrombosis and sclerosis, followed by syphilitic meningitis, but in the absence of a *post-mortem* examination I admit it is at least as likely that the terminal infection was tuberculous.

The second case is under my care in the hospital at the present moment.

Z., a boy aged 6. His mother is under treatment at the Royal Infirmary for tertiary syphilis of the nose. On August 6th, 1913, he dropped his cup and bread from his left hand at breakfast, and within the next few hours weakness was noticed in the left side of his face and dragging of the left leg on walking, while speech was distinctly affected. (Here we have the "silent onset.")

On admission, pulse, temperature, and respiration are normal. Intelligence poor; lies quietly, takes no notice, wets bed, cannot read letters nor answer questions intelligibly, and speech is indistinct. Can stand, but is weak on left leg and drags it after him. Left upper extremity also weak, but degree hard to estimate owing to his lack of intelligence. Left face, including the eyelid, parietic. Tendon reflexes increased on affected side, superficial normal. No wasting. Pupils equal and dilated, react sluggishly. Fundus: Right, normal; left, extensive patch of choroidal atrophy with interspersed pigment; disc = white atrophy. Vision nil.

Under treatment with mercury and iodide he improved rapidly, so that by September 1st, when he was discharged, he was running about the ward, with intelligence normal and no paralysis left, except in the face. On November 10th, in out-patient's department, I noted that there was only the slightest facial weakness left, but on December 31st he was readmitted on account of a recurrence of hemiplegia on the same side, preceded this time by severe headaches. The second attack is accompanied by much spasticity of the extremities—clasp-knife rigidity of the knee, adduction of the thumb, flexion at the hip, knee, wrist, and elbow, etc., but there is no fresh involvement of the face, nor is his mentality again impaired. He has recovered up to a certain point, but splints are now being used to prevent spastic contractures.

With evidence of such extensive and acute cerebral syphilis there is great risk of a catastrophic local reaction were he treated with salvarsan, and so far he has not had it. But I intend to administer either that or neo-salvarsan after a thorough course of mercury and iodide, and hope thereby to stop the disease at its present stage, and before it reaches that illustrated in the following case. I think this patient should be included here as an example of the pathetic termination of cerebral syphilis, though some might say it was juvenile general paralysis.

A boy, aged 11, sent to me by Dr. Sheridan, and admitted to the Liverpool Infirmary for Children on February 7th, 1911. Father admitted syphilis. Mother died with phthisis two years after the boy's birth. The child is said to have been quite well and bright up to 7 years of age. He then had some sort of left hemiplegic attack, without convulsions, upon which progressive weakness ensued. For the last twelve months he has been quite helpless, having lost the power of active movement and control over his bladder and rectum. The power of speech has been lost for about four months, and of late he has been going into "swoons," lasting as long as twelve hours, in which "the face twitches and the eyes roll."

Upon admission, temperature and respiration are normal; pulse accelerated, about 120. Upper central incisors notched. There is general rigidity, and the patient lies huddled up in bed and is unable to sit up. Expression usually vacant. Intelligence grossly impaired. He will sometimes answer "yes" and "no" to simple questions; generally no answer is obtained. When asked his name he tries to say some word, but it is quite unintelligible; there seems to be difficulty in articulation. Faeces and urine passed into the bed. There is convergent strabismus. Pupils unequal, dilated and inactive; but Mr. Shears reported the fundus practically normal and the squint alternating and due to hypermetropia. His neuro-muscular condition may be summed up by saying that he was diplegic, with extreme spasticity and exaggeration of reflexes.

A week after admission the patient had a convulsive seizure during the night, lasting an hour; movements general, but more severe on the left side of the body. This was succeeded by a state of general tremor with clammy skin, frequent pulse and shallow respiration, becoming Cheyne-Stokes before he died towards the close of the third day after the epileptiform attack. No *post-mortem* examination could be obtained.

In his condition, when I first saw the boy, the diagnosis between cerebral syphilis and general paralysis was unimportant, for recovery was impossible; but in the early stages it is momentous. For whereas the former may be more or less amenable to treatment, dementia paralytica is, or used to be considered, parasymphilitic, and is, at any rate as yet, upon the list of incurable and progressively fatal diseases.

Similarly, locomotor ataxy, though much rarer than general paralysis in children, is incurable, though sometimes subject to arrest. Remak,¹⁸ in 1885, first called attention to this disorder, and a fair number of cases is now on record. Many of them are brought to light by the

oculist,¹⁹ for optic atrophy is unusually frequent, while the ataxy is not marked. It has occurred at any age between 3 and adult life, but the majority seem to develop the disease at about 15. Enuresis is usually the first symptom. Unlike general paralysis, tabes is more frequent in females—in the proportion of 63 per cent. to 37 per cent.²⁰—though, as the known cases are comparatively few, this disproportion may be only coincidence. It is to be noted that locomotor ataxy in children has followed syphilis which has been accidentally acquired, not inherited, with a much greater frequency than is proportionate to the incidence of the two varieties. I have never recognized a case myself.

The discovery of the spirochaete and of the Wassermann and other diagnostic tests has thrown the whole subject of syphilis into the melting-pot, from which new conceptions of the disease are emerging. These are still malleable, and fresh investigations are daily entailing their remodelling. For instance, as was indicated above, it is more than likely that the term "parasymphilitic" may have to be dropped, the discovery of the spirochaete in the cerebral tissues of general paralytics compelling the inference that this disease and those related to it will now fall into place as pure syphilitic infections of the nervous system. Mott²¹ suggests the term "parenchymatous syphilis."

REFERENCES.

- ¹ Brown, *Amer. Journ. Dis. Child.*, 1913, vi, p. 171. ² Mulzer and Michaelis, *Berl. Klin. Woch.*, 1910, xlvii, No. 30, p. 1403. ³ Boas and Thomsen, *Arch. v. Derm. u. Syph.*, 1912, cxi, p. 91. ⁴ Holt, *Amer. Journ. Obstet.*, 1913, lxxvii, p. 1228. ⁵ Yerington and Holsclaw, *Amer. Journ. Dis. Child.*, 1914, vii, p. 32. ⁶ Churchill, *Amer. Journ. Dis. Child.*, 1912, iii, p. 368. ⁷ Carpenter, Wightman Lecture, 1909. ⁸ Bosanyi, *Jahrb. f. Kinderheilk.*, lxxviii, p. 177. ⁹ Carpenter, *Brit. Journ. Dis. Child.*, v, p. 161. ¹⁰ Browning, *BRITISH MEDICAL JOURNAL*, 1914, i, p. 81. ¹¹ Ager, *Amer. Journ. Dis. Child.*, 1912, iv, p. 311. ¹² Bradley, *BRITISH MEDICAL JOURNAL*, February 4th, 1871. ¹³ Carpenter, *Reports of the Society for the Study of Diseases in Children*, 1907, vii, p. 277. ¹⁴ Dean, *Brit. Journ. Dis. Child.*, ix, p. 392. ¹⁵ Grulee and Moody, *Journ. Amer. Med. Assoc.*, 1913, lxi, p. 13. ¹⁶ Lucas, *Amer. Journ. Dis. Child.*, 1912, iii, p. 264. ¹⁷ Box, *BRITISH MEDICAL JOURNAL*, 1911, i, p. 982. ¹⁸ Remak, *Berl. Klin. Woch.*, 1885, No. 7. ¹⁹ Stephenson, *Lancet* May 16th, 1908. ²⁰ Price and Shannon, *Amer. Journ. Dis. Child.*, 1912, iii, p. 236. ²¹ Mott, *BRITISH MEDICAL JOURNAL*, 1913, ii, p. 1270.

THE CLINICAL IMPORTANCE OF THE CEREBRO-SPINAL FLUID.*

BY

WILLIAM BOYD, M.D., M.R.C.P.E., DIPL. PSYCH.,
PATHOLOGIST TO THE GENERAL HOSPITAL, WOLVERHAMPTON.

THE cerebro-spinal fluid, when withdrawn, is clear, colourless, and limpid, with a faintly alkaline reaction, and a specific gravity of 1006 to 1008. It contains a trace of globulin, and a substance which reduces Fehling's solution, probably glucose. It contains no polymorphs, and only an occasional lymphocyte.

NAKED-EYE CHANGES.

The fluid may be turbid instead of clear. Excluding contamination by blood at the time of puncture, turbidity indicates the presence of polymorphs, which are never present in the normal fluid; their presence as a rule indicates acute inflammation of the meninges. Enormous numbers of lymphocytes, on the other hand, may be present without producing any turbidity. This is a point of practical importance, for in tuberculous meningitis, where the cells are mainly mononuclears, the clearness of the fluid might lead to the conclusion that there was no meningitis, but this microscopic examination would at once prove to be erroneous. The fluid may also contain large numbers of micro-organisms, and yet appear perfectly clear. Turbidity, then, is an indication of acute meningitis. A cerebral abscess, unless it involves the meninges or drains into the ventricles, produces no change in the fluid. Thus, if a patient shows signs of acute cerebral trouble accompanied by a high leucocytosis in the blood but a normal cerebro-spinal fluid, the probability is that he is suffering from abscess of the brain.

CYTOLOGY.

The number and nature of the cells present are of great importance. Using Widal's centrifuge method and a

* Abstract of a paper read at the Staffordshire Branch of the British Medical Association.