An Address ON WOUND INFECTIONS; AND ON SOME NEW METHODS FOR THE STUDY OF THE VARIOUS FACTORS WHICH COME INTO CON- SIDERATION IN THEIR TREATMENT. DELIVERED BEFORE THE ROYAL SOCIETY OF MEDICINE ON MARCH 30TH.

By COLONEL SIR A. E. WRIGHT, M.D., F.R.S.,
A CONSULTING PHYSICIAN TO THE EXPEDITIONARY FORCE.
(From the Research Laboratory attached to No. 13 General
Hospital, Boulogne-sur-Mer.)

THERE are a number of quite elementary problems which must be solved before we can arrive at any really effective treatment of wound infections. A very brief consideration of the facts will bring us face to face with the questions to which we have to find an answer.

In this war practically every wound is heavily infected. The chain of cause and consequence seems to be as follows. The clothes and skin of the soldier on war service become contaminated with all manner of filth containing pathogenic organisms and spores; the projectile takes these in with it, and it implants them far beyond the reach of any prophylactic applications of antiseptics.

A cultivation medium is now provided by the blood and lymph which are poured out into the track of the projectile; and we find in the wounds—I have in view here wounds examined immediately after arrival from the front—a mixed infection of a streptococcus with microbes derived from the faeces. This faecal infection is a special outstanding feature in this war.

Among many species of intestinal microbes which have been found in wounds, two have a quite special importance. One is the *gas-phlegmon bacillus*, or *Bacillus aerogenes capsulatus* of Welch—a large Gram-staining anaërobic and actively gas-forming microbe. It is found both in infiltrated superficial wounds, and in deep wounds, and is particularly abundant in the frothy and offensive faecal-looking discharges which anaërobic wounds furnish. The other is the tetanus bacillus. It is more rarely encountered, and is also much less abundant in the discharges. Sometimes, however, it may show up in every field of the microscope.

The presence of the streptococcus and these two faecal microbes makes the first period of the wound infection—the period of imprisoned discharges—a specially critical time for the patient. During this the streptococcus may invade the tissues, and set up cellulitis or, more rarely, erysipelas. Or the tetanus bacillus may find opportunity to grow out and manufacture its poison and induce tetanus. Or the bacillus of Welch may make its way into the body and set up a gas-phlegmon in the region round about the wound, and an obstructive gangrene in the distal portion of the limb. Or, again, the bacillus of Welch and the streptococcus may join forces, and may in conjunction produce the gas-phlegmon or cellulitis.

As soon as a free outlet has been provided, and aerobic conditions have been established in the wound, its bacterial flora changes. The ordinary pyogenic infection, which has up to this been in abeyance, now gains the upper hand, and instead of an “infection of the imprisoned discharges,” or, as the case may be, an infection of tissues, we have now an “infection of the granulating wound surfaces, and of the flowing discharges.” The chief bacterial agents here at work are the streptococcus and staphylococcus, and *Bacillus proteus*.

This pyogenic infection may, after lapse of time, subside, the wound healing up when this occurs; or the mixed infection may narrow itself down to a streptococcal infection and become chronic, the wound in this case remaining open indefinitely in the form of a discharging sinus. Or, lastly, when there is an obstructed outflow, the infection may go from bad to worse until the patient succumbs to continued suppuration and septicaemia.

Those are in very brief summary the facts with regard to the evolution of wound infections, and I would venture, in passing on to discuss with you their treatment, to remind you that the ideal we ought to approximate to

is the healing of the wounds by *first intention*—that is, without sensible interference by bacterial infection; and that we are so far from the attainment of that ideal that nearly all our wounded are suffering from bacterial infections; that very many are ill of these infections; and that not a few are, through these, in danger of their lives.

We have at disposal for the treatment of these wound infections three different therapeutic measures. Let me enumerate them to you in the order in which they would naturally suggest themselves to you.

First in that serial order would come *treatment by antiseptics*. After this would come what I would propose to call *treatment by physiological methods*—I mean procedures such as the opening and draining of the wound—which bring the antibacterial powers of the blood to bear on the infecting microbes. And lastly would come the reinforcement of the antibacterial powers of the blood, that is, *treatment by vaccine therapy* and similar methods. I believe it is really beyond all question that of these three the second is beyond all comparison the most important, and I would submit that, all loud talk about antiseptic treatment notwithstanding, this is at best an ancillary method of treatment. And of course the same applies also to treatment by vaccines.

Let me also here suggest to you another quite fundamental consideration. It is this: It will be clear that we cannot apply physiological treatment aright, nor can we apply any antiseptic or vaccine to best advantage in wound infections unless we first understand the physiological processes which go on in the wound. We do not understand these yet even in outline.

It will therefore be necessary to address ourselves to the task of discovering what goes on in the wound and of following out its biological evolution. And the only way of doing this will be to formulate to ourselves in clear terms the questions which want answers; then to consider how to set to work to get our answers—for merely looking at the wound will not help—and finally to take cognizance of the results which the experimental methods which I am about to describe to you have already yielded.

Our first question can be formulated thus:

1. *Can the microbes which are found in wound infections live and multiply in the unaltered blood fluids?*

In other words, if I take pyogenic microbes from the wound and implant these into the normal undiluted serum, will this give me a satisfactory amount of growth? If we are going to carry out this experiment, and to carry it out repeatedly, and deal with a number of different bloods; and if we are going to cultivate directly from the pus, we shall evidently have to work in capillary tubes with very minute quantities of pus and very small quantities of undiluted serum. The technique which I have arrived at for fulfilling these requirements is a very simple one. I may call that technique the *wet-wall method*.

Method of Making Cultivations of Pus in Serum.

The first step in the procedure is to make, by the technique I described in my book on *Technique*,¹ a graduated series of dilutions of pus, using for this purpose any indifferent diluting fluid, and arranging the successive dilutions on a slide in the form of a series of drops (Fig. 1, Slide I). I then take in hand a clean capillary pipette

Slide I.

Slide II.

Fig. 1.—Slide I: Graduated dilutions of pus arranged on slide. Slide II: Volumes of serum implanted with the above dilutions of pus.

fitted with a teat, make a mark upon the stem, and then draw up into it a series of unit volumes of serum, one unit volume of serum for every dilution of the pus. This done, I commence with the highest of these dilutions—that is, the one containing the smallest

number of microbes—and draw it up into the stem of the pipette, stopping off exactly at the fiducial mark (Fig. 2). I then expel this column, leaving, of course, as I do so, the walls wet with a quantum of microbial suspension—a quantum which would correspond roughly to that which would be left on the outside of a platinum wire of similar stoutness dipped into the suspension. I now expel from my capillary tube my first volume of serum—this in passing over the wet wall will take up its charge of microbes—and I receive this as it issues on to a clean slide. I repeat these manipulations with the next lower dilution of pus and the next unit volume of serum, and so on, until I have implanted my series of volumes of serum with my series of bacterial dilutions. At the end I find myself with a series of drops of serum containing graduated charges of microbes, ranged in order upon a slide (Fig. 1, Slide II); and I have in my hand a pipette which is contaminated up to the point indicated by my fiducial mark. I get rid of this contaminated segment of my capillary stem by resecting this just above the fiducial mark, and I now proceed to draw up into my pipette, separating off by bubbles of air, the whole series of drops of serum, beginning with that implanted with the fewest microbes. After sealing up the distal end of the pipette, and closing the butt end by bringing it down upon a bed of plasticine, the pipette is placed in the incubator. After an interval of six to twenty-four hours it is taken out, and the series of unit volumes are now, for purposes of microscopic examination or culture, expelled from the pipette in the order in which they have been taken up—that is, in succession from the lightest to the heaviest implanted. This is effected by cutting through the proximal end of the stem of the pipette and then the distal end, and afterwards luting the latter air-tight into the truncated end of the pipette. The luting is done by rolling a little plasticine into a ball, pressing the capillary stem into this pellet as we hold it between finger and thumb, doubling over the plasticine round the stem, and then invaginating the neck of the pipette into the plastic mass. We can now with a teat expel our series of unit volumes of serum on to a slide, obtaining them in the serial order which avoids the contamination of the lighter implanted with the heavier implanted.

Fig. 2.—Capillary pipette filled with five unit volumes of serum, and one unit volume of the last dilution of pus.

Method of Making Anaërobic Cultures in Capillary Tubes.

When we desire to make anaërobic serum cultures we do not employ the capillary pipette used for the original

Fig. 3.—Method for making anaërobic cultures in capillary tubes. implanting operations as a receptacle for our series of serum cultures, but employ instead a separate capillary

cultivation tube for each unit volume of serum. The capillary cultivation tubes here in question are made as follows: We take a portion of capillary stem, say some 8 cm. long, draw it out at one end in the flame of a by-pass into a hair-fine tube, and then break it off, leaving a certain length of throttle attached. We now introduce the other end of the capillary stem into the flame of our Bunsen, and let gravity bend it round into a siphoning curve.

Taking one of these capillary cultivation tubes we bring down the siphon end upon our drop of serum (Fig. 3, a), and let this run in, afterwards tilting the tube so as to take in, as a rear guard, a good sized dividing bubble of air. This done, we fill into our tube, by siphonage, from watch-glasses placed conveniently to hand, first a small quantity of pyrogallic acid, and then, without dividing off, a small quantity of caustic soda (Fig. 3, b and c). We now further incline the tube, and let the contained fluids gravitate down towards the distal end until the serum begins to enter the throttle. At that point we arrest it by sealing the tip, and we finally also seal up the butt end, leave the mixed pyrogallic acid and caustic soda to do their work and absorb all the oxygen from the dividing bubble air (Fig. 3, d).

Serum cultures made by this method furnish a very striking and uniform result. We obtain in the cultures implanted with the higher dilutions of the pus a pure culture of the streptococcus, and in the cultures more heavily implanted with the pus the streptococcus mixed with a certain number of other microbes; in particular, a few staphylococci and an anaërobic wisp-like diphtheroid bacillus which often is abundant in pus, being found both intracellularly and extracellularly. All the other pyogenic microbes appear to be inhibited in undiluted normal serum, and when they put in an appearance it is only after fairly heavy sowings with pus, and comparatively late.

Out of these facts would come what we shall presently see to be a practically important classification of pyogenic microbes—a classification into, on the one hand, *sero-phytes*, and, on the other, *sero-saprophytes*. The sero-phytes would be those which, presumably because they find their foodstuffs ready made in the blood fluids, are at home there, and can, in the absence of phagocytes, grow and multiply there without restraint, or practically without restraint. The sero-saprophytes would be those which cannot grow and multiply in the blood fluids until a change, which we may, pending nearer investigation, call simply a degenerative change, has passed over those fluids.

What holds true of the blood fluids themselves might perhaps justifiably be assumed to hold true also of the lymph which pours into the wound. None the less, it will be well specially to investigate this point.

2. Does the lymph which pours into the wound provide a favourable nutrient medium for the microbes which have been growing in that wound?

To pose this question is already to go a long way towards getting an answer to it. For we can, by the aid of a very simple device, obtain the lymph from the walls of the wound. I employ for this purpose what I may perhaps call a *lymph leech*. This consists, as you see, of a small glass tube. It is sealed up at one end, drawn out at the other in the form of a nozzle, and is furnished with a lateral mouth with a raised rim, the whole being very easily made out of a piece of glass tubing or small test-tube. To the nozzle we fit a piece of fairly thick walled rubber tubing, and this is blocked at the end with a piece of glass rod. When we want to obtain the exudate from a wound we bring down the lateral opening of the lymph leech upon a granulating surface, and then, transfixing the rubber tube with the needle of a hypodermic syringe, we draw out the air and make a negative pressure (Fig. 4). It will be appreciated that the lymph leech is in principle merely a small cupping glass, and that it will, by means of the vacuum we establish in it, hold on tight for whatever period may intervene between dressing and dressing, and furnish an exudate free from all contamination with residual pus left behind in the wound.

* I introduce this qualifying clause because I have on several occasions found the serum of the infected patient to give cultures of streptococcus with a planting of his pus much smaller than that which was required to give a culture in the serum of a normal man.

We can now proceed to compare the fluid in the cavity of the leech with the fluid in the wound outside, or, in connexion with work on antiseptics, to compare the contents of leeches applied respectively to treated and untreated surfaces in the same wound.

Fig. 4.—Lymph leech.

And lymph leeches also can be put to other uses. We can introduce for this purpose with a syringe any fluid we may select into the lymph leech, and investigate the effect that fluid exercises upon transudation and emigration. Again, where the nature of a deep-lying infection of a mucous or other surface remains uncertain, the application of a lymph leech might clear up the difficulty. We can also, in the case where we are testing the effect of a vaccine upon a wound, take to our aid the lymph leech. For the effect of a vaccine would probably show itself in the exudate in the cavity of the leech long before it would manifest itself in the outside wound. And, lastly, the application of a lymph leech to the site of inoculation might, perhaps, help to resolve the problem as to whether or no protective substances are then developed.

But it will be realized that for us at the moment that which is of chief interest is the comparison of the fluid in the cavity of the leech with the fluid in the wound outside.

When, after washing out a heavily infected wound with an antiseptic or simple saline solution, we apply a lymph leech to the walls, and then at the next dressing compare the contents of the lymph leech with the fluid outside, we think at first that there must be some mistake. Outside we have an opaque exudate presenting all the ordinary physical characters of pus containing leucocytes in all stages of degeneration, and swarming with all manner of pyogenic organisms. Inside we have a transparent and slightly blood-stained exudate containing streptococci in practically pure culture, and in addition a few leucocytes, all of which are actively phagocytic. Except in this latter respect, we have, in fact, identically the same result as when we made our thin implantations of pus into normal serum.

The problem now stares us in the face—What is it that makes all this difference between the contents of the lymph leech and the contents of the wound? How has the lymph which gives in the cavity of the lymph leech only a culture of streptococcus been converted in the wound outside into a fluid which is ideally favourable to the growth of a great number of different species of micro-organisms?

3. *What is the cause of that "corruption of the lymph" which converts it into a favourable nutrient medium for sero-saprophytic microbes?*

The proper way to go to work upon a problem of this kind is to keep it unremittingly before the mind. For then some hypothesis will in the end suggest itself. That found, there will invariably come to notice a certain number of accepted data which the hypothesis will fail to explain. These must then be carefully re-examined to see whether they will stand fast and discredit the hypothesis, or whether they also will come in and support it. And, finally, before launching our theory, we ought to think out all its consequences, and then take each of these and verify it.

At any other time one would wish, before promulgating a far-reaching hypothesis, to have verified it at every point. But we have on our hands in Europe, I suppose, already two or more millions of wound infections, and every other consideration must give way to that of accelerating

the researches, which we must look to to guide us in treating these infections.

I will therefore, before completing its verification, venture to put before you what is, I believe, the solution of the problem of the corruption of the lymph in the wound.

Let me take my departure from those facts which we were considering a moment ago, relating to the culture of pyogenic microbes in the blood fluids. You will remember that I suggested in connexion with serophytic microbes that these must find all the food materials they require ready formed in the blood fluids, and in connexion with sero-saprophytic microbes that for these no nutrient substances would be available until the albuminous substances of the blood had undergone some sort of preparatory transformation.

I conceive that that transformation could come only by a digestive process. Now, supposing this to be so, there would in the serum come into account a counter-acting influence. For we have there an antifermentative, or, as we usually style it, an antitryptic element, which would directly counteract any digestive element which might be struggling to come into operation. And it is clear that microbes which were dependent for their sustenance upon the products of digestive action could establish themselves in the blood fluids only on condition that this antitryptic influence was overborne.

This hypothesis furnishes, as it seems to me, an explanation for certain striking facts relating to the cultivation of microbes on blood fluids, and to bacterial infections. It, as it seems to me, explains the finding that heavy sowings of microbes into serum are effective in giving cultures, while light sowings are not only ineffective, but lethal to the implanted microbes. For it would be only natural that the resistance offered by the antitryptic power of the blood should be overcome by the mass effect of a number of microbes operating upon a restricted or—and this would come into consideration in localized infections—a mechanically isolated quantum of blood. And, again, it would be only natural that the mass effect of a large volume of antitryptic serum could not be overcome by the digestive activities of a very few microbes; and also, I think, in accordance with what we know, that microbes deprived of access to foodstuffs should perish of inanition.

Our hypothesis would also make intelligible, in connexion with infections by sero-saprophytic microbes, that frequent and heavy sowings into the blood should be required before a septicaemia can supervene upon a local infection. And again our hypothesis makes it comprehensible that there should be serious difficulty in obtaining haemocultures, even when the microbes have gained a footing in the blood stream.

And, lastly, our theory brings home to us that, in considering the defence of the body against bacterial infection, we have to take into account not only *active defence* in the form of phagocytes and bacteriotropic substances which make a direct attack upon micro-organisms, but also *passive defence*—that is, protection against infection obtained by preventing microbes converting to their uses the nutrient substances of the blood fluids.

These, however, are general considerations with applications far beyond the sphere of wound infections, and we must return to the particular problem of the corruption of the lymph in the wound.

When, after treating a wound with antiseptics and leaving it clean, we find it a very few hours afterwards teeming with microbes, we are in presence of something which, in my view, urgently stands in need of explanation. For our findings both in the lymph leech and in serum cultures made with the wet wall method from pus, would seem to teach us that the sowing of microbes left behind cannot be nearly heavy enough to produce the voluminous culture found in the wound, nor yet to account for the rapidity with which the sero-saprophytes have started to grow. And, moreover, consideration will bring out that another powerful factor must constantly come into operation in the wound. The factor in question is the tryptic ferment which is elaborated in the phagocytes and which is, when these break up, discharged into the surrounding medium. This tryptic ferment will come into operation under two different conditions. It will come in whenever a residue of pus—for this would contain free trypsin—is left behind in, or afterwards makes its way into, the

wound. Trypsin will again come into account whenever, after the washing of the wound, leucocytes once more begin to emigrate, and phagocytose, and break down.

I see in the reduction of the antitryptic power of the lymph thus effected the prime cause of its corruption. Before attempting to obtain confirmation of this from crucial experiments, there was a set of findings to be cleared up. I had found in connexion with pus implantations made into serum that when this was heated to 60° C. for ten minutes one no longer obtained the same differential growth of streptococcus as with unheated serum, but obtained instead mixed cultures of streptococcus with sero-saprophytes, in particular staphylococcus and the wisp-like diphtheroid bacillus already made mention of. It seemed at first sight as if this could not possibly be related with a reduction in the antitryptic power of the serum, for it has, in view of the high quality of Opie's work on this question, been generally accepted from him that the antitryptic power of the serum is unaffected by exposure to heat until a temperature approaching the coagulation point of serum is reached. In reality, however, when this is reinvestigated quantitatively,* it emerges that the antitryptic power of the serum is reduced by one-third to one-half when we subject the serum to a temperature of 60° C. for ten minutes.

That difficulty having been removed out of the way of our hypothesis, we may proceed to take cognizance of the results of the crucial test experiments—experiments in which graduated additions of trypsin are made to serum as a preliminary to the implantation of sero-saprophytic microbes. The event of these experiments can be summed up in a sentence. When we add trypsin in quantities sufficient to reduce appreciably the antitryptic power but insufficient to give us any free trypsin, the serum is converted into an eminently favourable nutrient medium for sero-saprophytic microbes.

Our hypothesis is thus very strikingly confirmed. It will be necessary hereafter to follow it into all its consequences. For the present it will, however, suffice if we ponder on the following points. The antitryptic power of the blood would appear to be increased in every case of severe wound infection. We have perhaps here a defensive reaction of the organism directed against a possible invasion of the blood by sero-saprophytes. And we may perhaps look in this direction for an explanation of the non-specific benefit which has been observed to follow upon the inoculation of bacterial vaccines. It is clearly not impossible that the inoculation of a bacterial vaccine might contribute both to active and passive defence—to the active defence of the body against a particular microbe by calling forth a production of specific bacteriotropic substances, and to the general passive defence of the organism by calling forth a production of antitrypsin. And these two forms of immunizing response would not necessarily be linked together. The production of specific bacteriotropic substances would no doubt depend upon the quantum of bacterial antigen incorporated; while the production of antitrypsin might perhaps depend upon the breaking down of phagocytes and the liberation of their trypsin.

REFERENCE.

¹ *Technique of the Teat and the Capillary Glass Tube*, Constable, London.

* The quantitative method here employed was in essentials that described in my *Technique of the Teat and the Capillary Glass Tube*. It was varied only in the respect that the series of trypsin, serum, and calcified milk mixtures which are employed in that method were taken up, not into a many-stemmed pipette, but into a long, unmounted wide-bored capillary stem, which had, for the purpose of convenient filling in, been bent round at one end in the flame of a Bunsen into a siphoning curve.

(To be continued.)

A SCHEME, drawn up by Drs. William and Charles Mayo, of Rochester, Minnesota, for the establishment of a £200,000 foundation for medical research, has been submitted to the consideration of the University of Minnesota Board of Regents. It is proposed that the interest of the fund shall be used in research work to be carried out by medical graduates at Rochester.

The Minnesota State Medical Society intends to ask the State Legislature to set aside 10,000 acres of land to be used as a leper colony. According to the *New York Medical Journal*, about fifty persons in the State are known to be suffering from leprosy. It is also proposed to establish a branch hospital at the State Hospital for clinical purposes.

CEREBRO-SPINAL MENINGITIS :

DIAGNOSIS AND PROPHYLAXIS.

IS LUMBAR PUNCTURE JUSTIFIABLE?

BY

A. LUNDIE, M.B., CH.B. EDIN.,

BACTERIOLOGIST, ALDERSHOT COMMAND;

D. J. THOMAS, M.R.C.S., L.R.C.P., D.P.H. OXON.,

SPECIALIST SANITARY OFFICER, ALDERSHOT COMMAND;

AND

S. FLEMING, M.B., CH.M. EDIN., D.P.H. CANTAB.,

SPECIALIST SANITARY OFFICER, ALDERSHOT COMMAND;

LIEUTENANTS, R.A.M.C.

Nobody thinks of performing lumbar puncture on slight grounds. The indications for and against are carefully weighed. The operation is decided on the evidence obtained by a careful clinical examination of the case.

The indications for the operation are perfectly clear. An ingravescent headache, absolutely intolerable to the patient, is a very good reason for lumbar puncture at epidemic seasons. The concomitant symptoms may be diverse. There may be fever or no rise of temperature at all. The pulse may be accelerated, normal, or infrequent. There may be vomiting or no gastric disturbance at all. There may be a rash of one type or another, or the skin may be unaffected. There may be head retraction, or even flexion of the head. The knee-jerks may be normal, exaggerated, or unequal. Kernig's sign may or may not be present. The plantar reflex may be flexor or extensor. (The headache itself may be absent, and the symptoms present may be anomalous, and indicate delay.)

The cranial nerves ought to be examined systematically as far as possible, but it will be best first of all to study the patient's power of attention, before putting questions to him that he may answer at random. If his headache is really intolerable, it will often be observed that in spite of his being able to answer questions, the patient speedily lapses after each one into an apathetic state. He may, indeed, continue to mutter and groan, but he is not alert and alive to his critical condition. This condition contrasts most strikingly with the mental state prevailing in yellow fever. Here the patient is extremely alert. If he is of a certain mental calibre, he may clearly diagnose his own illness, and direct its treatment with the utmost non-chalance. He may fully realize his dreadful danger, and determine to do his level best to fight it out, or he may sink into a state of terror, and discuss the mortality of his disease with the attendant. He remains alert to almost the very end. On the contrary, in cerebro-spinal meningitis, there is early dulling of all the senses, and it is well worth while to stand at the bedside and make these observations, as they are of great diagnostic significance.

In testing the cranial nerves, it is best to adhere to the old system of beginning at the first and proceeding in the proper order. The sense of smell may be dulled or lost, optic neuritis should always be looked for. The movements of the eyeball should be tested. The other special senses should, if possible, be also tested. Hearing is commonly found to be defective, and if so, the deafness must not be mistaken for stupor. Hyperacusis has been observed.

A careful study of the nervous system will show the existence of some random process, and in each case there will be differences, proving that the lesion is not the same in every case, and that it is not possible to systematize the symptoms, so as to draw a classical picture of a typical case from nervous symptoms alone. Each case must be taken on its own merits. The acceleration or retardation of the pulse may indicate the point of greatest inflammation at the time the symptom is observed. The thermic phenomena are probably due to the same cause. It may be a matter of chance whether the thermic centre is involved in an inflammatory process or not. There probably is often a pan-encephalitis, rather than a meningitis; in one *post-mortem* examination the corpus callosum was across without being cut, simply owing to its own softened condition and the weight of the cerebral hemispheres. In another case there was a complete breaking down of the tissue forming the wall of the descending cornu of the lateral ventricle. There was no special reason why the other ventricle should not be similarly affected, unless the