4. Free circulation of air about the wound.
5. Easy dressing.
6. By utilizing it in the transverse position it may be of use in cases of incontinence of urine and faeces.

The apparatus is in use at the Royal Herbert Hospital, Woolwich, and Major Spencer, F.R.C.S., R.A.M.C., reports "that it fulfils its purpose admirably, and enables continuous irrigation of a wound in any part of the body to be carried out efficiently and easily without discomfort to the patient or disturbance when dressings are changed, and without risk of leakage into the bed."

Messrs. Down Bros., London, have made the apparatus for me.

The Croonian Lectures

ON

TRYPANOSOMES CAUSING DISEASE IN MAN AND DOMESTIC ANIMALS IN CENTRAL AFRICA.

DELIVERED BEFORE THE ROYAL COLLEGE OF PHYSICIANS OF LONDON,

By SIR DAVID BRUCE, C.B., F.R.C.P., F.R.S.,

SURGEON-GENERAL, A.M.S.; LATE DIRECTOR OF THE ROYAL SOCIETY'S COMMISSION ON SLEEPING SICKNESS.

LECTURE III.—TRYPANOSOMA GAMBIENSE AND CONGO SLEEPING SICKNESS.

HISTORICAL.

Trypanosoma gambiense was first seen in the blood of an Englishman who had been employed for some six years as master of a Government boat on the River Gambia in West Africa. He was admitted to the Government Hospital at Bathurst on May 10th, 1901, fourteen years ago. Dr. R. M. Forde, the Colonial Medical Officer in charge, examined his blood and saw actively moving bodies, but was unable to recognize their nature. He therefore asked the late Dr. J. E. Dutton, of the Liverpool School of Tropical Medicine, to come to his help.

In the meantime, however, the patient had been invalided to England, where he was treated in a Liverpool hospital for some time. Thinking himself sufficiently recovered to resume duty, he returned to Bathurst in December, 1901, and on December 15th his blood was submitted to Dr. Dutton for examination. Dr. Dutton saw the parasite, described it, and named it *Trypanosoma gambiense*.

But at this time it did not enter into the minds of any one that this trypanosome was the cause of sleeping sickness. It was looked upon as an almost harmless haematozoon, causing no inconvenience to the native and only a slight ephemeral fever, known as "trypanosoma fever," to the white man.

At the same time the finding of a parasite of this kind for the first time in the blood of man raised a good deal of interest, and Drs. Dutton and Todd were sent out in the following year to investigate the question further.

They arrived at Bathurst on September 2nd, 1902, and proceeded to examine a large number of natives for the presence of trypanosomes in their blood. They examined in all 1,043 natives on the Gambia, and found the parasite only in six cases.

They summed up their report by saying that, taking all the facts into consideration, they believed that trypanosoma fever, as it occurred in natives, was a peculiarly mild one, and they suggested the possibility that the natives might bear the same relation to the Europeans as the wild game of Africa to domestic animals in nagana—the tsetse fly disease; in other words, that the black man was immune to the disease, and merely acted as a reservoir of the virus to the more susceptible white man.

For some time, then, the important pathogenic part this trypanosome plays remained unknown, and it was not until 1903, when sleeping sickness was being investigated in Uganda by the Royal Society Commission, that the discovery was made that *Trypanosoma gambiense* was in truth the cause of sleeping sickness.

GEOGRAPHICAL DISTRIBUTION.

Reference to the map published to illustrate my first lecture will show that this species of trypanosome has a wide distribution in the tropical zone of Africa. It extends on the north from St. Louis at the mouth of the River Senegal to the Bahr-el-Ghazal in the Egyptian Soudan; on the east it reaches to the eastern shore of Victoria Nyanza; and on the south to the southern end of Lake Tanganyika, the River Luapula in North-West Rhodesia and Donguela in Portuguese West Africa.

MORPHOLOGY.

When in Nyasaland I received from Dr. Kleine, German East Africa, three monkeys infected with *Trypanosoma gambiense*. These animals had been inoculated with blood from natives suffering from "Congo sleeping sickness" on Lake Tanganyika, and it is from these three strains that this description is taken. We sent for these strains because we wished to have an opportunity of comparing *Trypanosoma gambiense* with the trypanosome causing disease in man in Nyasaland, side by side and under similar conditions. It presents a strong resemblance to the nagana parasite. There are the same short and stumpy, intermediate, and long and slender forms. In size and general appearance these two species so closely resemble each other that one might easily believe them to be varieties of the same species.

Shape.—The shape of the long and intermediate forms is the same as in the nagana parasite, but among the short there is an absence of the blunt-ended forms, which have been seen to be so marked a feature in the Nyasaland trypanosome.

Contents of Cell.—The protoplasm often shows many chromatin granules in its substance.

Nucleus.—In *Trypanosoma brucei*, among the short and stumpy there are large numbers of posterior-nuclear forms. In the Tanganyika strains not a single example of this peculiarity was seen.

Micronucleus.—Small and round, and situated on an average 1.1 microns from the posterior extremity in the short, 1.3 in the intermediate, and 1.8 in the long forms.

Undulating Membrane.—This, as in the other members of this group, is well developed, and thrown into bold folds and undulations.

Flagellum.—The flagellum in the long and intermediate forms is free. There is no free flagellum in the short forms.

Length.—In regard to length there is practically no difference between the two species, as will be seen from Table I.

TABLE I.—Showing Length of *Trypanosoma gambiense* Compared with *Trypanosoma brucei*, Zululand, 1913.

Strain.	Average.	Maximum.	Minimum.
<i>T. gambiense</i> , Uganda	22.1	33.0	13.0
<i>T. gambiense</i> , Tanganyika I	22.7	34.0	15.0
<i>T. gambiense</i> , Tanganyika II	25.8	36.0	16.0
<i>T. gambiense</i> , Tanganyika III	21.3	34.0	16.0
<i>T. brucei</i> , Zululand, 1913	21.0	35.0	12.0
Breadth of <i>T. gambiense</i>	2.31	4.75	1.25

Trypanosoma gambiense is therefore very similar in size and shape to *Trypanosoma brucei*, but it would appear to be possible to distinguish them by the presence of the blunt-ended, posterior-nucleated forms which are so common in the blood of animals infected by the nagana parasite and quite absent in animals infected by the other. But as these posterior-nucleated forms are absent or scarce in the blood of man, this method of diagnosis requires the inoculation of experimental animals, and the study of many preparations of their blood. It would appear to be impossible at present to distinguish between the two species by the microscopical examination of preparations made from the blood of man alone.

SUSCEPTIBILITY OF ANIMALS TO TRYPANOSOMA GAMBIENSE.

I now pass to the pathogenic action of this species on various animals. In this it differs markedly from the nagana parasite, which is essentially a trypanosome of the lower animals, seldom attacking man; whereas the reverse is true of the species under consideration, which under natural conditions is almost wholly confined to man.

When *T. gambiense* is taken directly from a patient suffering from sleeping sickness there is often a good deal of difficulty in passing it on to the lower animals—that is to say, it is difficult to infect experimental animals by the inoculation of infected human blood. This is very different from the behaviour of *T. brucei*, which infects most animals with the greatest readiness. It is almost impossible at first to give this disease to goats, monkeys, dogs, and guinea-pigs. The rat is the animal which is least refractory. However, after the trypanosome has become accustomed by several passages through a particular kind of animal, its virulence for this species is heightened. It must also be noted that there is reputed to be a good deal of difference in the virulence of different strains.

This difference in regard to virulence between the two species is shown in Table II, where the average duration in days of the disease caused by *Trypanosoma gambiense*, Tanganyika, is compared with that caused by *Trypanosoma brucei*, Zululand.

TABLE II.

	Monkey.	Dog.	Guinea-pig.	White Rat.
<i>Trypanosoma gambiense</i> ...	159	96	264	137
<i>Trypanosoma brucei</i> ...	26	34	67	30

The disease in animals caused by *T. gambiense* is thus much more chronic than that caused by *T. brucei*, and this character, combined with the morphology already described, affords the surest and safest means of separating these species.

GLOSSINA PALPALIS THE CARRIER OF TRYPANOSOMA GAMBIESE.

As we have already seen, the carrier of *T. gambiense* is the tsetse fly, *Glossina palpalis*, whose habitat is the wooded shore of lakes and rivers.

When the Royal Society Commission in 1903 had convinced itself that in all probability a trypanosome was the cause of sleeping sickness, naturally, on the analogy of the old nagana work, a tsetse fly was looked for. It is a curious fact that at that time the presence of tsetse flies on Lake Victoria was unsuspected.

I have already described the habits of this species of fly, and shown how very numerous it is in Uganda, swarming on the lake shore in such numbers that our fly boys, who collected about 500 daily for months together from one or two spots, did not seem to make any impression on the supply.

Infectivity of Wild *Glossina palpalis*.

Some of these flies were found to be naturally infected with *T. gambiense*, and I shall therefore, in the first place, consider the natural infection which exists among the wild *Glossina palpalis*.

It will be remembered that the proportion of tsetse flies in Nyasaland infected with Nyasaland sleeping sickness was 1 in 500. It will be interesting, then, to compare the infectivity of the wild tsetse flies in Uganda.

It must be granted that although the shores of Victoria Nyanza and the banks of the Nile and other rivers swarmed with these tsetse flies before 1898, it was only about this time that they became infected with sleeping sickness. Before this time sleeping sickness was unknown in Uganda. It seems probable that the disease was introduced by Emin Pasha's men, who were brought into the country from the Belgian Congo some little time before.

I am sorry that the same method of examination was not used in Uganda as in Nyasaland. In Nyasaland the flies were fed on three different species of animals for nine days. In Uganda the flies were caught on the lake shore, and when brought up to the laboratory were only fed on one species of animal—the monkey. In 1903 at Entebbe, the Government cantonment, the tsetse flies had plenty of opportunity of becoming infected, since they were caught in the vicinity of the hut tax labourers' camp. These men came in thousands to Entebbe to work for Government for one month in lieu of paying hut tax. They lived in rudely built grass huts near the lake shore, and on

examination of their blood some 30 per cent. of them were found to harbour the parasite.

In 1903, while these highly infected labourers were living on the lake shore, the proportion of infective flies was found to be as high as 11.2 per 1,000. The Government removed the hut tax labourers from the vicinity of the lake, which became deserted, and a year afterwards the proportion of infected flies fell to 1.2 per 1,000.

When the Commission returned to Uganda in 1908 and took up camp at Mpumu at the north end of Lake Victoria we found the lake-shore flies in the vicinity still infective, although the population had been removed early that year. The examination of 7,200 flies gave a proportion of 1.8 per 1,000.

But we had given the Government to understand that as soon as the natives were removed the flies would become harmless. It was therefore important to find out how long the lake-shore flies remained infective, and why they remained infective. For this purpose they were examined every year until 1912, and the result is given in Table III. It will be seen that although there had been a steady decrease in the proportion of infective flies, a few remained, and these showed no sign of disappearing.

TABLE III.—Infection of Wild *Glossina palpalis*.

Year.	Locality.	No. of Flies Examined.	No. of Flies Infective.	Proportion of Infective Flies per 1,000.	Remarks.
1903	Entebbe	(?)	(?)	11.2	
1904	"	(?)	(?)	1.2	
1908	Mpumu	7,200	11	1.8	1 in 651
1909	"	18,691	7	0.4	1 in 2,670
1910	"	27,179	4	0.14	1 in 6,795
1911	"	23,899	1	0.04	1 in 23,899
1912	"	28,279	4	0.14	1 in 7,070

The mistake made by the Commission was, first, in believing that the transmission of the *Trypanosoma gambiense* was mechanical, and that a fly lost its power of infection within three days after feeding on an infected animal; and, secondly, in believing that man was the sole reservoir of the virus. It was found that a fly may remain infective for several months, and that man is by no means the only source of the virus.

This prolonged infectivity which some flies possess is due to the fact that in these the trypanosomes do not die off, but proceed to further multiplication, and I must now describe the cycle of development which the parasite passes through in the fly, from its ingestion with the blood until its appearance in the salivary glands of the fly in an infective form. When dealing with *Trypanosoma brucei*, I mentioned that its development in the fly is identical with what takes place in the case of *T. gambiense*, and that it would be more convenient to describe the cycle at this point. I did not describe the development of the nagana parasite in *G. morsitans*, as one description will suffice for both species.

The Cycle of Development of *Trypanosoma gambiense* in *Glossina palpalis*.

It is to Kleine that the honour is due of being the first to show that a tsetse fly could convey the infection some fifty days after the fly had fed on an infected animal, and when we come to consider the difficulties of this observation we must feel that he deserves the very highest credit for making this important advance in our knowledge of trypanosome diseases. He used the tsetse fly, *G. palpalis*, and the nagana trypanosome in his first successful experiment, which was carried out at the end of 1908.

A few months later, in March, 1909, we had our first successful experiment in Uganda with *T. gambiense* and *G. palpalis*.

On March 5th, 1909, sixty *Glossina palpalis* were placed in two cages, thirty in each. The flies were fed on two infected monkeys for two days. They were then starved for 72 hours to get rid of the danger of mechanical transference. The following five days they were fed on a healthy monkey, and every successive period of five days, or thereabouts, on a fresh, healthy

monkey, up to 86 days, when the experiment came to an end. The result was that the first two monkeys remained healthy, and all the following monkeys up to 75 days became infected with sleeping sickness. If five days be deducted for the incubation period, then the flies became infected 20 days after their first feed on the infected animal.

There is some evidence that among the 60 flies only one was infective. Fifty-four days after the beginning of the experiment, each cage was placed on a separate monkey. Up to that time both cages of flies had been fed on the same animal. Cage A contained, after fifty-four days, 11 flies; Cage B, 4 flies. Cage A continued to infect monkeys for twenty-one days more, making a total of seventy-five days. Cage B did not infect. Again, as was natural, the flies gradually died off during the experiment, and as each fly died it was carefully dissected and examined for trypanosomes. Not a single trypanosome of any kind whatever was seen in any dissected fly up to 75 days, when a fly died in Cage A, and this fly, on examination, was found to be swarming with trypanosomes. After the death of this fly, Cage A ceased to be infective, and when the experiment was stopped the remaining flies were killed off and dissected, but among them not a sign of a trypanosome could be seen. In the same way the flies in the non-infective Cage B were examined, with a similar negative result.

This was a most interesting and successful experiment. It was evident that a single infective fly did all the mischief and by good luck this fly was detected.

Not only did one fly do all the mischief, but judging from the incubation periods it would appear that in all probability it infected each animal on the first day it bit it, showing how dangerous such an infective fly is. When this fly came to be dissected, the gut was found to be crammed with innumerable trypanosomes, and in addition the salivary glands contained large numbers.

This is the first record of these parasites being found in the salivary glands, and led up to the discovery that the invasion of the salivary glands is an integral part in the cycle of development of this group—Group A—of trypanosomes.

After this, numerous experiments were made on the same lines, and with several species of trypanosomes.

The Development of Trypanosoma gambiense in Laboratory-bred Glossina palpalis.

In 42 experiments performed to ascertain what proportion of laboratory-bred flies become infective when fed on an infected animal, only 8 gave a positive result, as many as 34 were negative. In the 8 positive experiments 371 flies were used, an average of 46; in the 34 negative experiments 1,323, an average of 40. The shortest time which elapsed before a laboratory-bred fly became infective with *T. gambiense* was 27 days, the longest 53 days, and the average 36 days. In the 42 experiments, 1,694 laboratory-bred flies in all were used.

If we consider that in each of the eight positive experiments only one fly became infective, then only 8 flies in 1,698, or 1 in 212, (0.5 per cent.) became infective. This is a very small proportion and helps to account for the fact that fifteen years elapsed before this cycle of development was discovered. This development of the trypanosome in the fly is somewhat similar to what occurs in the test-tube in artificial cultivation. A thousand tubes are inoculated, let us say with *T. brucei*, the trypanosomes all appear to die off, but twenty days later a peculiarly resistant individual is found in one tube of the thousand, which has adapted itself to the new environment, and soon multiplies into myriads.

But, although it is evident that in these 42 experiments only 8 or 10 flies became infective, a larger number were found in which development had proceeded to some extent—that is to say, the flies had become infected, but not infective. As each fly died it was dissected. In all, 39 flies were found with trypanosomes in the alimentary tract—that is to say, about 2 per cent. We may therefore summarize this series of experiments by saying that 0.5 per cent. become infective and 2 per cent. infected. It must be noted, however, that a good deal of difference exists in different series of experiments, the proportion of infected flies rising sometimes to 8 or 10 per cent.

The Cycle of Development of Trypanosoma gambiense in Glossina palpalis.

To find out what occurs after a trypanosome has been swallowed by a tsetse fly, we fed laboratory-bred tsetse

flies on an animal whose blood contained numerous trypanosomes, and, at the end of various times, killed the flies and dissected them. This was done for periods of one day, two days, three days, and so on, for fifty-six days.

The flies were examined first in the fresh condition, when the trypanosomes could easily be seen by their movement; afterwards stained specimens, made for more minute examination, were examined day after day, and coloured drawings, at a magnification of 2,000 diameters, made of all the different forms met with. The drawings of the trypanosomes found in the fore-gut, mid-gut, hind-gut, proctodaeum, and salivary glands were kept separate, so that a series of drawings of trypanosomes taken from any one part, from the first day of infection to the fifty-sixth day, could be compared. More than 600 drawings were made in this way, so that it seems impossible that any important form could have been left out.

General Consideration regarding the Development of the Trypanosomes in the Fly.

For the first three or four days after flies have had a feed of infected blood, trypanosomes are found in them all. These are those originally ingested with the blood, and at the end of six or seven days, when the process of digestion has been completed, they are found to have disappeared from most of the flies; that is to say, it is only in a certain percentage that further development takes place. In one series we saw that this was as low as 2 per cent.; in another it rose to 8 per cent. In other words, in some 95 per cent. of flies which imbibe infected blood, the trypanosomes simply degenerate and die out within the first few days. In some 5 per cent., on the other hand, the trypanosomes find conditions more favourable for development, and increase, filling the whole of the fore-gut, mid-gut, and hind-gut with countless swarms of multiplication forms.

How long this infection of the fly continues is not absolutely known. It is considered probable that in many cases it continues for the rest of the fly's life. In one case a fly was found to retain its infectivity for ninety-six days, but in a few cases there was evidence that an infective fly might in its lifetime lose its infectivity and become harmless.

Types of Trypanosoma gambiense found in the Alimentary Canal of the Fly.

It would serve no good purpose to describe separately, day by day, the variously shaped trypanosomes found in the different parts of the alimentary canal, since the different forms or shapes run into each other in such a way as to make any classification of them seem impossible. The results with regard to the number of trypanosomes found in the different parts of the alimentary canal were as follows:

The Proboscis.—In our experience *Trypanosoma gambiense* is never found in the proboscis of the fly, except immediately after an infected feed, when for a short time blood containing trypanosomes may be seen in the lumen of the proboscis.

Proventriculus.—This part of the alimentary canal is sometimes found empty when the remainder of the gut is swarming.

Fore-, Mid-, and Hind-gut.—It is here that the greatest development of the trypanosomes is found. Among the extraordinary numbers and diversity of type it is difficult or impossible to find one's way. Generally speaking, the trypanosomes found during the first few days are merely degenerated blood forms.

After this there appears a type of trypanosome which remains dominant throughout the whole developmental period. This is a long, moderately broad form, the protoplasm staining well, without granules or vacuoles, having an oval compact nucleus situated in the centre of the body, a small, round micronucleus lying at some distance from the elongated snout-like posterior extremity. The undulating membrane is narrow and simple, and the flagellum proceeds little, if at all, beyond the protoplasm of the cell. The flagellum also appears very frequently to arise from a pink-coloured body situated near the micronucleus, an appearance never seen in the normal blood trypanosomes.

This seems to be the normal developing type in the intestine of the fly. It is seen in all parts of the intestine and at all times.

It forms masses of innumerable individuals alike in size and shape.

When a fresh supply of blood is taken by the fly, this type can be imagined to multiply with extraordinary rapidity. When the blood supply runs low then this type can also be imagined as degenerating and disappearing just as rapidly. The host of divers forms which thus arise

their way by the proboscis and salivary duct or hypopharynx to the glands. They have never been found in the body cavity.

Another curious fact is that as soon as the trypanosomes reach the salivary glands they revert to their original blood form and become infective. What causes or leads up to this reversion to the blood type in the salivary glands is quite unknown, but as we shall see, the *G. palpalis* does not become infective until this invasion of the salivary glands has taken place.

ILLUSTRATIONS OF VARIOUS MODIFICATIONS IN SHAPE OF *TRYPANOSOMA GAMBIENSE* IN *GLOSSINA PALPALIS*.

Fig. I.—*Trypanosoma gambiense* in fly.

1-3, Normal blood parasites (*T. gambiense*). 4, Twenty-four hours after ingestion by the fly. 5 and 6, Forty-eight hours after ingestion by the fly. 7, Ninety-six hours after ingestion by the fly. 1-7 represent the trypanosomes as they appear in the intestine of *Glossina palpalis* during the first few days. 1-3 are ordinary blood forms, as seen immediately after the fly has fed, and before any change has taken place. 4-7 represent the process of degeneration which takes place during the first four days. The body swells up, the nucleus breaks up, and the cytoplasm becomes vacuolated.

Fig. III.—Part of a mass of *Trypanosoma gambiense* from the mid-gut.

Fig. V.—Masses of protoplasm.

45, 46, and 47, *T. gambiense* from hind-gut, ten days after infected feed. 48, *T. gambiense* from fore-gut, seventeen days after infected feed. 49, *T. gambiense* from proventriculus, thirty days after infected feed. 50, *T. gambiense* from mid-gut, forty-six days after infected feed. 45-50 represent some of the more exaggerated types of degenerative forms. As will be seen from the drawings, they are huge, misshapen masses of protoplasm, multi-nucleated and, as a rule, multi-flagellated.

beggars description. Some are round or oval in shape, 3 or 4 microns in diameter, with or without flagellum, and from this simple form all shapes and sizes can be seen up to the huge, shapeless mass of protoplasm, multi-nucleated and multi-flagellated.

Salivary Glands.—Trypanosomes did not appear in these glands until the twenty-fifth day, but after this time they were usually present. This invasion of the salivary glands is the most interesting phase in the development of Group A, and brings to mind the development of the malarial parasite in the mosquito. It differs in this, however, that whereas in the latter spores are found, which pass to the salivary glands across the body cavity, in the tsetse fly the trypanosomes apparently find

Fig. II.—Developmental forms.

8, *T. gambiense* from fore-gut, eight days after infected feed. 9, *T. gambiense* from proventriculus, fourteen days after infected feed. 10, *T. gambiense* from fore-gut, eighteen days after infected feed. 11, *T. gambiense* from mid-gut, twenty-five days after infected feed. 12, *T. gambiense* from mid-gut, thirty-four days after infected feed. 13, *T. gambiense* from fore-gut, forty-four days after infected feed.

Fig. IV.—Rounded forms.

25, *T. gambiense* from mid-gut, eight days after infected feed. 26, *T. gambiense* from fore-gut, eleven days after infected feed. 27 and 28, *T. gambiense* from mid-gut, sixteen and seventeen days after infected feed. 29, 30, and 31, *T. gambiense* from fore-gut, seventeen and eighteen days after infected feed. 32, *T. gambiense* from hind-gut, twenty days after infected feed. 33 and 34, *T. gambiense* from fore-gut twenty-four days after infected feed. 35, 36, and 37, *T. gambiense* from mid-gut, twenty-four, forty-four, and forty-six days after infected feed. 38, *T. gambiense* from fore-gut, forty-six days after infected feed.

Fig. VI.—Infective forms.

73, *T. gambiense* from salivary glands, thirty-four days after infected feed. 74, *T. gambiense* from salivary glands, forty-two days after infected feed. 75, *T. gambiense* from salivary glands, forty-three days after infected feed. 76, 77, and 78, *T. gambiense* from salivary glands, forty-six days after infected feed. 79 and 80, *T. gambiense* from salivary glands, fifty-six days after infected feed. When alluding, generally, in a previous part of this lecture, to the types of *Trypanosoma gambiense* found in the salivary glands, it was said that in the salivary glands, and here alone, the trypanosomes are found to revert to the normal type found in the blood. 73-80 illustrate this reversion. By comparing them with 1, 2, and 3 (Fig. I), which represent normal blood trypanosomes, it will be seen that they are very similar to the short and stumpy form found in the blood.

The salivary glands first become invaded twenty-five days after the infecting feed. In all the salivary glands from flies which gave a positive result, trypanosomes similar to the short blood-type were invariably present.

Experiments to Ascertain if *Trypanosoma gambiense* during its Development within *Glossina palpalis* is Infective.

In Zululand it was observed—and at the time it seemed to be an astonishing fact—that when tsetse flies, a short time after feeding on infected animals, were chopped up and injected under the skin of healthy dogs no infection of nagana took place.

It seemed strange that a fly, swollen out with blood in which numberless active trypanosomes could be seen under the microscope, if injected under the skin of a susceptible animal, should not give rise to the disease. As I left Zululand at this time I was unable to follow this observation up, but in Uganda I carried out experiments on the same lines with *T. gambiense* and *G. palpalis*.

If tsetse flies (*Glossina palpalis*) are fed upon an animal whose blood contains trypanosomes, the parasites can be seen living and moving in the contents of the intestines of all the flies for a few days, and in a small percentage (2 per cent. to 8 per cent.) of the flies active trypanosomes may continue to be found swarming in their intestines on any day between the seventh and fiftieth day or even longer.

A series of experiments was undertaken in Uganda to ascertain if *Trypanosoma gambiense* retained its power of causing sleeping sickness when inoculated subcutaneously into monkeys during its period of multiplication within the fly, especially during the interval of some twenty or thirty days in which the bites of infected flies had been proved to be harmless.

The flies were fed upon a monkey whose blood contained many trypanosomes. After a pre-determined time had elapsed the wings and legs of the infected flies were cut off and the bodies were either chopped up or brayed in a mortar with saline solution. In most instances the gut was proved by microscopical examination to be heavily infected with trypanosomes before the whole chopped-up fly or flies were placed under the skin of the experimental animal. Of twenty-nine experiments carried out with *G. palpalis* and *T. gambiense*, eighteen gave a positive result; they were almost uniformly successful up to eighteen hours after the infected feed.

Further experiments proved that this parasite only retains its virulence for a period of about eighteen hours after it has been swallowed by the tsetse fly; it then loses its power of infecting animals for a period of some twenty-two days. This means that during their period of development in the tsetse fly the trypanosomes lose their virulence, and do not regain it until the developmental cycle is completed.

Summary.

Trypanosomes taken into the alimentary canal of tsetse flies retain their shape and infectivity for some eighteen hours. They then degenerate and lose their power of infection, and, as a rule, have disappeared altogether from the majority of the flies by the fifth or sixth day. In a small percentage of flies, male as well as female, the trypanosomes maintain their position; they continue to multiply, and in a short time swarm in the alimentary canal of the fly. These multiplication forms bear little or no resemblance to the original trypanosomes. After some twenty days the developing flagellates find their way into the salivary glands, resume their original blood form and regain their infectivity.

THE RESERVOIR OF TRYPANOSOMA GAMBIENSE (CONGO SLEEPING SICKNESS).

We have seen that in *T. brucei* the cause of "Nyasaland sleeping sickness," the reservoir of the virus was undoubtedly the wild game, and that in all probability man played a very small part, if any, in its spread. I have already described *T. brucei* as essentially a parasite of animals, and *T. gambiense* as a parasite of man.

In Uganda cases of sleeping sickness were limited to the edge of the lake and the islands; I say "were," because the epidemic has died out. There was formerly a great deal of trade intercourse between the islands and the mainland. Under the shade of the trees, on the shore of the lake, the natives met, and here the fly was thickest. Now we know that in the sleeping sickness area in Uganda anything from 20 to 50 per cent. of the lake shore population had sleeping sickness trypanosomes in their blood, and it therefore appeared extremely probable that the fly conveyed the virus from native to native.

We believed at first (1903) that man was the only reservoir of practical account, and that if he was removed from the fly area, the fly would soon—within a few days—become non-infective. When we returned to Uganda in 1908 we were much surprised to find the fly still infective, although the native population had been removed from the

lake shore for some months. It was evident, therefore, that the fly had some other reservoir from which to draw the virus, and experiments were instituted to throw light on this. The cattle were first examined. These animals, as we have seen, are practically immune to sleeping sickness—that is to say, they show no signs of illness, but it was possible the trypanosome might live in their blood, and be capable of infecting flies without affecting the health of the cattle.

Cattle as a Reservoir.

On September 9th, 1909, an ox was inoculated with a quantity of blood containing large numbers of the parasites. The blood of the ox was examined daily, and eighteen days after the injection was found to contain the trypanosome. Its identity was established by injecting a healthy monkey with some blood from this ox. The monkey showed *T. gambiense* in its blood on the sixth day. Cattle are therefore capable of being infected with sleeping sickness by the injection of infected blood, although they show no signs of illness. Cattle were also infected by the bites of artificially infected flies.

Next, freshly-caught flies, brought up to the laboratory from the lake shore, were fed on healthy cattle. In the first experiment, 2,195 freshly-caught flies were fed on an ox during sixteen days, when the animal was found to be infected. In the second, 1,370 flies, and in the third, 459, gave a positive result.

The next question to be asked was, Is it possible to infect tsetse flies by feeding them on these infected cattle, and afterwards to transmit the disease by means of these flies to healthy animals? Five experiments were carried out. Two of them were positive.

It was therefore proved that laboratory-bred flies can be infected by feeding them on infected cattle, and afterwards the disease can be transmitted to healthy animals by means of these flies.

Finally, it had to be ascertained whether it was possible to find in the sleeping sickness area cattle which had been naturally infected.

Seventeen cattle from various sources were examined with this point in view. Most of them came from places where the tsetse flies were numerous. In one out of the seventeen the parasite was found. This ox came from the island of Kome in Lake Victoria, where sleeping sickness was prevalent and where tsetse flies abound. A small quantity of its blood injected into a healthy monkey gave rise to sleeping sickness in the monkey, and showed that *T. gambiense* was really being dealt with.

The conclusion to the whole matter, then, was that in addition to man, cattle may act as a reservoir of the virus of sleeping sickness.

Antelope as a Reservoir.

We next turned our attention to antelope. At this time, the beginning of 1910, man and his domestic animals had been removed from the lake shore of the mainland for some two years, and from the islands since the previous September. The effect of this depopulation had been to make an area along the northern shore of the lake virtually a game reserve, in which waterbuck, bushbuck, reedbuck, Speke's tragelaphus, hippopotami, wild pig, and other game abound. Before this time we had a strong preconceived idea that cattle and antelope were immune to sleeping sickness, and these preconceived ideas are stubborn things. But after the cattle experiments we set ourselves to ask the same questions regarding antelope.

Eleven antelope in all were employed in this experiment—four bushbuck, six reedbuck, and one waterbuck. A cage of flies which was known to contain an infective fly or flies, was fed on each animal daily for several days. The result was positive in all cases, proving that antelope may be readily infected with sleeping sickness by the bites of infective flies.

The next question was, Can these infected antelope transmit the infection to clean laboratory-bred flies, and if these flies become infected, can they transmit the virus to susceptible animals?

Clean laboratory-bred *Glossina palpalis* were fed for several days on an infected antelope. After an interval of starvation of twenty-four hours or more the flies were transferred to healthy animals and fed daily. When the healthy animal showed *Trypanosoma gambiense* in its

blood the experiment was stopped, and the surviving flies were dissected as soon as possible.

The result of twenty-four experiments carried out on these lines is given in Table IV.

TABLE IV.—Result of Feeding Laboratory-bred Flies on Antelope Infected with Sleeping Sickness.

No. of clean flies used.	Species of antelope flies fed on.	No. of days flies fed on antelope.	No. of days before flies became infective.	Result.	Remarks.
160 100 70 100	Bushbuck 2328	12 8 6 8	29 28 29 39	+ + + +	Buck 2328 never showed <i>T. gambiense</i> in blood. In spite of this, flies fed on it became infected 55 days after the buck's infection.
100 100 100	Reedbuck 2357	7 8 5	41	+	Buck 2357 showed <i>T. gambiense</i> in its blood for 5 days only.
200	Reedbuck 2359	6	44	+	Buck 2359 showed <i>T. gambiense</i> in its blood for 7 days only.
50 60	Bushbuck 2371	6 6	20	- +	Buck 2371 showed <i>T. gambiense</i> in its blood for 3 days only.
100	Bushbuck 2372	8		-	Buck 2372 showed <i>T. gambiense</i> in its blood for 2 days only.
95 60 50	Waterbuck 2378	6 6 4	30	+ - -	Buck 2378 never showed <i>T. gambiense</i> in its blood.
110 60 100	Reedbuck 2427	6 4 6	24 33 30	+ + +	Buck 2427 showed <i>T. gambiense</i> in its blood for 4 days only.
50	Bushbuck 2428	7	28	+	Buck 2428 never showed <i>T. gambiense</i> in its blood.
50 100	Reedbuck 2429	4 6	27 49	+ +	Buck 2429 showed <i>T. gambiense</i> in its blood for 3 days only.
55 90 100	Reedbuck 2431	3 6 5	28 36 43	+ + +	Buck 2431 showed <i>T. gambiense</i> in its blood for 4 days only. In spite of this, flies fed on it became infected 81 days after its infection.
50	Reedbuck 2445	4		-	Buck 2445 showed <i>T. gambiense</i> in its blood for 6 days only.

The most significant of these experiments is the one in which it is shown that fifty-five days after the last feed of the infected flies on bushbuck 2,328 the blood of the buck was still capable of infecting clean laboratory-bred flies, although *Trypanosoma gambiense* had been so scarce in the blood that they had never been seen microscopically.

These experiments show that antelope of the waterbuck and bushbuck species, when infected with the trypanosome of sleeping sickness, can transmit the infection to clean laboratory-bred *Glossina palpalis*. The infected antelope's blood was in one case capable of giving rise to infection for at least eighty-one days, and in another for at least fifty-five days. These experiments further show that the flies, when infected, are capable of transmitting the disease to susceptible animals.

From these experiments on antelope, the curious fact seems to emerge that a higher proportion of flies become infected by feeding on infected antelope in whose blood the parasites are scanty, than, for example, on infected monkeys, whose blood swarms.

Another curious fact is that even when trypanosomes were never found by microscopical examination in the blood of an antelope, flies fed on that antelope became infected. It would almost lead one to believe that the trypanosomes exist in the blood of these animals in some

other less easy recognizable form than the ordinary adult trypanosome form.

The next thing we set ourselves to do was to find out if the antelope living in the fly area are naturally infected with sleeping sickness. Positive evidence on this point would naturally complete the chain of evidence proving that antelope living in the fly area may act as a reservoir of the virus of sleeping sickness. So far it had only been proved that they were potential hosts.

The easiest method by which it could be proved that they are naturally infected would be the old one of shooting the animals, and at once sending samples of their blood into some healthy area where the blood could be injected into healthy susceptible animals. But those who know the local conditions in Uganda will recognize the difficulty of doing this. The cover is so thick at the edge of the lake that it is exceedingly difficult to see the game. The sun is almost unbearably hot, the country very difficult, and the exposure to the bites of the fly very great.

It may be said at once that we were only able to shoot five antelope on the lake shore. The blood of these, when injected into susceptible animals, gave negative results. A large drive was being organized, which might have thrown the desired light on this subject, but unfortunately the Commission was ordered to leave the country before the drive could take place. It was, therefore, perforce left to a later worker to solve this difficult problem.

In 1911 Dr. H. L. Duke took the question up. Four years had now elapsed since the lake shore was depopulated, and two years since the islands had been emptied of their population. He chose the uninhabited island of Damba for the experiment. This he did because the Commission had shown in 1910 that the fly on the island were still infective, and again the same result was obtained in May, 1911, by Dr. Carpenter.

With the exception of certain patches of papyrus, practically the whole shore line of the island swarms to a greater or less extent with fly. At one part there is the famous fly beach, from which for several years some thousands of pupae have been brought monthly to the sleeping sickness camp at Mpumu.

The only species of antelope found on the islands of Lake Victoria is the sitatunga (*Tragelaphus spekei*), a species of bushbuck adapted for living in marshes. Dr. Duke shot four sitatunga on the island of Damba, and from these *T. gambiense* was obtained. I cannot say whether all four antelope harboured the parasite, because the blood of the first two was injected into one monkey and the blood of the last two into another. Both monkeys showed the *T. gambiense*. Dr. Duke examined these trypanosomes with meticulous care in order to exclude any possible source of fallacy, and there cannot be any reasonable doubt that the sitatunga on Damba Island were acting at that time as a reservoir of this parasite.

Another proof that the trypanosome found in the sitatunga was really *T. gambiense* has lately been given by Dr. Duke. He states that two of the fly boys who have during the last three years worked with Dr. Carpenter on the islands have developed sleeping sickness of the Uganda type, and trypanosomes have been demonstrated in their glands. These boys have only been exposed to the bite of *G. palpalis*, as they had for eighteen and thirty months respectively resided on the lake shore and islands, where there are no other species of tsetse fly.

This shows conclusively that *T. gambiense* still exists in these lake shore and island flies five years after the removal of the population, and is another striking proof that the antelope are acting as a reservoir.

Taking everything into consideration, then, we may look on it as proved that the antelope on the shores of Lake Victoria act as a reservoir of the virus of sleeping sickness, and that this accounts for the fact that the flies have retained their infectivity in spite of the removal of the native population from the lake shore and the islands.

The prophecy of the Commission that the fly would become harmless shortly after the natives were removed from the lake shore has unfortunately proved wrong, and before the islands are repopulated some other measure will have to be taken to get rid of the fly danger.