tissue (Figs. 1, c, 2, c, and 3, c) which owing to thrombosis of the neighbouring vessels soften and are converted into cysts. To this abnormal vascularity also must be attributed the pain which some individuals feel in the smaller

Fig. 3.—Section through the epiphysis and adjoining part of the diaphysis of a first metatarsal bone. a, Fistulous opening through which a glass rod has been passed into a cyst b, which has formed in the area of fibrous tissue, c.

bones of the hands or feet after the consumption of even one glass of a generous wine. This pain is commonly regarded as a sign of gout and attributed to uric acid, whereas in reality it proceeds from an undue relaxation and consequent over-distension of the vessels which are deprived of their normal support by the osseous trabeculae.

When once this process has reached the articular cartilage and has perforated it at several spots, the surface of the bone is apt to be worn off and laid bare, in consequence of which it becomes inflamed and rarefied. The first

phalanx may then be deflected outward. A frequent complication of this condition is a lymphangitis, and the whole leg may thus become the seat of a leucophlegmatic oedema. It is easily intelligible what mischief may be done if an obscure affection of some internal organ be supposed to be due to "misplaced gout," and if on the advice of some authorities the alleged materia peccans is then directed by mustard poultices to its supposed normal place of exit by the big toe. Such naïve practice presumes that the redness due to the irritation by mustard is equivalent to an attack of gout and serves to relieve the blood of its contained impurity.

This cystoid degeneration is readily distinguished from gout by the absence of swelling of the regional lymphatic glands. It may, however, become gouty if from a concomitant disease of the kidneys uric acid is retained and accumulates in the blood. The tissues, especially those of an inflammatory or a metaplastic nature, then undergo hyaline degeneration, and favour the deposition of uratic crystals.

Such change in the type of the disease is then shown by the accession of the diagnostic element of gout—namely, the enlargement, as just mentioned, of the regional lymphatic glands, which have also become hyaline, as I have already pointed out.¹ My observations of the involvement of the lymphatic system in these circumstances have been confirmed by Minkowsky, who apparently was unaware of my statement to that effect, which had been published ten years before the appearance of his work on gout.

REFERENCE.

¹ Vide *Lancet*, December 16th, 1893, Fig. 3.

WE learn that work has been begun on the hospital which is to form part of the School of Tropical Medicine in Calcutta, and that it is hoped that the building of the hospital will be completed next year, the necessary funds having been collected.

THE United States Public Health Service has recently set on foot an active campaign of education for the prevention of typhoid fever. The proportion of typhoid cases in the United States is said by the *Bulletin* of the Service to be from two to five times that in many European countries where systematic preventive measures are enforced. The number of deaths from the disease in the United States last year is stated to have been 30,000, while 400,000 persons were incapacitated. During the last forty years, although there has been a considerable reduction in the prevalence of the disease in some cities, the United States as a whole are a generation behind the time in respect to the reduction of the typhoid rate.

THE CARE OF THE ILLEGITIMATE BABY.

BY

A. DINGWALL FORDYCE, M.D., F.R.C.P. EDIN.,
EXTRA PHYSICIAN, ROYAL HOSPITAL FOR SICK CHILDREN,
EDINBURGH.

I AM unaware of any reports of investigations into the welfare of poor illegitimate babies throughout the early years of life, and consequently the results of an investigation undertaken in Edinburgh in the early part of 1914 may be of interest.

The object of the investigation was to trace the history of illegitimate babies in the city for three years after birth in order to gain an idea of the care bestowed upon them as compared with legitimate babies of their own class. Several ladies very experienced in social work most kindly assisted me in the work, and their aid proved invaluable, as is evident from the very considerable number of cases in which a thoroughly satisfactory and first-hand history was obtained.

Conditions in Edinburgh.

Illegitimate infants born in the Royal Maternity Hospital, Edinburgh, January 1st to June 30th, 1911, and their history for three years—that is, till June 30th, 1914:

(a) Dead-born or died practically at birth...	14
(b) No trace of mother or child ...	38
(c) "Reports" about ...	7
(d) Satisfactorily traced ...	82
Total ...	141

Of 82 cases traced for three years:

Child known to be alive and cared for ...	54
Child presumably cared for ...	9
Cases in which child died and mother known to have been good ...	6
	69

So that in 69 out of 82 cases the babies had fair attention.

Cases in which baby died in questionable circumstances ...	5
Cases in which the conditions prior to death were uncertain ...	8
	13

What do these statistics show?

1. It was possible to trace 82 cases for three years after the birth of the child out of 127 cases in which the child survived birth.

2. Of these 82 cases, the mother afterwards married in 36 cases.

3. In 69 out of these 82 cases the baby had fair attention.

4. Forty-five cases could not be traced satisfactorily, or at all.

It would seem that of cases under heading 4 a certain number could not be traced because of shame on the part of a good mother, but in many cases it was certainly because an undesirable mother wished to leave no tracks. It may be taken that in 75 per cent. of cases the illegitimate baby received fair attention, and that in 40 per cent. of cases the unmarried mother later married.

It seems reasonable to conclude that an illegitimate baby in 1 out of 4 cases does not receive the ordinary attention given to his legitimate brother. This does not mean that 3 out of 4 are properly attended to. At best it means that 3 out of 4 receive the ordinary treatment of legitimate children of that class. In the out-patient department of any sick children's hospital in a large city the time of the physician is to a very large extent occupied by attending to children under 3 years of age who suffer from preventable diseases. Their mothers are well-meaning and kindly, but, either from ignorance on their part or the effect of an unwholesome environment, or both combined, the children become unhealthy. But, distinct from these well-meaning though ignorant mothers are most of the unmarried mothers. In some cases they are as unselfishly devoted to their child as any mother could be. More commonly they are fond of their child, but the times of stress they have been through have left their indelible mark; they are laden, overborne, oppressed by the difficulties of their situation, they recognize the claims of the child on them and would respond if they could; but they cannot, and though they sincerely regret it, there seems little hope

for the future except by the death of the child. Other illegitimate babies there are who are brought to the hospital when practically moribund, in whose cases death is obviously wished for, and the doctor is looked upon not as a curer of disease, but as a necessary means of securing a death certificate!

Let us not salve our conscience with the unctious that three out of four illegitimate babies are properly attended to. They are not. But all I have been able to show is that one out of four is improperly attended to.

In Hungary since 1903 every homeless child—every child for whom there was no obvious means of suitable support—has had a right to support by the State, and has been so supported.

The result has been that during the years 1904 to 1908 the number of births in the country increased by over 2 per cent., while the number of illegitimate births decreased by over 4 per cent.!

In Edinburgh an infants' home was opened in 1912. Unmarried girls with their first illegitimate baby are admitted to it on quitting the maternity hospital, and on signing a promise to remain in the home for at least two months, and to suckle their babies. Each girl pays 5s. weekly for the first two weeks, and thereafter 2s. 6d. weekly. Girls who spend three months in the home may leave their babies there at a charge of 5s. weekly until they are a year old. The mothers of babies who have been left in the home visit them almost invariably twice a week, and stay to bath, feed, and put them to bed.

The home saves many girls from finding themselves in a hopelessly difficult position on leaving the maternity hospital at the expiry of ten days. Some have no home, and others are forbidden by their relations to return with their babies to the home they have disgraced. The only course open to them is to board out their young babies with the first person who offers, and to seek work, however unfit physically they may be. In the home they have a refuge. A certain amount of the money in payment is sometimes obtained at first from the father of the child, and otherwise the amount is usually obtainable from relatives, or, if necessary, from charitable sources.

The babies are cared for and nursed by their own mothers. They usually thrive well and become strong and healthy. When the baby cannot later go with his mother or to her relatives, the mother has time and the assistance of the matron to find a good home, where he may be boarded out or adopted. This "good home" is, however, extremely difficult to find.

As regards the mother, the first object of the home is to strengthen and develop her character generally, so that when she leaves she may be able to lead a better life.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

PLUMBISM IN A BRANCH OF THE HOSIERY INDUSTRY.

THE following brief report may lead, in other places where the making of military pants is carried on, to the detection of further cases of plumbism.

A. K., a man aged 28, came to see me on May 31st, 1915, complaining of abdominal pains, so severe at times that he could not sleep at nights. On the night of May 30th he was "up all night" in pain. He had been very costive for some weeks. There had been no vomiting, but his appetite was poor and he had lost flesh lately. He referred the pain to the umbilical region. The abdomen was flaccid, not tender to touch; in fact, the pain was somewhat relieved by pressure. He had also "rheumaticky" pains in the left arm and leg during this period of poor health, of some weeks' duration. There was a typical blue line on the gums, but no ankle or wrist drop.

The condition at once suggested plumbism, but his employment, which he described as that of cotton spinner in the hosiery trade, seemed at first to negative this possible source of the poison. On closer inquiry, however, he explained that, in passing the cotton through a rolling machine, it was impregnated with a dry ochre pigment, which gave the pants the regulation tint.

In visiting the "blowing room" where he worked, it

was seen that a cloud of this pigment pervaded the room in the neighbourhood of the machine, and even at some considerable distance from it, covering the walls and roof. The workers stated that they were frequently spitting up the pigment "from their lungs." The strong air draught from below upwards is necessary in order to drive the cotton into the machine. Since a new machine had been introduced some weeks previously the dust had been much more dense. There had also been a much heavier pressure of work owing to large Government orders.

Two other rooms in the same factory have similar machines, and many of the operatives within the dust radius, including some women, are reported to have had poor health lately.

A sample of the ochre revealed the fact that it consisted of "almost pure basic chromate of lead." This induced me to make further inquiry.

A. K. rapidly recovered from his acute symptoms under treatment, while absenting himself from work, and has now returned to the blowing room.

Another worker, G. S., a man aged 40, was seen by me at his home on June 20th. He had been under another medical man's care for some weeks for symptoms similar to those of A. K., accompanied by marked loss of weight and weakness. When I examined him I found a well-marked blue line on his gums.

As no other source of plumbism could be discovered (the water supply of Nottingham being now above suspicion) both cases were notified to the Home Office.

Subsequent inquiry was made by the Medical Inspector of Factories, and I learn from A. K. that precautions are being taken by using another pigment.

Nottingham.

A. CHRISTIE REID, M.D.

Reports of Societies.

OXFORD OPHTHALMOLOGICAL CONGRESS.

THE annual meeting of the Oxford Ophthalmological Congress was held on July 15th and 16th in the Department of Human Anatomy of the University of Oxford, by the kind permission of Professor Arthur Thomson. Notwithstanding the inevitable absence of some of the regular attendants owing to military duties and other reasons connected with the war, a good number were present during the proceedings, which were opened at 9 a.m. on July 15th by a hearty welcome from the Master, Mr. Sydney Stephenson. The morning of this, the first day, was devoted to addresses by Sir ST. CLAIR THOMSON, Sir MACKENZIE DAVIDSON, Lieutenant-Colonel R. H. ELLIOT, Messrs. W. H. H. JESSOP, HARRISON BUTLER, and STEPHEN MAYOU. The subjects chosen were of considerable interest, and each address was followed by discussion. Lieutenant-Colonel ELLIOT prefaced his remarks with words of congratulatory welcome to Mr. Sydney Stephenson on his first appearance as Master of the Congress in place of Mr. Robert W. Doyne, now the Past Master. The afternoon was spent at the Oxford Eye Hospital, where interesting cases were shown by Messrs. P. H. ADAMS, WILLIAM ROBINSON, N. C. RIDLEY, E. H. EDWARDS STACK, and B. CRIDLAND. Messrs. RIDLEY, STACK, and MADDOX also gave demonstrations. The whole of the second day (July 16th) was occupied by a discussion, opened by Dr. FRANK SHUFFLEBOTHAM, on *Industrial diseases and accidents*. The subjects dealt with comprised coal-miner's nystagmus, eye injuries caused by occupation with their prevention and first aid treatment, industrial cataract, plumbism as it affects the eye, siderosis, and the rarer industrial diseases and injuries of the eye, and were introduced by Messrs. PERCIVAL J. HAY, EDGAR COLLIS, WILLIAM ROBINSON, S. McMURRAY, J. GRAY CLEGG, and SYDNEY STEPHENSON respectively. The papers were extremely interesting and evoked considerable discussion, in which many members took part. During the two days demonstrations were given by Mr. SYDNEY STEPHENSON, Lieutenant-Colonel R. H. ELLIOT, Messrs. J. BURDON-COOPER, R. J. COULTER, W. H. H. JESSOP, P. H. ADAMS, N. C. RIDLEY, J. GRAY CLEGG, C. T. REES WOOD, and S. E. WHITNALL, comprising pathological specimens, apparatus, and instruments of much interest. An exhibition of ophthalmological instruments and apparatus was open daily. The official dinner of the Congress took