Illness and convalescence. The results were of no particular importance; during the fever there was usually a leucopenia with a relative decrease of polymorphs and increase of lymphocytes and mononuclears; during convalescence there was in some cases, but by no means in all, an eosinophilia. One patient in particular I remember, who had only two days' fever, and whose eosinophilia rose from normal on the first day to 22 per cent. on the fifth. I should mention that in every case malaria was excluded, and in every case of eosinophilia the faeces were examined microscopically.

With regard to the transmission of the disease, I noticed that *Culex fatigans* was the most obvious biting insect about the barracks at the beginning of the epidemic, but that later on *Stegomyia* predominated. Sandflies were present throughout the period, but were never obvious.

Another interesting point about the epidemic was the fact that the first two British officers to be attacked were the only two who did not use mosquito nets, they were both attacked within four days of landing in Rangoon. All the other British officers were attacked later, and one when examined on the first day of illness, showed no less than 42 mosquito bites on the knees, the result of working in "shorts" in the office the night before.

This suggested the possibility of examining mosquitos fed on patients at different stages of the disease; accordingly cultures were taken from the contents of the stomachs of mosquitos (all *Culex fatigans*) fed in this way. I would not have believed it possible to obtain pure cultures in this way, nevertheless in 18 out of 23 cases a pure culture of a bacillus was obtained giving nearly all the reactions of Colonel Rogers's bacillus, but differing from it in its greater motility and smaller tendency to filamentous forms. It was not agglutinated by the serum of the goat mentioned above. I mention this bacillus only as a point of interest.

With regard to the nature of the epidemic itself I am not in a position to judge, as I had not previously seen either dengue or sandfly fever; I take it that the epidemic was a mixed one.

AN ADAPTABLE EYE SHADE FOR MICROSCOPIC USE.

By PROFESSOR S. G. SHATTOCK, F.R.C.S.

My excuse for drawing attention to this device is the advantage (with which I have long been impressed) to be derived from cutting off the access of direct light to the eye above the ocular.

In working in a darkened room with a hooded lamp, the drawback mentioned does not, of course, arise. But when daylight is used the admission of light to the eye above the ocular is a distinct hindrance to the study of fine detail. The difference can be at once brought home by temporarily shading the eye with the hand so as to improvise a dark chamber above the ocular. The ability to see more minute details with an eye shade, and a consciousness of diminished strain, will become so apparent that, once used, I feel sure the device will be afterwards invariably resorted to for any prolonged microscopic study.

Mr. Swift, who makes it to my design, tells me that binocular microscopes were occasionally fitted with shades, but these (as in the case of field glasses) were fixed adjuncts. As the use of binoculars for histological purposes became obsolete, the advantage of the shade appears to have been lost sight of. In some form it seems to be well worth reviving.

The draw-tube of all microscopes at the present time is

Fig. 1.—A diagrammatic section (natural size) showing the shade in situ, as kept in position between the flat, upper edge of the draw-tube A, and the rim of the top of the ocular. The form of the upper opening of the shade is indicated by the dotted line.

made of a standard inside diameter, known as No. 1—namely, 23.3 mm.; and this is the size adopted for the aperture in the floor of the shade through which the ocular drops; but the aperture is made also of a larger size to correspond with standard No. 2—namely, 26 mm.—and it can, of course, be cut so as to take an ocular of any other dimension.

It is hardly necessary to add that the shade can be used for either eye by rotating it so that the higher part corresponds with the outer, receding margin of the orbit. Its cost in black matt brass is 3s. 6d.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

THE ANTITOXIC POWER OF HYPOCHLOROUS ACID.

For some time past experiments have been carried out with a strongly antiseptic liquid consisting of hypochlorous acid and calcium biborate, in order to determine its antitoxic power. The best proportions to use are not yet definitely fixed, but will be given in a fuller account to be published later.

The harmlessness of the liquid in question for guinea-pigs and rabbits is shown in the following table:

TABLE I.

Guinea-pigs.	Quantity of Liquid Injected Subcutaneously.	Effect.
1	1 c.cm.	Slight inflammation at the place of inoculation.
2	2 c.cm.	
3	3 c.cm.	
4	4 c.cm.	
Rabbits.	Quantity of Liquid Injected in Ear Vein.	None.
1	1 c.cm. + 4 c.cm. salt sol. 8 %	
2	1 c.cm. + 1 c.cm.	
3	1 c.cm.	
4	4 c.cm.	
5	5 c.cm.	
6	8 c.cm.	
7	10 c.cm.	

The antitoxic effect of the above-mentioned liquid on diphtheria toxin is demonstrated in Tables II and III.

Neutralization of Diphtheria Toxin in Vitro.

TABLE II.

Guinea-pigs.	Mixture of Toxin and Antitoxic Liquid Injected Subcutaneously.	Result.
1	4 L.D. of toxin + 1 c.cm. of antitoxic liquid	Animals remain healthy.
2	4 L.D. „ + 1 c.cm. „ „	
3	4 L.D. „ + 1 c.cm. „ „	

Three guinea-pigs inoculated with the same quantities of toxin died after less than forty-eight hours.

Neutralization of Diphtheria Toxin in Vivo.

The antitoxic liquid was injected five minutes after the toxin had been inoculated; further doses were given after twenty-four hours and forty-eight hours. Both toxin and antitoxic substance were given subcutaneously on different sides of the body.

TABLE III.

Guinea-pigs.	Quantity of Toxin Inoculated.	Quantity of Antitoxic Liquid Inoculated—			Results.
		After 5 Min.	After 24 Hours.	After 48 Hours.	
1	4 L.D.	1 c.cm.	1 c.cm.	1 c.cm.	Animals remain healthy.
2	4 L.D.	1.5	1.5	1.5	
3	4 L.D.	2	2	2	
4	4 L.D.	2.5	2.5	2.5	
5	4 L.D.	1	1	1	
6	4 L.D.	1	1	1	
7	4 L.D.	1	1	1	
8	4 L.D.	1	1	1	
9	4 L.D.	1	1	1	+ 5 days. Healthy.
10	4 L.D.	1	1	1	

Guinea-pigs inoculated with 4 L.D. of toxin died after less than sixty hours.