The Goulstonian Lectures

ON SPINAL INJURIES OF WARFARE.

DELIVERED BEFORE THE ROYAL COLLEGE OF PHYSICIANS
OF LONDON.

By GORDON HOLMES, M.D., F.R.C.P.,

ASSISTANT PHYSICIAN TO CHARING CROSS HOSPITAL, AND NATIONAL
HOSPITAL FOR PARALYSED AND EPILEPTIC, QUEEN SQUARE;
LIEUTENANT-COLONEL (TEMPORARY) R.A.M.C.

[WITH SPECIAL PLATE.]

I. THE PATHOLOGY OF ACUTE SPINAL INJURIES.

I MUST first express my appreciation of the honour the College has done me by inviting me to give the Goulstonian Lectures for the year 1915.

It is with much diffidence that I approach the task, more particularly on recalling the list of lecturers who have preceded me, and the valuable contributions they have made to medicine, and especially to that branch of it in which my chief interests lie. The conditions under which I have had to work during the past fourteen months have made me realize that the ambition to attain their high level is scarcely practicable; I can only hope that the number of clinical observations I have the opportunity of presenting to you will in some measure atone for the absence of completeness and detailed analysis you will find in my lectures.

And even what I can offer has been possible only through the co-operation and help I received from the medical officers of the numerous hospitals in France, in which the material was collected. I am more especially indebted to my colleague, Lieutenant-Colonel Sargent, in association with whom the work has been done. In addition to operating on many of the cases he took a coequal part in the necessary clinical investigations, and his name must be consequently associated with mine in any observations of value I can place before you.

The chief interest of our observations lies in the symptoms produced by different types of spinal injury and when different portions of the cord are affected, but in order to interpret and estimate their significance correctly it is first necessary to study the nature of the medullary lesions to which they are due. We have been hitherto able to obtain *post-mortem* examinations in about fifty cases, and the histological changes have been studied in a certain number of these.

The spinal cord may be injured directly by the projectile and either completely or incompletely divided, but more commonly it escapes direct damage by the missile and is injured only by displaced fragments of bone, which either compress or lacerate it; frequently, however, it is not injured directly either by the projectile or by indriven fragments of bone, and the structural changes in such cases can be attributed only to the concussion or commotion effects produced in the cord by a missile which has struck some portion of a vertebra. Spinal concussion is most commonly seen when the projectile has touched either a spinous or transverse process, which it may have fractured or not, but it may be also produced by a bullet which penetrates or perforates the body of a vertebra.

It is obvious that this classification into *direct injury*, *contusion*, and *concussion* cannot be exact, as contusing or compressing fragments of bone frequently lacerate the cord as a missile may, while concussion effects are liable to occur whenever the missile strikes the vertebral column, and whether or not it injures the cord directly or drives fragments of bone on to it.

DIRECT INJURIES.

On examining a case in which the spinal cord has been completely divided, we find as a rule a considerable amount of clot and often pieces of bone between its two ends. A few days after the infliction of the wound these are swollen, irregular, and very soft to touch for at least 1 cm. from the point of division; indeed they may be more or less diffuent, and on handling semifluid disintegrated material, frequently stained with blood, which S.

George Makins, from his experiences in the South African war, very accurately described as custard-like, may extrude. After hardening these portions still seem swollen, softened, and oedematous; the outlines and details of their cross section are obscured, and there are often minute or larger haemorrhages within them, chiefly in the grey matter.

Not infrequently the spinal wound is infected, but these appearances are then modified only by the existence of a septic meningitis which may spread rapidly upwards and downwards from the lesion. It occasionally happens, however, that haemorrhages and early adhesions between the arachnoid and dura, and in the subarachnoid space limit the infection to the wound. Subdural haemorrhages of considerable size also occur, but they are rarely sufficiently large to compress the spinal cord. Haemorrhages, which are, however, generally small and insignificant, are more common in the soft meninges.

Microscopical examination always shows that there are severe and relatively extensive changes in the spinal cord immediately above and below the lesion; for the distance of half a segment at least and often further the tissue is completely softened and none of its normal elements are recognizable. Under the microscope only ruptured disintegrating axis cylinders and globules and irregular masses of myeline can be seen; the latter rapidly become smaller and smaller and are absorbed by the numerous scavenger cells which quickly pervade the tissue. Haemorrhages, generally small but occasionally of considerable size, are usually found in these softened areas, but they are more common and, as a rule, larger in the grey than in the white matter. In many cases, however, haemorrhages have been rare or very small, and they are evidently not an essential feature of the condition. The blood vessels are always engorged, but they are, as a rule, thrombosed only in the immediate neighbourhood of the primary wound.

The most striking feature in these sections, however, is the presence of large spherical or irregular cells, which are frequently, apart from the blood vessels, the only recognizable tissue elements (Plate, Fig. A).

They have, as a rule, a single round or oval nucleus, larger than that of a lymphocyte but smaller than those of endothelial cells, which is granular and stains darkly, and is usually situated to one side of the cell; two or three nuclei in a cell are not uncommon. The cytoplasm is finely granular and stains deeply in the early stage, but when the cells are fully developed they are usually filled with coarse fatty granules which are derived from the disintegrated myeline. Later they become larger, and more irregular in shape and their cytoplasm frequently contains large vacuoles. The nuclei are now shrunken or deformed, stain more deeply and usually lie immediately on the periphery of the cell. Eventually these cells seem to extrude their nuclei and break up and pour their softened and digested contents into the tissue in which they lie. They are usually smaller and more tightly packed in the grey than in the white matter; when the tissue is severely necrotic they may be found only in the neighbourhood of vessels.

These are obviously compound granule cells, the origin of which has been much discussed; by some they are attributed to connective tissue, endothelial or leucocytic origin, but others believe they are wholly, or in greater part at least, derived from the neuroglia. The study of our specimens certainly supports the latter view, as the rarity of cell infiltrations or proliferations in the neighbourhood of the vessels has been a striking feature in most of our sections, and there has been no evidence that they develop in those regions only in which either connective tissue or endothelial cells are to be found, that is in the vessel walls. On the other hand, one can frequently see in less damaged portions of the tissue that a neuroglial nucleus accumulates cytoplasm around it, which at first takes an irregular or angular shape, but rapidly swells into a spherical element that is identical with an immature granule cell. Further, gradual transitions between neuroglial nuclei and those typical of the granule cells can be frequently seen.

A proliferation of connective tissue elements around the smaller blood vessels also occurs. They take the form of elongated spindle-shaped cells which appear to be engaged in the formation of new capillaries.

These secondary changes, which occur in the neighbourhood of a laceration or division of the cord, evidently produce further destruction of it. They are obviously degenerative rather than inflammatory, and are due to the oedema and circulatory disturbances that occur in the bruised and necrotic tissue on the borders of the injury, spreading into and involving parts which were not directly damaged by the missile. Oedema is the most important factor; it seems to affect the vitality of the neuroglial matrix as well as of the nerve cells and fibres, and combined with circulatory disturbances leads to their disintegration.

Even infection of the wound does not as a rule materially affect these histological changes; leucocytic infiltration is rarely extensive, and when pus cells are found they appear only as an accidental complication in the section. Evidences of infection are always most prominent in the meninges, and it is a striking fact that invasion of even the softened tissues by pus cells which extend along the vascular sheaths is rare.

In addition to these pathological changes in the immediate vicinity of the wound more distant and more irregularly distributed lesions are found in the majority of cases, and often extend over four or five segments in either direction.

Distant Lesions.

Oedema of both grey and white matter with some swelling and softening of the cord is the most constant of these changes. It gradually diminishes away from the wound, and often seems to bear no definite relation to its severity.

Haemorrhages of various sizes are often associated with it, but these are less constant; they are generally small punctiform extravasations of blood, which give a mottled appearance to the cross section of the cord, but they are frequently larger (Plate, Fig. B); a large central haemorrhage with a tendency to spread longitudinally in the cord, such as is generally understood by the term "haematomyelia," was not present in any of the fifteen cases in which the microscopical examination has been completed. These haemorrhages are found particularly in the grey matter and about the central canal; one of the most common sites is the dorsal horn, where they can obviously interrupt the reception of afferent impulses. In the grey matter they are liable to break up and destroy the tissue, but when small they produce surprisingly little change in the white columns, the extravasated cells merely tracking along the vessels or between the fibres; occasionally, however, there is some local softening, and later neuroglial proliferation, around larger extravasations.

The extent of these small disseminated intraspinal haemorrhages is occasionally surprising; in one case they spread over two and a half segments on each side of the wound. They are found with lesions of all regions of the cord, but they are usually most prominent when the cervical region is wounded and probably least so with injuries of the lower dorsal and lumbar segments. Owing to the relatively slight destruction they produce in the tissues their importance in the production of clinical symptoms may be easily over-estimated.

Interesting changes which cannot be directly connected with the trauma or with the occurrence of these small disseminated haemorrhages are frequently found in the nerve cells and fibres at a considerable distance from the wound. The most striking of these is swelling of the axis cylinders. Frequently all those in a considerable area of the white matter are found slightly swollen in the neighbourhood of the wound where the tissue is oedematous, but a more unexpected condition is the relatively enormous increase in size of either isolated axones or of groups of axones, unassociated with any other obvious changes in the tissue, and often at a considerable

distance from the injury (Plate, Fig. C); in fact, this is usually the most far-spreading lesion recognizable, and may be often seen throughout the four or five segments adjacent to the lesion. The swollen axis cylinders may be in either ascending or descending tracts, and there is no evidence to suggest that the change is related to a secondary degeneration of the fibres. They are most commonly seen in the white columns, but also occur in the grey matter and in the dorsal roots, generally, however, in proximity to the wound. When the increase in size is moderate the swollen fibres appear circular in cross section and either granular or homogeneous, but as soon as they become larger their cross section is often oval or irregular, and not infrequently they contain more darkly staining central cores.

In longitudinal sections the swollen fibres may appear of almost uniform calibre for a certain distance, but they are more commonly moniliform or bead-like; in such fibres Bielschowsky preparations frequently show that the component fibrils of the axis cylinders are separated in the swollen parts by a great increase of the interfibrillar substance (Plate, Fig. D). The most swollen fibres may be frequently seen degenerating and undergoing disintegration; some apparently break up into a granular debris, others become vacuolated and are occasionally invaded by small phagocytic cells (Plate, Fig. E).

In sections stained by haematoxylin a thin irregular ring may be generally seen around them, which, when stained by Weigert's method, is obviously the myeline sheath. This is occasionally ruptured, so that an incomplete ring surrounds the axone; and in longitudinal sections it is often found broken and discontinuous. In other parts, especially when oedematous, the sheaths are swollen and irregular in outline and calibre, and they are frequently disintegrated into globules of myeline, especially in the neighbourhood of the lesion. In more distant regions, however, apart from any other

pathological alteration, the myeline sheaths of isolated or small groups of fibres are found in parenchymatous degeneration.

Two types of change are seen in the nerve cells of the grey matter: in the region of the wound they are often shrunken, stain darkly, and present none of the normal details, or if the tissue is oedematous they may be swollen. Some of them at least break up rapidly, often owing to invasion by neurophages, but when less severe these changes are evidently recoverable. A more common type of change is chromatolysis, with some swelling of the cell body, eccentricity of the nucleus and disappearance of the Nissl bodies, especially from its centre. Bielschowsky preparations often show irregular swelling of the dendrites of these cells.

In addition to these changes in the cells and fibres, irregular focal lesions in which all elements of the tissue may be involved often occur for a considerable distance from the wound, sometimes as far as four or five segments.

Occasionally there are small circumscribed patches of necrosis in which both the fibres and the neuroglial matrix are involved, or in which only the larger strands or neuroglia persist, without any evidence of haemorrhage or inflammatory reaction around them (Plate, Figs. F, G); there is, in fact, simple uncomplicated necrosis of the tissue, generally accompanied by considerable oedema. The vessels may be slightly engorged, but as they are rarely thrombosed the necrosis cannot be attributed to vascular lesions; further, they are generally in the periphery of the white matter in the region in which its blood supply by the pial vessels is most complete, and neither in distribution or shape are they such as might be expected from vascular lesions.

These focal necroses are generally smaller in the grey matter, and when not complete produce only a rarefaction

DESCRIPTION OF SPECIAL PLATE.

Fig. A.—Granule cells in an area of complete softening. No normal tissue is recognizable in the section.

Fig. B.—Multiple haemorrhages in both grey and white matter, two segments above a complete division of the cord.

Fig. C.—A cross section of swollen axis cylinders in one lateral column two and a half segments above the level of the maximal damage produced by concussion. There was no other abnormality in the white matter here.

Fig. D.—Swollen axis cylinders in longitudinal section stained by Bielschowsky's method.

Fig. E.—Swollen axis cylinders, lying in round or oval spaces, which are undergoing vacuolation and disintegration.

Fig. F.—Irregular foci of softening and necrosis in the white matter two segments above a complete division of the cord.

DR. GORDON HOLMES: SPINAL INJURIES OF WARFARE.

FIG. A.

FIG. B.

FIG. C.

FIG. D.

FIG. E.

FIG. F.

DR. GORDON HOLMES: SPINAL INJURIES OF WARFARE.

FIG. G.

FIG. H.

FIG. J.

FIG. K.

FIG. L.

FIG. M.

of the matrix with severe degenerative changes in the nerve cells. In two cases, however, there was extensive necrosis of the ventral horns and the adjacent parts of the ventral columns, with disappearance of or degenerative changes in the nerve cells and fibres, and invasion of the affected area by granule cells.

Less complete and more irregular lesions of the white matter, in which all fibres are not completely destroyed, are more common, and were, in fact, present in all the cases in which a histological examination has been made.

The periphery of the cord is most frequently involved, and the foci have often an irregular wedge shape, with the base on the periphery and the apex directed centralwards. They are found most frequently in the dorsal and lateral columns, more rarely on the ventral surface, except perhaps in the angle between the ventral periphery and the ventral mesial fissure.

In these affected areas only the coarser neuroglial trabeculae and the dense subpial layer persist, and in section the tissue has consequently a sieve-like or reticular appearance (Plate, Fig. H); in early cases the neuroglia in the process of disintegration may be recognized by its characteristic colour in van Gieson preparations. There is usually very little evidence of oedema in or around the foci, and a coagulated transudate can rarely be seen within the meshes. There is never any cellular infiltration, or at the most only an insignificant increase in the number of free cells in the sheaths of the vessels. A considerable neuroglial proliferation is often visible around the foci, and spider cells can be frequently seen here and in the persisting trabeculae.

The majority of the meshes are often empty, but they frequently contain distended or disintegrated myeline sheaths and axis cylinders, which are often enormously swollen. It is, however, a striking fact that relatively normal axis cylinders often persist within them; indeed, these seem to escape more frequently than any other part of the tissue. Other meshes contain large granule cells, which are often loaded with fat and debris.

This change is sometimes more general and diffuse, and may affect the greater part of the periphery of the cord, producing either the appearance of confluent foci or a general sieve-like rarefaction of the tissue. It varies also in degree; in places only the larger glial strands persist with disintegrated detritus between them. Occasionally there are traces of haemorrhage into these areas, but they are rare.

Frequently, however, the only abnormality in a part of the cross section of the cord is the presence of more or less circular vacuoles, either isolated or groups, which may be either empty or contain granule cells or a swollen axis cylinder (Plate, Fig. J). The tissues around generally appear compressed excentrically from the vacuole, so as to suggest that the latter was produced by some distensible force. Owing to the loose and porous structure of the neuroglia this cannot have been due to an accumulation of fluid within it, and in sections they rarely contain a coagulum. On the other hand, a careful examination of the development of these vacuoles shows that they occupy the site of a single myelinated fibre, and the frequency with which swollen fibres are found within them suggests that they are generally due to the swelling of an axis cylinder or of its sheath. In other instances they are obviously filled by granule cells, but this may be due to secondary invasion.

In a surprisingly large proportion of the cases, in fact in more than half those which have been examined in microscopical sections, and apparently in an equal proportion of those in which only a naked eye examination has yet been made, there were curious cavities, roughly cylindrical or oval in cross section, in the segments of the spinal cord

adjoining the lesion, either in those above or below it, or in both (Plate, Figs. K and L). These often extended a considerable distance, in some cases over four or five segments, and at their ends furthest from the wound they were occasionally the only indication of any abnormality. In three cases they were found on each side of the primary injury, and in four cases only below it. In one cord the cavity lay in and occupied the greater part of one dorsal horn (Plate, Fig. M), but with this exception they were always found in the dorsal columns, generally to one side of the middle line; they did not involve the grey matter or approach the central canal. Their size was always considerable, the greatest diameter usually varying between a quarter and half the dorso-ventral breadth of the dorsal columns. They were consequently distinctly visible to the naked eye when cut across, and generally contained greyish or brownish gelatinous matter, sometimes tinted with blood.

Their contents on microscopical examination are seen to consist of broken up and degenerating fibres and disintegrated neuroglia, which are not, except in cases of long duration, quite necrotic, though shrunken; darkly staining, pyknotic nuclei and degenerating granule cells were generally the only recognizable elements. The latter were usually laden with fat. A certain number of red cells could be seen in many cavities, but excepting that in the dorsal horn none contained much blood.

In some cases this disintegrated matter filled the cavity,

but in most instances, after fixation and the preparations of sections at least, the cavities were partly empty. There can be no doubt that in every case the contents during life existed under pressure, as the unaffected parts around them were compressed, and the dorsal mesial septum was generally displaced towards the opposite side when the cavity lay in one dorsal column only (Plate, Fig. K).

The cavities were, as a rule, very sharply delimited from the tissues around them, which were often quite intact, but in places there was occasionally an irregular softening

on their margins. In certain cavities the degenerating material in them came into direct contact with, or had simply sloughed off from, the normal tissue, but the walls were frequently formed by one or more layers of granule cells, the outer of which remained well developed, while the inner degenerated and added to the disorganized contents.

When a cavity is examined in serial sections it is usually found that it becomes larger away from the lesion and terminates in a rounded or obtuse apex; its other end rarely extends down to the primary lesion, though it is generally connected with this by a haemorrhage or a small track of softening, or in a few instances by an apparently artificial slit, which might be attributed to retraction and shrinkage of the oedematous tissues. There was as a rule very little vascular disturbance around the cavities, apart from the oedema which spreads from the neighbourhood of the injury; haemorrhages were not as a rule found in or around them, and apparently bore no causal relation to their development. In one case a vessel which passed horizontally through a cavity was thrombosed, but this seemed to be a secondary effect, and due to pressure or tension upon it.

Although the cavities contained disintegrated myeline and swollen and disrupted axis cylinders, an examination shows that in relation to their size they involve but little functional tissue; their chief effect must be consequently due to the pressure they exert upon, and the distortion they produce of, the unaffected fibres in the neighbourhood.

Either the dorsal or ventral roots or both are frequently injured by the missile or by displaced fragments of bone, but apart from this considerable changes are often found

DESCRIPTION OF SPECIAL PLATE.

Fig. G.—A patch of complete necrosis in the left lateral column three segments above the level at which the cord was lacerated by a bullet.

Fig. H.—Sieve-like rarefaction and vacuolation of the periphery of one lateral column two segments above the level of the maximal injury produced by concussion.

Fig. J.—A central softening in the cord with small foci of softening in the periphery of the white matter, and extensive vacuolation.

Fig. K.—A cavity filled with necrotic material in the left dorsal column three segments below the level of maximum injury produced by concussion.

Fig. L.—A cavity in the left dorsal column three segments above a direct injury to the cord. It contains only necrotic detritus. There are also a patch of complete necrosis in the left lateral column and areas of partial softening in the rest of the white matter.

Fig. M.—A descending cavity in the left dorsal horn two and a half segments below a complete division of the cord.

in them, and especially in the dorsal roots near their entry into the cord. Their neuroglial portions are usually swollen and oedematous, and their myelinated sheaths are distended or partially disintegrated. The proportion of fibres actually broken across is usually insignificant, and the change does not usually extend beyond the line of the glial and connective tissue junction. These changes in the dorsal roots may have some significance in relation to certain clinical phenomena we will describe later.

These spinal changes, associated with a direct injury of the cord, may be illustrated by one typical case:

CASE I.

Pte. C., 198, was wounded by a rifle bullet on November 27th, 1914. He lost power in his legs at once and developed retention of urine. He had had haemoptysis at first, but spat up blood-stained sputum only when he was admitted to the base hospital ten days later. He also complained of a stinging girdle pain just above the level of his umbilicus.

The bullet entered 6 cm. to the left of the tenth dorsal spine and made its exit in the fifth right intercostal space in the mid-axillary line. He had a right-sided haemothorax, cystitis, an early bedsore, and trophic blisters on the front of his ankles and on the inner sides of his knees. His legs were flaccid and paralysed, and no reflex movement could be obtained on stimulating his soles. The abdominal muscles were paralysed to midway between the umbilicus and the xiphoid, and the lower four intercostals did not contract on either side. The knee and ankle jerks, as well as the plantar and the lower abdominal reflexes, were absent, and all forms of sensation were completely lost to 8 cm. above the umbilicus.

He died eighteen days after the infliction of the wound. A small right-sided haemothorax was found in addition to the spinal injury. The bullet had passed through the arch of the tenth dorsal vertebra, lacerating both the dura mater and the cord; the latter, after hardening in formalin, was found to be completely destroyed between the upper part of the twelfth dorsal and the middle of the third lumbar segment. There was some meningitis both above and below the injury.

Below the Lesion.—In the fourth lumbar segment the dorsal columns were almost completely destroyed, as well as the inner margin of the left dorsal horn, and the tissue around was softened and infiltrated by granule cells (Fig. 1). The grey matter was oedematous, and its nerve cells were in advanced chromatolysis. The greater part of the right lateral column was necrotic and pervaded by haemorrhages, while the rest of the white matter presented a sieve-like appearance, and contained swollen axis cylinders, granule cells, and proliferating spider cells. From the fifth lumbar segment a large cavity, which contained disorganized fibres and neuroglia as well as granule cells, extended through the dorsal columns downwards into the conus. The remaining portions of the dorsal columns were oedematous and vacuolated or sieve-like in the fifth lumbar, but normal in the sacral segments. The white matter of the ventro-lateral columns was also vacuolated, but showed only a slight loss of fibres below the first sacral segment; while the grey matter was oedematous and all its cells were swollen and in advanced chromatolysis.

Above the Lesion.—As high as the upper part of the eighth dorsal segment there were irregular patches of necrosis and sieve-like rarefaction of the tissue, in which only the coarser neuroglial strands and a certain number of axis cylinders, normal or swollen, persisted (Fig. 1); there was also some infiltration by granule cells and minute haemorrhages. It was chiefly the

peripheral portions of the lateral and dorsal columns that were involved. A small softening, in which there was an apparently artificial slit, extended upwards in the left dorsal column through the eleventh, tenth, and the lower part of the ninth dorsal segments, and in the upper portion of the latter became continuous with a large cavity which reached the upper margin of the eighth segment; it was sharply defined from the surrounding normal tissue and contained disintegrated debris and degenerating granule cells.

It may be noted that, although the clinical symptoms—the reflex changes as well as the motor and sensory paralysis—indicated a lesion of the ninth dorsal segment, the twelfth was the highest segment in which the spinal cord was actually damaged by the wound; the histological examination, however, explains the discrepancy.

As the immediate and more distant effects of a spinal wound have been described, the anatomical changes produced by compression or contusion, and those due to concussion, will require less time, as in both these classes we

again meet the histological lesions which we have already found associated with injuries produced directly by the missile.

CONTUSION OR COMPRESSION OF THE SPINAL CORD.

When a portion of a vertebra or a detached spicule of bone is driven into the spinal canal, it frequently lacerates both the cord and the theca, and causes lesions which may differ only in degree from those produced directly by a projectile. Frequently, however, there is no obvious external injury to the cord and the dura mater is not torn, even though the clinical symptoms indicated a complete transverse lesion. Small haemorrhages into the meninges are, however, common, and on palpation the cord at the level of the contusion is soft, and if the pia mater is incised or pricked semifluid custard-like material may escape.

Fig. 1.

When the injury is less severe the normal appearance of the cross section is only obscured, and there are frequently minute haemorrhages throughout it. The damaged area and the segments on either side of it are usually swollen by oedema, and the cord may be indented by the indriven bone. When the lesion is examined under the microscope changes are found very similar to those in the parts adjoining a direct injury, but their intensity naturally varies much. The most important is softening and disintegration of the tissues, always greatest in the region which was directly contused. If the injury is severe the whole cross section may be softened, but more commonly there are discontinuous foci in the ventro-lateral and dorsal columns. The grey matter may be also completely destroyed in whole or parts, but it is more usually extremely oedematous and only partly disintegrated, with its nerve cells necrotic or in advanced chromatolysis. Occasionally only a diffuse or focal necrosis is found in which no stainable elements persist. The affected areas are gradually invaded by granule cells, and a considerable proliferation of neuroglial cells is visible around them, but, apart from a pronounced congestion of the vessels and an occasional increase of the

cells in their walls, there is no evidence of any inflammatory process. The amount of haemorrhage into the injured region also varies very much.

The distant lesions differ in no respect from those which are so commonly associated with direct spinal injuries. Diffuse foci of necrosis and softening, vacuolation due to falling out of fibres and the sieve-like rarefaction produced by the disappearance of fibres and of the finer glial matrix, as well as minute scattered haemorrhages, are found in the white matter of the adjoining segments, while the grey matter is also oedematous, and contains similar haemorrhages and perhaps foci of softening. The central cylindrical cavities, which have been already described, also occurred in the dorsal columns relatively as frequently as with direct injuries.

CONCUSSION OF THE SPINAL CORD.

In cases of concussion, when the cord is not damaged by the fracture or dislocation of a vertebra, there may be no external signs of injury or only a more or less uniform swelling opposite the site of impact, and even to touch no definite abnormality may be recognizable.

On microscopical examination the vessels are found engorged, and there are generally punctiform haemorrhages, especially in the grey matter. The most striking change, however, is the oedematous swelling of the most affected segments with either diffuse or focal necrosis and softening, which, at least in the cases that have been examined microscopically, has been most pronounced in that part of the cross-section nearest the point of impact. In these areas there may be complete destruction of the functional elements, but more usually only a proportion of the fibres have disappeared, while the myeline sheaths and axis cylinders of others are swollen. Focal softenings also occur in the grey matter, but are usually unrelated to the haemorrhages which this frequently contains. There is, often, however, some softening and disintegration of the tissue, as well as degeneration of the nerve cells, around these haemorrhages.

The distant lesions in cases of concussion are similar to those found associated with direct and contusion injuries, but they are often very marked in relation to the changes found at the site of maximum damage. Scattered haemorrhages, irregular foci of necrosis, and softening and cavity formation occur, but the most pronounced feature is the extensive parenchymatous changes that often extend over four or five segments in either direction. These consist in the swelling of fibres, either isolated or in groups, in the midst of tissue which is otherwise normal or only slightly oedematous; as a rule the axis cylinder is more swollen and the myeline sheath surrounds it as a distended and attenuated ring, but in places the sheaths are more affected, and are often broken up. The disappearance of these swollen fibres and the partial disintegration of the neuroglia leave vacuoles and round or oval cavities in the white matter and give it a reticular or sieve-like appearance. There are also often considerable histological alterations, either degenerative or chromatolytic, of the nerve cells at some distance from the level of the injury, but these are found most commonly where the tissue is oedematous.

If these distant and diffuse lesions are compared with those found at the point of impact it will be seen that they differ from them only in degree, and as they occur whether the cord has been directly injured or not by the projectile or by indriven bone, it is probable that in each case they are due to the same cause, which we must seek in the effects of concussion of the spinal cord produced by a sudden violent impact on the vertebral column.

We must now consider shortly the nature of these changes which have been described, the causes to which they are due, and their significance in the production of the clinical symptoms that characterize these cases. The conclusions and views expressed must not be regarded as final, as there is still much material to be investigated, and the amount of time which could be allotted to the histological preparations which are already available has been necessarily very limited.

Spinal Concussion.

The most important and obscure factor is that which we understand by *concussion*—that is, functional or

anatomical disturbances produced indirectly in the spinal cord by a sudden and violent impact on the vertebral column. The nature of spinal concussion has been much discussed, and it has been, in fact, questioned if spinal lesions, such as those described above, do occur apart from temporary dislocation or fracture of a vertebra, compression by fragments of bone, or an extra or subdural haemorrhage causing direct trauma to the cord.

But many cases in which none of these possible causes existed have been recorded, and we have had the opportunity of observing cases in which they could not be demonstrated. Further, certain of the distant lesions that we have already described, which often extend over several segments on either side of the primary injury, or beyond the level of the impact on the vertebral column, cannot be due solely to a direct trauma, and they are identical in nature to those attributed to concussion. The changes are irregular focal softenings or patches of necrosis, sieve-like vacuolation of the white matter, disseminated haemorrhages and local lesions of the myelinated fibres.

It is in the first place necessary to insist that these changes are not, as oedema may be, continuous with those in the neighbourhood of the spinal wound; the haemorrhages, for instance, are usually discrete and are often only minute extravasations of blood in the Virchow-Robin or perivascular spaces, while the focal necroses and softenings do not, as a rule, spread longitudinally in the cord from the region of the trauma. Further, when a group of fibres is affected, the lesion is usually focal, and it is not necessarily those of one tract only that are involved; it cannot consequently be either a manifestation of a secondary degeneration, or of a pathological change which has spread from the point at which the fibres were directly injured.

The special character of these lesions is, therefore, their diffuse and irregular distribution and their tendency to diminish gradually from the point of maximal disturbance. They are not due to haemorrhages, as these bear no constant relation to them, and as vascular occlusions are also rare these obviously cannot be the main causal factor. On the other hand, as the chief lesions are foci of primary necrosis and parenchymatous change in the cells and nerve fibres, the essential changes may be described as primary disturbances in the vitality of certain portions of the tissue, associated with oedema and frequently with small scattered haemorrhages.

It is difficult to offer a complete and satisfactory explanation of how a blow on the vertebral column can produce these lesions in the cord, protected as it is within the canal. Certain structural alterations found in cerebral concussion are attributed to the violent oscillations produced in the cerebro-spinal fluid, especially in that of the ventricles, but the spinal cord is only surrounded by fluid, and is able to swing to some extent within the dural sac with its oscillations. The waves of pressure thus set up may, however, produce physical effects within the cord, and possibly disturbance of its lymph circulation, but the most probable explanation is that put forward by Fickler, according to which the cord is made to oscillate within the canal by the impact on the vertebral column, and as its movements will obviously not be synchronous with those of the column it may be directly bruised against the walls of the canal, while at the same time the sudden jarring of the cord produces a physical disturbance in its tissues, and especially in the fluid axoplasm of its fibres. There is much in favour of this explanation, and the factors it hypothecates cannot be neglected. The histological changes in the spinal roots may be, in fact, partly due to the strain thrown upon them by the displacement of the cord. But if it is the whole explanation, the structural lesions should be greatest at the position of *contre coup*; we have, however, so far always found them most pronounced immediately under the site of impact. Further, it must be remembered that the spinal roots and the ligamentum denticulatum limit the movement of the cord within the canal, and in many levels at least must make *contre coup* bruising impossible.

Whatever may be the exact mechanism of spinal concussion, it must be admitted that a sudden violent impact on the vertebral column can produce diffuse, irregular, and severe structural changes within the spinal cord. The factors which determine the severity of these lesions must

be the momentum of the projectile, the part and surface area of the vertebra which it strikes, and the region of the spine which is wounded.

Secondary Changes.

Finally, we must consider the secondary changes which may occur in the neighbourhood of the primary injury and in those portions of the cord which have suffered from concussion.

It has occasionally happened that when the wound is not severe the patient was at first able to perform some movements in his legs, but lost the power to do so within the following two or three days. Further, we have seen new symptoms develop or their level alter under observation. This may be due to a secondary haemorrhage into the affected tissues or to progressive softening, of both of which we found possible evidence in our sections.

But the most striking secondary change was the development of the cylindrical cavities we have described. Their exact pathogenesis is obscure, but certain features they present must be emphasized in attempting an explanation. In the first place, they seem to involve the destruction of very little tissue, but rather separate and compress the fibres around them; their contents must be consequently under much pressure. In the second place they evidently develop away from the lesion, as the material they contain is always less necrotic at their upper or lower end if they are respectively ascending or descending cavities; and finally they do not as a rule extend to the maximal lesion, but are generally connected with this by a track of oedematous or softened tissue, or by a narrow channel of softening or by a fissure (Fig. 1). It seems, therefore, probable that they originate from the accumulation under pressure of transuded fluid and degeneration products in a small projection of the primary lesion, which tracks upwards or downwards along the lines of least resistance through either normal or oedematous parts, destroying only a relatively small amount of tissue but increasing in size probably under the same principles as a retention cyst. The granule cells which frequently line their walls, or are contained within them, must be due to a reactionary proliferation of the neuroglia in the tissue through which they track.

Finally, it must be emphasized that they are not due to infection, as in several cases in which they occurred the theca had not been lacerated, and there was no sign of infection in either the cord or the meninges.

They are obviously not produced by haemorrhages, though a certain number of red blood cells may be found within them, nor by vascular lesions, as there has been no evidence of these, and the position of the cavities does not correspond with the distribution of any spinal vessel. The frequency with which they occupy the ventral portion of one or both dorsal columns is striking, but its significance is not clear. This region is, however, a watershed area between the distribution of the anterior spinal arteries and of the small arteries that supply the dorsal columns, and as such may have a relatively poorer blood supply than other parts of the cord. In many cases, too, the dorsal columns seem to suffer more severely with oedema and softening than the ventro-lateral columns.

It is obvious that these irregular and diffuse changes which are found in various types of spinal injury must be taken into account in interpreting and estimating the significance of the clinical symptoms that are observed in these cases. The following conclusions at least can be drawn:

1. The structural lesions in the spinal injuries of warfare are rarely sharply limited or circumscribed, and cannot be compared to those produced experimentally in a physiological laboratory. The level of the lesion, as indicated by the clinical symptoms, for instance, often does not correspond with the level of maximal damage.

2. The lesions are so irregular in distribution and severity when the spinal injury is not complete that much care is necessary in drawing conclusions from the clinical symptoms alone on the functions of parts which it may be assumed have been involved.

3. Secondary changes may occur later in the cord which can alter or modify the clinical symptoms.

MALIGNANT TUMOUR OF THE KIDNEY.

AN ANALYSIS OF SOME SYMPTOMS AND SIGNS OF A SERIES OF CASES.

By R. J. WILLAN, M.S., F.R.C.S.,

HONORARY ASSISTANT SURGEON ROYAL VICTORIA INFIRMARY, NEWCASTLE-ON-TYNE; STAFF SURGEON R.N.V.R., ATTACHED TO THE ROYAL NAVAL HOSPITAL SHIP "DRINA."

THIS article, written from rough notes made some time ago, is divided into four portions. Part A contains an analysis of age, sex, side, and the onset symptom. Part B deals with the association of pain, haematuria, and a palpable renal tumour to each other during the illness. Part C mentions some complications, and Part D includes illustrative clinical cases. The material is extracted from my own case notes; I am also indebted to my colleagues on the staff of the Royal Victoria Infirmary, Newcastle-on-Tyne, for the use of their records.

A.—AN ANALYSIS OF AGE, SEX, SIDE, AND THE ONSET SYMPTOM.

1. Age.

Ten out of the series occurred in children under the age of 10; in each case the condition was sarcoma. No case occurred between the ages of 10 and 29 inclusive—that is, the second and third decades of life; 31 out of the 59 cases occurred between the ages of 45 and 65; or, if children are excluded, and only the 49 adults reckoned, 31 out of the 49 occurred between these ages—that is, 63.2 per cent.

A curious fact, which may only be a coincidence, is that in the fourth, fifth, sixth, and seventh decades the preponderance of cases occurred in the first half of the decade, for 5 cases were between 30 and 35 years old, 7 of the cases were between 40 and 45, 10 of the 17 cases were between 50 and 55, while 9 of the 12 cases were between 60 and 65.

Of the 49 adult cases, 14 occurred between the ages of 45 and 55, a percentage of 28.5.

TABLE I.

Under 9.	10-19	20-29	30-39	40-49	50-59	60-69	70-79	Not Stated.
10	—	—	5	11	17	12	1	3

Total, 59.

2. Sex.

63.8 per cent. of the series were males, there being 36 against 23 females. Of the 10 children, 6 were boys.

3. Side.

In 31 instances the malignant growth affected the left side and in 26 the right side. In 2 cases the side affected was not stated.

4. Onset Symptom.

In 7 cases of the series the actual onset symptom is not stated; these will therefore not be taken into account in calculating percentages. Of the remaining 52 cases, the commonest initial evidence was painless haematuria. This was observed in 14 cases.

In 10 cases the illness was ushered in by acute renal colic, while in 10 other cases a painless swelling found in the flank was the first evidence of anything wrong; 9 cases began with chronic pain in the flank, while 5 patients suffered from cachexia before any sign of malignant growth of the kidney became manifest. In 2 cases there was increased frequency of micturition at onset, another went to his doctor for a varicocele, while the remaining one had "swelling of the bowels."

Therefore the separate symptoms of (1) painless haematuria, (2) symptomless flank swelling, (3) acute renal colic, and (4) chronic renal pain, account for the onset symptoms in 43 cases, a percentage of 82 per cent.

Of the 7 children included in the above, in 3 cases a painless tumour was found in the flank, in 2 cases there was painless haematuria, in 1 case a chronic aching pain, in 1 case cachexia; while in the 3 further cases there was no statement.