patients, but invariably hand it to someone who is doing it frequently.

A blood sugar test may be necessary in some of the cases; in others it will be advisable. Some may decide to carry on without having the test done. It seems to me that it would be worth while for one of the younger or less busy practitioners in each of the two larger towns of the division to make himself efficient in it, preferably by first doing it under the supervision of some expert, and then his knowledge might, by arrangement, be made available for his fellow practitioners.

There is no quick and easy way for the treatment of diabetes. Unless some of the fundamental conceptions of the disease are appreciated it is impossible to treat it rationally. Insulin has come as a great aid, but it has not replaced the need for that close attention to diet which has been the foundation of any success that has attended treatment in the past. Added to dietetic treatment insulin has improved the outlook for the diabetic patient. It will give him a sense of well-being which he did not obtain under the older methods, it will make his diet more palatable, it will increase his physical efficiency, it may avert coma, and it will prolong life, but it is not devoid of danger, and it is not a cure for diabetes mellitus.

DIABETES AND INSULIN.*

BY

J. A. NIXON, C.M.G., M.D., F.R.C.P.,

PHYSICIAN, BRISTOL ROYAL INFIRMARY.

The Provincial Board of Health, Ontario, has made arrangements to supply insulin free to physicians for patients in Ontario who are unable to pay for the drug themselves, and has issued a pamphlet which contains many valuable instructions. The following points of importance are worth noting:

1. The majority of patients suffering from diabetes do not require insulin and can better be treated by dietetic measures alone.
2. In order to decide whether insulin is necessary the patient should, wherever possible, be placed in hospital for three weeks to determine the severity of the disease.
3. The following work should be done in hospital:
 - (a) Estimation of the quantity of urine passed in twenty-four hours.
 - (b) Examination of urine and quantitative determination of sugar in urine.
 - (c) Blood sugar determinations.

The patient should then be placed on a weighed maintenance diet calculated from his basal metabolic rate, and examinations repeated daily to determine the power of carbohydrate utilization. Then if insulin be required the amount and time of the dosage should be scientifically worked out.

While in hospital the patient should be taught:

- (a) How to weigh and prepare his diet.
- (b) How to examine his urine.
- (c) The symptoms and danger signals of his disease.
- (d) How to administer insulin to himself.
- (e) The symptoms and treatment of overdose of insulin.

It is pointed out, further, that insulin is not a cure for diabetes. It is not a substitute for dietetic treatment; indeed, dietetic treatment in cases requiring insulin must be more rigidly, more carefully, enforced than in most cases requiring dietetic treatment only.

Method of Use.

The one danger to be avoided in the administration of insulin is the production of hypoglycaemia—the lowering of the blood sugar level below the normal (100 mg. per 100 c.cm.). It must never be given to any patient until it has been ascertained that there is an actual excess of sugar in the blood. Glycosuria does not in every instance mean that a person has diabetes. In the condition known as "renal glycosuria" the kidneys become abnormally pervious to sugar, so that glycosuria occurs although the blood contains less than the normal amount of sugar.

The first step in investigating a case of glycosuria, there-

fore, is to determine the amount of sugar in the blood. If the blood sugar (collected four hours or more after a meal) on an ordinary diet amounts to 200 mg. per 100 c.cm., the procedure I adopt is as follows:

Macleod's Trial Diet.¹

The patient is placed on a trial diet consisting of two parts: A, containing no carbohydrates, and B, an all-starchy part. A consists of eggs, fish, green vegetables, and fat; B of 100 grams of white bread, distributed between breakfast and dinner. This diet is continued until the glycosuria disappears or reaches a constant level. If it disappears the case is to be classified as mild, provided the blood sugar level is below 200 mg. The daily amount of bread may then be increased by 50 grams a day until sugar reappears; then three-fourths of the limit reached should be taken. A patient who can tolerate 100 grams of white bread (or 50 grams of glucose) does not need insulin, provided that the blood sugar is within normal limits when the glycosuria disappears. If 100 grams of bread cannot be tolerated the allowance must be brought down to 50 grams (or 25 grams of glucose) and if after three days the urine is not sugar-free (with normal blood sugar) the case is classified as severe and insulin may be given.

INSULIN TREATMENT.

The patient is placed on a "basal requirement diet" or the minimum diet adequate for the individual under treatment. Banting gives this as being 3/4 gram of protein per kilogram of body weight and a total of calories calculated according to the patient's height, weight, and basal metabolism. A rough estimate allows from 30 to 35 calories per kilogram of body weight.

In practice Woodyatt's² diet forms an excellent working basis; it contains 50 grams of protein and has a total value of 1,679 calories. It is adequate for a person weighing 140 lb. or 10 st. Growing children need relatively more protein than adults. Carbohydrate in the proportion of 1 gram of carbohydrate for every 1.5 grams of fat is required to enable the fat to be "burnt up." If fat is in excess in the diet ketosis occurs. Woodyatt has evolved a simple formula for regulating proportions in the diet:

Fat should equal the sum of twice the carbohydrate plus half the protein ($F. = 2C. + 0.5P.$).

This formula takes into consideration a fact not always borne in mind—that protein contains a large amount of potential glucose. The total potential glucose in a diet equals all the carbohydrate plus half the protein and one-tenth of the fat ($G. = C. + 0.58P. + 0.1F.$).

Reckoning according to these formulae, Woodyatt's basal requirement diet of 1,679 calories may be made up of: fat 139 grams, carbohydrate 57 grams, protein 50 grams. Since each gram of fat yields 9 calories and each gram of carbohydrate and of protein 4 calories, in order to calculate the number of calories in this diet we proceed as follows:

Fat	...	...	...	...	139 × 9 =	1251
Carbohydrate	...	...	...	...	57 × 4 =	228
Protein	...	...	...	...	50 × 4 =	200
Total	...	...	...	...		1679

These figures are easily remembered and from them any total number of calories can be reckoned for a smaller or larger diet. In order to reduce or increase the diet we may subtract or add gram by gram from the fat, protein, and carbohydrate until the necessary total of calories is reached. In ordering the weights we can convert grams to ounces by allowing 30 grams to 1 ounce.

The basal diet having been settled it is never reduced. Insulin is given in sufficient quantity to enable the patient to utilize the diet. It is advisable to administer the insulin in two doses a day as its effect is short lasting: one dose in the morning and another in the evening seems the best plan.

I commence with a dose of 5 units of insulin. It must be administered by hypodermic injection; by mouth it is inert. It should be given before a meal containing carbohydrate—Banting advises one hour before the meal—because carbohydrate in the food may stimulate the islet cells of the

* The substance of a paper read before the Bath and Bristol Branch of the British Medical Association.

pancreas before the insulin has had time to exert its full effect. If the dose of insulin is too great, symptoms of hypoglycaemia may develop; these are faintness, sweating, tremor, short shallow respirations, accompanied occasionally by nervousness, anxiety, emotional and mental disturbances, and disorders of speech, with, finally, delirium and coma. The antidote for these symptoms should always be at hand on beginning insulin treatment—namely, a sterile 5 per cent. solution of glucose. This may be given by mouth (1 pint at the earliest appearance of any of the symptoms), or if collapse ensues rapidly, by intravenous injection. Kellogg estimates that 5 units of insulin will increase sugar tolerance 10 or 11 grams.

In estimating the proper dosage of insulin it is imperative to determine the effect on the blood sugar. Glycosuria may disappear while the blood sugar still remains at a dangerously high level. The best time for determining the blood sugar is three hours after a meal. The safe level is 100 grams per 100 c.cm. Below this it is not wise to go; a rise above 150 indicates that the dose of insulin is insufficient. The dose must be increased by 5 units at a time until a satisfactory blood sugar level is obtained.

I begin with 5 units twice a day, then increase to 10 units in the morning and 5 in the evening; if this is not enough, then 10 units night and morning, and so on, until the blood sugar is 130. When a dosage of insulin sufficient to enable the patient to metabolize the minimal diet has been found, the diet may be gradually increased. Woodyatt's diet is only adequate to maintain equilibrium in a person at rest. If he is to do a day's work the diet must be raised.

Regulation of Diet.

Starting from Woodyatt's "basal requirement diet" the problem of increasing the food may be made comparatively simple.

The total calorie content of Woodyatt's diet can be reckoned as 1,680, made up thus:

Fat	...	...	...	140	grams
Carbohydrate	...	...	...	57	"
Protein	...	...	...	50	"

Successive increases of one-eighth of each constituent will allow of 210 calories being added:

Fat	...	...	...	17.5	$\times 9 = 157.5$
Carbohydrate	...	...	...	7	$\times 4 = 28$
Protein	...	...	...	6	$\times 4 = 24$

209.5

Or a diet containing 2,100 calories is arrived at by adding a quarter of each constituent—that is to say, 420 calories. Doubling the constituents brings the total up to 3,360.

But as our patients will not be accustomed to weighing their food in grams, the following tables may be employed as ready reckoners.

WOODYATT'S DIET (1,680 CALORIES).

Diet may be chosen from the following groups:

A. Fats. (Select from these up to 140 grams.)

			Grams of Fat.
5 per cent. vegetables	...	6 ounces	0
10 per cent. vegetables	...	3 "	0
Bacon	...	2 "	30
Butter	...	2 "	50
Cheese	...	2 "	20
Milk	...	2 "	2
Cream	...	2 "	48
Egg	...	One	6
Meat or poultry	...	4 ounces	12

B. Carbohydrates. (Select from these up to 57 grams.)

			Grams of Carbohydrate.
5 per cent. vegetables	...	6 ounces	3½
10 per cent. vegetables	...	3 "	6
White bread	...	4 "	56
Milk	...	2 "	3
Potato	...	2 "	11
Rice	...	2 "	11
Oatmeal porridge	...	2 "	9
Sugar	...	1 ounce	33

C. Proteins. (Select from these up to 50 grams.)

			Grams of Protein.
5 per cent. vegetables	...	6 ounces	3
10 per cent. vegetables	...	3 "	1½
Lean meat	...	4 "	32
Lean ham	...	4 "	32
Poultry	...	4 "	32
Fish	...	3 "	18
Milk	...	2 "	2
Egg	...	One	8

Beverages.

The following may be taken without reckoning: Tea, coffee, oxo, bovril, lemco, water, soda or seltzer water.

Method of Ordering Diet of Given Calorie Content.

Calories.	Grams of		
	A.	B.	C.
840	70	28	25
1260	105	42	38
1680	140	57	50
1965	165	70	50
2100	175	71	63
2295	195	85	50
2525	225	100	50
2520	210	85	75
2930	250	110	60
3360	280	114	100

The majority of patients may not require more than 50 grams of protein; an amount adequate for a weight of 10 st.

Specimen Diets of 1,680 Calories.

The whole carbohydrate is divided between two meals.

Breakfast.	A.	B.	C.
Tea or coffee.			
Milk ½ oz.	1	1½	1
Bread 1½ oz.	1	18	1½
Butter 1 oz.	25	—	—
Bacon 1 oz.	15	—	5
Egg (one)	6	—	8

Dinner.

Bread 1½ oz.	Bread 1½ oz.	Bread 1½ oz.
Roast mutton 2 oz.	Tinned tongue 2 oz.	Sardines 1½ oz.
Cabbage 1½ oz.	Cauliflower 1½ oz.	Cheese 1 oz.
Butter 1 oz.	or	Butter 1 oz.
Bovril.	Lettuce 3 oz.	

Tea.

Tea.	Tea.	Tea.
Milk ½ oz.	Cream 2 oz.	Milk ½ oz.
Butter 1 oz.	Butter 1 oz.	Butter 1 oz.
Bran biscuits.	Egg (one).	Egg (one).
	Bran biscuits.	Bran biscuits.

Supper.

Cheese 1 oz.	Cheese 1 oz.	Boiled ham ½ oz.
Butter ½ oz.	Butter ½ oz.	Butter ½ oz.
Bran biscuits.	Bran biscuits.	Cabbage 1½ oz.
	Oxo.	Bran biscuits.

(At breakfast cream 2 oz. in place of milk.)

Vegetable Groups.

5 per cent. Group.

Cabbage	Marrow	Lettuce (raw)
French beans	Tomatoes	Watercress (raw)
Seakale	Grape fruit	Carrots
Broccoli	Leeks	Parsnip
Spinach	Celery	Beetroot
Brussels sprouts	Asparagus	Lemons
Cauliflower	Turnips	

10 per cent. Group.

Carrots	Green peas	Blackberries
Onions	Melons	Cranberries
Mushrooms	Oranges	Pineapple
Salsify	Strawberries	Peaches

Progress of the Case.

By first establishing the correct dosage of insulin and enabling the patient to take the "optimal diet" it is possible to dispense with frequent blood sugar estimations. Disappearance of glycosuria may imply that the dose of insulin is unnecessarily and even dangerously large. On the other hand, if the patient has a low threshold for sugar glycosuria may be present after the blood sugar has fallen to a safe level.

It seems to be a good working rule to estimate the blood sugar at the end of twenty-four hours of the "basal diet" with insulin administered. The blood should be withdrawn three hours after a carbohydrate meal; if it has not fallen to 100 mg. per 100 c.cm. the dose of insulin is increased; if it has fallen to 80 or lower, the insulin must be diminished.

When the diet is being increased each step in the increase should be continued for three days before estimating the blood sugar. After the patient has reached what is considered the "optimal" diet and the proper dose of insulin ascertained the patient should be instructed how to administer the hypodermic injection of insulin for himself, and also how to test the urine for sugar.

It has been recommended that the patient should be left with a trace of glycosuria. Banting, however, has pointed out that unless the threshold for sugar in the blood has been proved to be constantly below 150 grams it is unsafe to permit even a trace of glycosuria. In the majority of diabetics whose disease is severe enough to warrant the use of insulin glycosuria is likely to indicate a dangerous amount of blood sugar.

The patient must be instructed in the recognition of symptoms of hypoglycaemia, such as weakness, sweating, tremor, nervous apprehension, and disorders of speech. He must also be instructed that if any of these symptoms appear he should at once drink some glucose solution or eat some sugar.

The action of insulin consists not only in enabling the patient to utilize the carbohydrate in the food; it also affords rest to the islet cells of the pancreas. Thus it happens that in some cases there is a recovery of pancreatic function; the pancreas begins to share in the sugar metabolism, so that the quantity of insulin needed is decreased. A number of patients treated with insulin have found that after a time the insulin dose can be reduced or, in a few fortunate instances, even stopped altogether. It is not possible, unfortunately, to promise this in every case; but it demonstrates the importance of making blood sugar estimations at intervals in all patients who are having insulin. Probably intervals of three months will suffice for these estimations, but there is no reason why slight reductions in the dose should not be tried and the effect judged on clinical grounds.

The feeling of well-being that insulin produces is immediate and patients become good judges of what is sufficient for their needs. It may be possible to keep patients in health by giving short courses of insulin with alternating periods of discontinuance. But we have little information on this point.

One drawback to the use of insulin is the expense it entails. The drug itself is costly and the repeated blood sugar estimations add to the bill. At present only the millionaire or the insurance patient can afford insulin treatment. There is no reason why this should continue. As already said, the Ontario Provincial Board of Health issues insulin free to those who cannot afford to pay for it. In England we have enjoyed for many years free diphtheria antitoxin, free calf lymph for vaccination, free novarsenobenzol for the treatment of syphilis, and in addition free blood examinations for patients suspected of having contracted syphilis. Free blood-sugar tests and free insulin for diabetics might be seriously considered by the Ministry of Health.

REFERENCES.

¹ *Physiology and Biochemistry in Modern Medicine*. By J. J. R. Macleod. First edition, 1918, p. 654.

² *Endocrinology and Metabolism*. By Lewellys F. Barker, 1922, vol. 4, p. 263.

INSULIN TREATMENT WITHOUT BLOOD SUGAR ESTIMATIONS.

BY

ARTHUR INNES, M.B., F.R.F.P.S.G.,

GLASGOW.

THE following brief report of a case successfully treated by insulin without any blood sugar estimations seems to me to bring this very efficient substance to the use of the busy practitioner to whom the stipulated numerous estimations are impossible by reason of both lack of time and lack of technique.

A young man, aged 25, a civil servant of more than average mentality, had been under my care for diabetes for more than three years. At first, on a strict régime of fasting and "ladder diet," he did well, but latterly response to this was *nil*, and in March of this year, on a diet of only 400 calories, his glycosuria was severe, acetone was present, thirst very great, skin harsh, dry, and cold, and he always felt drowsy. His weight was 8 st. 4½ lb.

His father could afford the cost of the insulin, but not the expense of a nursing home or specialist supervision. I failed to obtain him admission to those wards where this treatment was then being investigated, and resolved to undertake the treatment in his lodgings and without blood sugar estimations, for his condition was daily becoming more grave.

As already stated, the urine contained acetone, and not knowing at this time the danger of giving insulin in such cases unless glucose be injected at the same time, I gave 5 minims, equivalent to 7 units of insulin. A most alarming collapse occurred, which, however, was promptly overcome by giving a tablespoonful of sugar and water solution, which we had prepared beforehand in case any untoward symptoms should develop. He received this very small dose at 1 p.m. and had a meal of soup and beefsteak immediately thereafter. At 3.30 he wrote in a diary which I asked him to keep, "Entire cessation of thirst. Inclined to perspire. Palms of hands moist. Feeling of great warmth all over." This latter feature impressed him tremendously. From that date to now he has never had acetone in his urine, and he has done hundreds of examinations of it for me.

The doses were increased daily until on the fifth day he was receiving 20 units all in one dose. No ill effects followed this dose, but it did not cause the disappearance of sugar from the urine excepting from the specimen passed three hours after the injection. On the seventh day we gave him 20 units intravenously without any bad effects and with complete disappearance of glycosuria for ten hours thereafter; but it recurred as bad as ever, so that though the immediate effect was excellent it had no better sustained action than the subcutaneous.

On the next day we reduced the starches in his food a little more and increased the fat, as we had now no fear of acetone, and on the next day after his injection we observed palpitation, nausea, and profuse sweating, but these symptoms disappeared at once on sipping a little sugar and water. We learned from this to hasten slowly in the sugar reduction, and kept him on a diet of approximately 57 carbohydrate, 114 protein, and 107 fat—giving about 1,620 calories. He was feeling well and his weight was now 9 st.

On the twenty-sixth day from the start of the treatment no insulin was given, and as a result copious glycosuria and thirst returned at once, and, strangely enough, daily doses of 22 minims of insulin with a diet of only 25 carbohydrate, 58 protein, and 98 fat failed to keep the patient well, and I proposed, and he agreed, to observe two days' fasting followed by a week's "ladder diet," and after this to resume the insulin injections. The results of this method were as follows:

June 13th. Fasting. Sugar all day.

June 14th. Fasting. Sugar cleared in the evening.

June 15th. Carbohydrate 10, protein 4, fat 3—calories 83. No sugar.

June 16th. Carbohydrate 16, protein 8, fat 6—calories 150. No sugar.

June 17th. Carbohydrate 14, protein 16, fat 12—calories 228. No sugar.

June 18th. Carbohydrate 14, protein 28, fat 18—calories 330. No sugar.