Finally, I have tried to bring forward arguments to show that clubbing is the result of defective oxidation in the tissues of the extremities, whether produced mechanically by obstruction to venous return, or as the result of a general lowering of the oxygen tension of the blood affecting parts of the body where normally the circulation is slow. This seems to offer the best explanation of the recorded facts concerning this very interesting phenomenon, and incidentally suggests that Pigeaux's¹⁴ conception of the development of the deformity was correct.

TYPES OF DEXTROCARDIA.

(With Special Plate.)

BY

H. WALLACE JONES, M.Sc., M.D. LIVERP.,

M.R.C.P. LOND.,

HONORARY ASSISTANT PHYSICIAN, LIVERPOOL ROYAL INFIRMARY.

(From the Cardiographic Department of the Royal Infirmary.)

DEXTROCARDIA in its widest sense is the term applied to the condition in which the heart beats in a more or less permanent manner in the right half of the thorax.

This abnormality has been known from very early times, and the first recorded case was that reported by Severinus of Rome in 1643,¹ while Riolano,² also in the seventeenth century, reported two cases, one of which was Queen Marie de Médicis. Even as early as 1749 dextrocardia had been divided by Senac³ into congenital and acquired groups. Most of these early cases were based on *post-mortem* findings, and it was not until the beginning of the nineteenth century, when auscultation and percussion became perfected, that many cases were discovered during life. In more recent times, owing to the application of x rays and the electro-cardiograph, the diagnosis has become comparatively easy.

Cases of dextrocardia may roughly be divided into con-

FIG. 2.—Case I. Transposition of heart and other organs. Lead I inverted, transposition of Leads II and III.

FIG. 4.—Case II, a case of complete transposition. The usual record of a transposition case—that is, Lead I inverted.

genital and acquired, but it is more usual to subdivide them in the following way:

I. Inversions.—“Mirror image” of the heart.

- (a) Complete.
- (b) Incomplete.

1. Transposition of viscera with heart in the normal position.
2. Complete inversion of the thoracic organs and a liver situated in the middle line.
3. Inversion of the heart only.

REFERENCES.

- ¹ Buhl: *Lungen-Entzündung, Tuberkulose und Schwindsucht*, 1872, 159.
- ² Freytag: vide Symes-Thompson, *Med. Chir. Trans.*, London, 1904, lxxxvii, p. 85.
- ³ Liebermeister: vide Shaw, *Allbutt and Rolleston's System of Medicine*, 1907, iii, p. 67.
- ⁴ Buzzard, F.: *BRITISH MEDICAL JOURNAL*, 1901, i, 1333.
- ⁵ West: *Clin. Soc. Trans.*, xxx, 60.
- ⁶ Gay: *Path. Soc. Trans.*, xxii, 111, 1871.
- ⁷ Smith: *Path. Soc. Trans.*, xxiii, 78, 1872.
- ⁸ Stadie: *Journ. Experim. Med.*, 30, 215, 1919.
- ⁹ Haldane: *Proc. Roy. Soc. Med.*, 13, Therap. Sect., 66, 1920.
- ¹⁰ Stokvis: *Internat. Beitr. z. inner Med.* (Leyden's Festschr.), Berlin, 1902, i, 597.
- ¹¹ Van der Bergh: *Nederl. Tijdschr. voor Geneesk.*, 1901, i, 761.
- ¹² Schwartz: *Arch. Int. Med.*, 1916, xvii, 459.
- ¹³ Béclère: *Bull. et Mém. Soc. Méd. des Hôp. Paris*, 1901, xviii, 283.
- ¹⁴ Pigeaux: *Arch. gén. de Méd.*, 1832, xxix, 174.

II. *Dextroversions*.—Heart displaced to the right without inversion of the cavities.

- (a) Occurred during foetal life.
- (b) Acquired since birth.

INVERSIONS.

The complete variety is comparatively common; over two hundred cases are recorded in the literature.⁴

The incomplete type is, on the other hand, extremely rare.

Type 1 is seen in the case reported by Royer and Wilson⁵—a boy, 6 years old, who was found to have complete transposition of the viscera with the heart in the normal position.

The lungs were included in the transposition, the three lobes being on the left side. The heart had a patent foramen ovale; pulmonary stenosis, and a rudimentary left ventricle.

Type 2 is represented by the celebrated case of Abernethy,⁶ in which the heart of a 10 months old infant was on the right side with the liver in the middle line, the rest of the organs being in the normal position.

Of Type 3 only two cases are recorded—that reported by Moffet and Neuhof⁷ which was associated with patent interventricular septum, and that reported by Meyer⁸ which was not associated with any other developmental error. This variety is very much rarer than one would suppose from the frequent way in which it is described.

All these cases of inversion of the heart have what may

FIG. 5.—Case II. The normal record obtained when the right arm and left arm leads are reversed.

be termed a “mirror image” of the normal heart—that is, with inversion of the cavities.

The electro-cardiographic record of this variety shows complete inversion of lead I with minor changes in leads II and III, in distinction to the record obtained in the second main type, or dextroversions, where the heart is merely displaced and which give a normal cardiographic record.

It is by means of this characteristic record of the transposition cases that they are most readily distinguished from

the cases of displaced heart or dextroversion. The electrocardiogram is therefore of great help in those cases in which the transposition is confined to the heart, and in which the x-ray appearances are very similar.

DEXTROVERSIONS.

Type (a) may be due to inflammatory conditions during foetal life or to defects in developments, such as absence of a lobe of the lung, diaphragmatic hernia, or to atelectasis.

Type (b) may be due to changes in the left side of the chest pushing over the heart—such as pneumothorax, pyopneumothorax, haemothorax, and occasionally mediastinal tumours—or to changes in the right side, such as pleurisy, fibrosis of the lungs and pleura, or bronchiectasis drawing over the heart.

In the congenital type there are often associated abnormalities of development, defects of the septum being common—a case with defect in the auricular septum was recorded by Foggie⁹ and one with a defect in the ventricular septum by Moffet and Neuhof.⁷ Ordinary valvular disease is occasionally associated with the condition, as is seen in the case of complete transposition with mitral stenosis described by Owen¹⁰ and by the fourth case of our series.

Clerc and Bobrie⁴ have drawn attention to the interesting fact that a heart beating to the right is often more irritable than one beating in the normal position, and patients often complain of palpitation and many of the symptoms of the "effort syndrome" without any definite evidence of cardiac failure. This is very well shown in the third case of this series. Occasionally dextrocardia tends to run in families, and a case of brother and sister with this condition is related by Neuhof.¹¹

In the following four cases of dextrocardia two of them belong to the transposition group and two to the acquired variety.

CASE I.

A boy, aged 10, with complete transposition. He has no symptoms; the condition has been present since birth. Heart dullness to the right of the sternum, liver dullness to the left, splenic dullness to the right. The stomach resonance is on the right.

X rays show transposition of heart, liver, and stomach (Fig. 1). The electrocardiogram shows typical inversion of deflections in lead I, with transposition of leads II and III (Fig. 2).

CASE II.

A woman, aged 40—a case of complete transposition. I am indebted to Dr. Murphy and Mr. R. E. Kelly for being able to include this most interesting case in this series. She has been operated on by Mr. Kelly for gall stones and an hour-glass stomach—of course, on opposite sides to the normal patient. He has drawn attention to the interesting fact that if dextrocardia occurs about 1 in 4,000, gall stones 1 in 10, and an hour-glass stomach 1 in 1,000, the chances of any one individual possessing all three is about 1 in 40,000,000, and this patient therefore ought to be unique in the British Isles.¹²

Her heart is normal in size, with an apex beat $3\frac{1}{2}$ in. to the right in the fifth space, and her liver dullness is on the left side.

X rays show complete transposition (Fig. 3).

The electrocardiogram shows inversion of lead I with the usual transposition of leads II and III (Fig. 4). Fig. 5 shows a record from the same patient after the leads from the right and left arms have been reversed. The record shows a normal deflexion in all leads.

CASE III.

A woman, aged 35, who has known for the last twelve years that her heart has been beating on the right side. She can remember a time, fifteen years ago, when her heart was examined and said to be normal. She had influenza and pleurisy two years ago, and most of her symptoms date from this time, when attention was again drawn to the fact that her heart was not in the normal position. She now suffers from pain over the heart which radiates to the right shoulder and neck, with shortness of breath on exertion. There is occasional sweating at night and

FIG. 6.—Case III. Acquired dextrocardia. (Electro-cardiogram normal.)

she has had a winter cough for the last year or two. Her apex beat is in the fifth space, $4\frac{1}{2}$ in. from the middle line on the right side; the heart sounds are normal. There is slight dullness over the lower lobe of the left lung with slightly deficient breath sounds and diminished vocal fremitus and vocal resonance. The liver dullness is in the normal position.

X rays show the heart on the right side, no evidence of pulmonary tuberculosis or any other lung disease. The electrocardiogram shows normal deflexion (Fig. 6).

CASE IV.

A man, aged 27, who was suddenly seized with shortness of breath in 1917 while on active service in the East, and when examined by his medical officer was found to have his heart beating on the right side. He now suffers from shortness of breath on exertion with pain across the chest and has occasional sweating at night. There is slight bulging of the precordium; the apex beat is in the fifth space $3\frac{1}{2}$ in. to the right of the middle line. There is a well marked presystolic thrill and bruit with a low rumbling diastolic bruit at the apex. No abnormal physical signs were detected in the chest.

X rays show the heart displaced to the right with no signs of any change in the lungs to account for it. The electrocardiogram shows a normal deflexion (Fig. 7).

FIG. 7.—Case IV, a case of acquired dextrocardia.

In conclusion, I have to thank Dr. John Hay for his very kind help in preparing this series; and also Dr. R. E. Roberts for taking the x-ray photographs.

REFERENCES.

- 1 Upson: *Amer. Journ. of Roentgenology*, 1921, vol. 8, p. 385.
- 2 Riolo, Giovanni: Cité par Otto. *Nouvelles observations rares d'anatomie, de physiologie et de pathologie*. Berlin, 1829.
- 3 Senac: *Traité de la structure du coeur, de son action et de ses maladies*, 1749.
- 4 Clerc et Bobrie: *Archives des Maladies du Coeur*, 1918.
- 5 Royer and Wilson: *British Journ. of Children's Diseases*, vol. 5, p. 176.
- 6 Abernethy: *Phil. Trans.*, 1793.
- 7 Moffet and Neuhof: *Journ. of Amer. Children's Diseases*, July, 1915.
- 8 Meyer: *Arch. des Maladies du Coeur*, sixteenth year, No. 4, p. 254.
- 9 Foggie: *Edin. Med. Journ.*, vol. 5, No. 5, p. 428.
- 10 Owen, Sydney: *Heart*, vol. 3, No. 1, p. 113.
- 11 Neuhof, Selian: *Journ. Amer. Med. Assoc.*, April, 1913, p. 1064.
- 12 Kelly, Robert E.: *British Journ. of Surgery*, vol. 8, No. 29, 1920.

SIR ROBERT WOODS : RETROPHARYNGEAL LIPOMA.

FIG. 1.—C, Cricoid. H, Hyoid.

FIG. 2.—C, Cricoid. H, Hyoid

H. WALLACE JONES : DEXTROCARDIA.

FIG. 1.

FIG. 3.