syphilis of the central nervous system so far as we have been able to ascertain.

5. Definite changes (other than a complete positive, and in the absence of blood) indicate apparently some definite pathological change in the cerebro-spinal fluid; but all pathological fluids do not necessarily show any colloidal benzoïn reaction.

AN EFFICIENT INTESTINAL ANTISEPTIC:

THE PERCHLORIDES OF MERCURY AND IRON
IN COMBINATION.

BY

T. STACEY WILSON, M.D. EDIN., F.R.C.P.,
CONSULTING PHYSICIAN, BIRMINGHAM GENERAL HOSPITAL.

SOME recent correspondence in the *BRITISH MEDICAL JOURNAL* has prompted me to bring forward the following evidence that a combination of the perchlorides of mercury and iron constitutes an efficient intestinal antiseptic.

My attention was first drawn to the value of this remedy by a short notice in this *JOURNAL* for May, 1894, by Dr. W. B. Wedgwood of King's Lynn, who wrote recommending its use in enteric fever, and suggesting a dose of 30 minims of liquor hydrargyri perchloridi and 20 minims of tinctura ferri perchloridi. Ever since that time I have used these two drugs as an intestinal antiseptic, combined with a little glycerin and chloroform water, but have found that 15 minims of each is quite sufficient to give a good result in those cases in which it is indicated, and I have every reason to be extremely thankful to Dr. Wedgwood for having drawn my attention to this remedy.

In Enteric Fever.

I first tried this prescription in a comparative test in enteric fever, for I took a hundred consecutive cases, of whom one-third had been treated with mercury and iron, one-third with sulphurous acid, and the remainder in various other ways. They were divided into groups according to the number of days that the pyrexia lasted after their admission to the hospital. The average height of the temperature was then ascertained for each group by dividing the week into two portions and taking the average morning and evening temperature for each case during each half-weekly period. The general run of the temperatures as thus averaged proved to be wonderfully uniform in all the groups, and so far as the morning temperature was concerned the average rate of fall of the temperature was not appreciably influenced by the treatment adopted. In the case of the evening temperature, however, where mercury and iron had been given, the average was more than a degree lower than where other treatment had been adopted.

The reliability of the test in this series of cases was enhanced by the fact that they were all in charge of the same two ward sisters, and the excellence of the nursing was shown by the fact that there were only three deaths out of the hundred consecutive cases.

Owing to the fact that few of the patients came into the hospital during the first week it was not possible to decide as to the effect of the mercury and iron treatment upon the total duration of the pyrexia, but where this treatment was adopted the case appeared to run a shorter and more favourable course. One of the fatal cases gave evidence as to the potency of the antiseptic action of this combination. It was complicated by acute nephritis, and during the first ten days of his illness the patient had given himself a liberal allowance of roast beef and mutton. He died somewhat suddenly a few hours after an injection of morphine given by a resident on his own initiative, and at the *post-mortem* examination over two hundred ulcers were found in the intestine, many of them quite deep, but all free from sloughs and all looking remarkably healthy and showing signs of healing. The unusually satisfactory condition of the ulcers considering their size and depth seemed to be distinctly traceable to the mixture which he had been taking every four hours for nearly two weeks.

In Colitis and Dysentery.

Evidence of the antiseptic value of the mercury and iron mixture is given also by its efficiency in acute and subacute colitis, especially when of the ulcerative type, and it often causes very rapid improvement in cases which have resisted other methods of treatment. The mixture proves of great service also in the treatment of dysentery. While I was acting as visiting physician to the First Birmingham War Hospital at Rubery, a convoy of over a hundred cases of dysentery was received and put under the charge of myself and the late Major Orr (of Ludlow). In consequence of my experience with ulcerative colitis I decided to give all my cases with three exceptions the mercury and iron mixture, and ordinary house diet consisting of beef and mutton and potatoes, etc., although not a few of them were having twelve to eighteen stools a day, and some still had a little pyrexia. Major Orr decided to put his cases on the same treatment.

The result of these somewhat heroic measures was, however, perfectly satisfactory, for there was not a man who did not make rapid progress to recovery. The three exceptions mentioned above were cases with high fever and extremely frequent evacuations; they also were given the mercury and iron mixture, but fed on finely divided raw meat and carbohydrate in the form of fresh juice expressed from pounded carrots after steeping for two hours in water at a temperature of 120° F. Carrot juice thus prepared contains albuminous material and vitamins free from vegetable fibre, and also inositol—a non-fermentable sugar which is of much value in such cases. The patients on house diet got on so well that no bacteriological investigations were made owing to the restricted laboratory service available, but the three severe cases were found to be due to Shiga's and to Flexner's bacilli. The *Amoeba coli* was not found.

Although I have had no experience of the treatment of acute dysentery I have little doubt that this remedy will prove of service, and this conviction is strengthened by the knowledge that a nurse whom I had treated with mercury and iron for colon trouble and who went out to a mission hospital in North Borneo introduced this mode of treatment of acute dysenteric conditions there with most satisfactory results. It is a remedy which proves of great value in cases where muscular disturbance of the colon is giving rise to tenderness and hardness of the bowel together with pain or other nervous symptoms. It is of special value where this type of colon trouble is causing mental depression, phobias, or obsessions, in place of pain, and in such cases its curative action is sometimes extraordinarily rapid.

In Addison's Anaemia.

The latest theories respecting the causation of "Addison's anaemia," as published by Dr. Arthur Hurst in this *JOURNAL* for January 19th, make it highly probable that this combination of mercury and iron may prove of value in this hitherto fatal disease. Naturally there has not yet been time to put this suggestion to a clinical test; I can, however, refer to a patient who recovered from a very severe second relapse of "Addison's anaemia," during which he was taking the mercury and iron for colon disturbance. I thought the cure of the relapse was due to the vaccine and dietetic treatment he was given, and it was a great disappointment to find that a year or two later this same treatment proved quite ineffective in averting the fatal termination of a third relapse of this complaint. On this occasion colon disturbance was not a prominent symptom, and if I remember right the mercury and iron mixture was not persevered with throughout the attack. Considering the powerful antiseptic action of mercury given in this way, it is possible that it may sometimes prove curative in this hitherto incurable disease, if it be really due to the action of streptococci in the duodenum or small intestine.

Precautions.

When using mercuric perchloride as an antiseptic certain things must be borne in mind.

1. Combination with albumin or alkalis will destroy its antiseptic value. It must therefore be given when the

stomach is empty, and must also be combined with some drug which has more powerful affinities for alkalis (such as ferric perchloride) in order to prevent its neutralization by the alkalis of the duodenum. Where there is retention of the gastric contents owing to atonic dyspepsia or pyloric spasm its use is contraindicated owing to the certainty of the mercurial salt being converted into an albuminate.

2. It must be remembered that the two perchlorides in some cases tend to irritate the gastric mucous membrane, and therefore the remedy is best given in circumstances which ensure as rapid a passage into the duodenum as possible. This object is best attained by giving the drugs fifteen or twenty minutes before meal-times when the stomach is already becoming active in preparation for the reception of a meal.

In acute cases, when definite meals are not given, the remedy can be given every four hours, and is usually quite well tolerated. This combination of perchlorides is as a rule wonderfully well borne, even by patients who are quite unable to take iron as usually administered after meals. I have not infrequently known patients take this mercury and iron mixture continuously for six months or more, except that sometimes, but not always, it is in women found advisable to omit it for a day or two at the commencement of the menstrual period.

Another fact to be borne in mind is that sometimes the continued use of this drug causes a loss of weight, suggesting the possibility that the antiseptic interferes with some of the bacteriological processes upon which digestion depends; it is therefore well to watch the weight in patients who are taking it regularly.

PERNICIOUS ANAEMIA:

SOME MORPHOLOGICAL AND ETIOLOGICAL CONSIDERATIONS.

BY

A. PINEY, M.B., CH.B. BIRM., M.R.C.P.,

DIRECTOR OF THE INSTITUTE OF PATHOLOGY, CHARING CROSS HOSPITAL.

THE paper by Dr. A. F. Hurst in this JOURNAL (January 19th, p. 93) has once more called attention to the haemolytic conception of pernicious anaemia. He states that—

"The most characteristic features of the blood in Addison's anaemia are now recognized as being the great inequality in the size of the red corpuscles—the anisocytosis—due to the presence of microcytes and megalocytes, and the increase in their average size in spite of the presence of some cells far smaller than the smallest cells found in normal blood."

These features are not specific unless they are associated with a large amount of haemoglobin. Anisocytosis without such excess of haemoglobin can be observed in all the secondary anaemias. Dr. Hurst has made use of the graphic method of determination of the size of the red corpuscles, and, valuable as this is, it is less accurate than the viscosimetric method of estimation of the average volume of the red corpuscles. This defect of the graphic method is well seen in acholuric family jaundice, where the average size of red corpuscles is high although the cells look small; this is due to their globular form. The viscosimetric method demonstrates that the high average volume persists throughout the remissions of pernicious anaemia.

Dr. Hurst states that megaloblasts may be found in the blood in conditions other than pernicious anaemia—for example, lead poisoning. The difficulty which arises here depends upon faulty nomenclature. The various types of nucleated red cell seen on occasions in the course of examining stained films are, first, normoblasts, which are fully haemoglobinized red cells of normal size (incompletely haemoglobiniferous cells are also seen); secondly, macroblasts, which are large pale specimens of the former; and, thirdly, megaloblasts, which are supplied with a full complement of haemoglobin, and, when young, have a finely reticular nucleus. The megaloblast does not in any way resemble a very immature form of nucleated red cell, as does the pale macroblast. In the course of its development the

megaloblast loses its nucleus, and the resulting cell is a megalocyte, which is supplied with a full complement of haemoglobin. Normally a macroblast is converted into a normoblast, which loses its nucleus and gives rise to the ordinary red corpuscle, but under some circumstances the macroblast loses its nucleus and gives rise to a pale (polychromatophilic) red cell of large size (macrocyte).

The megalocyte is thus seen to be a mature corpuscle, while the macrocyte is incompletely developed. The difference between these cells lies in the fact that, while both are giants, the macrocyte is only morphologically gigantic, whereas the megalocyte is also a functional giant. Osler called attention to a peculiar feature of pernicious anaemia:

"A surprising fact is that there are many patients with extreme anaemia who are remarkably vigorous. One may see patients with a count of about two million red cells who insist that they are able to do everything as usual, except for a little shortness of breath."

No other anaemia presents this astounding feature, and in no other anaemia of adult life are megaloblasts and megalocytes, which are the functional giants accounting for this peculiarity, formed. Hurst agrees with those writers who regard the presence of megaloblasts and megalocytes as being due to "an unusual degree of regenerative activity," although a little later he attributes the leucopenia of the disease to defective function of a poisoned bone marrow. The reconciliation of the two statements appears to be impossible.

Hunter's conception of pernicious anaemia as a haemolytic anaemia receives Hurst's support, but there is no resemblance between it and the known haemolytic anaemias—namely, acholuric icterus. There is definite morphological evidence of defective function of the marrow in that the hyaline leucocytes are always reduced in number and that there are always a varying number of polymorphonuclear leucocytes of large size with hypersegmented nuclei. All the known toxic conditions cause fine changes in the polymorphonuclear leucocytes—for example, diminution in number of the granules and irregular size and distribution; these changes are absent in pernicious anaemia, and confirm the opinion that the toxin of the disease acts on the bone marrow, not on the circulating corpuscles.

Achlorhydria is regarded by Hurst as an invariable concomitant and precursor of the disease, but one of the cases recorded by Levine and Ladd¹ showed some free acid in the gastric juice when first observed, and the diagnosis was confirmed at autopsy. I have observed a case of subacute combined degeneration of the cord, in which free acid was present; there was no anaemia, but viscosimetry showed that the average volume of the red corpuscles was already high. The diagnosis was confirmed by the later history of the case and by autopsy. Achlorhydria, when once developed, is persistent, but not more so than the increased average volume of the red cells, the reduction of hyaline cells, and the hypersegmentation of polymorph nuclei.

Cases of pernicious anaemia occurring after gastrectomy for carcinoma are attributed by Hurst to the resultant achlorhydria. These cases may be of great value, but no detailed blood counts are published, and there must therefore be some doubt whether these cases were pernicious anaemia or the "pseudo-pernicious" anaemia associated with secondary deposits in bones, which may closely simulate the disease.

There is obviously excessive haemolysis in the course of the disease, but this, apparently, depends upon the peculiar type of cell which is presented to the haemolytic organs; the haemolysis is the cause of the anaemia but not necessarily of the disease.

Zadek,² who trephined the tibia during a remission of the disease, found that there had been reversion to fatty marrow and very few cells were present. Myeloblasts are an invariable constituent of the bone marrow in pernicious anaemia, but are not present in the marrow of the normal adult or in other anaemias—that is to say, they are embryonic cells. The megaloblast as defined above is also an embryonic cell, and thus the two types of cell together conform to the old statement of Ehrlich that there is a reversion to an embryonic type of blood formation. Zadek has only