complication, with resulting contamination of the milk by living bacilli.

The following is a very brief account of my seven cases:

CASE I.

A girl, aged 17, well nourished and robust, was admitted to hospital with extensive tuberculous glands on both sides of the neck. She had been operated on several times for removal and abscess formation, but the infection proceeded downwards to the axillary glands, and on the surface of the chest wall, towards the breast, which became hard but not painful. In a few weeks an abscess formed which was opened and scraped. A persistent sinus resulted, which only healed after a long course of human tuberculin. The girl made a good recovery, all the glands subsided, the sinuses and ulcers healed, and she was discharged from the hospital after ten months' treatment, apparently cured.

On examination of pus no tubercle bacilli could be demonstrated under the microscope, but scraping from the breast tissue showed them in large numbers. An emulsion inoculated into a rabbit produced an acute general tuberculosis, and growths from the rabbit's spleen showed typical bovine cultures.

CASE II.

A woman, aged 37, seen in consultation. She had suffered from tuberculous neck glands for over twenty years, healing up and breaking down at intervals. A year ago she noticed the right axillary glands to be enlarged, the infection spreading to the right breast. The lungs were apparently healthy. Two large sinuses were seen in the lower quadrant of the breast, discharging greenish-yellow pus.

Examination of the pus showed tubercle bacilli, and animal inoculation (rabbit) produced a general tuberculosis with typical bovine cultures. I advised amputation of the breast, but she declined, and I have not been able to learn what happened to her.

CASE III.

A girl, aged 19, was admitted to hospital with very extensive tuberculosis of the right neck glands. She had been brought up in Canada and had drunk a very large quantity of cow's milk during the last twelve years. The infection had spread by direct lymphatic extension to the axillary glands, which had suppurated, and thence to the breast, which was in a condition of general tuberculous mastitis with five sinuses discharging a large amount of pus daily.

The sinuses were scraped and packed with iodoform emulsion, and she was put on a course of human tuberculin. After a prolonged suppuration—ten months—they gradually closed, and she was sent to a sanatorium almost well. After a residence of six months, during which time she increased over 2 st. in weight, she was discharged with all sinuses and ulcers soundly healed, and she has remained quite free of the disease to the present time.

CASE IV.

A woman, aged 33, was admitted to hospital with extensive mastitis of the breast, numerous discharging sinuses, and a very fetid discharge.

History.—At the age of 11 she first developed tuberculosis of the neck glands, which had been treated by incision and scraping on several occasions. There were several sinuses in the posterior triangle on the right side; the left side of the neck was quite clear. For over twenty years the glands had been more or less active and she had lost weight during the last six years. The infection had spread down the superficial lymphatics of the right side of the chest, over the clavicle and direct to the breast. Numerous ulcers and scars were present over the chest wall.

Treatment.—The sinuses were opened up and scraped and the cavities cleansed with a strong antiseptic, and afterwards packed with thymol. She was given a prolonged course of human tuberculin, and very soon began to show improvement. The progress was slow but progressive, and after a residence in hospital of fourteen months she was discharged, soundly healed, and has remained quite well up to the present time.

CASE V.

A male, aged 27, was admitted to hospital suffering from tuberculosis of the shoulder (left). The head of the humerus had been excised seven years before, but the wound had never healed. He complained of pain, especially on pressure over the left breast. There was a uniform swelling of the breast tissue 3 inches in diameter, with the nipple as the centre. There was fluctuation, and a needle showed thick pus; over an ounce was withdrawn and found to be quite sterile. No tubercle bacilli could be seen under the microscope.

Animal Inoculation.—A guinea-pig and a rabbit were inoculated with 1/2 c.cm. each of the pus, under the skin of the abdomen. Both animals developed progressive tuberculosis and died in four months.

Post mortem the rabbit showed a generalized tuberculosis of all the organs, with enormous caseation of the mesenteric glands. The guinea-pig showed a generalized infection, with deposits in liver and spleen, but no affection of lungs. Cultures made from the spleen of the rabbit on glycerin agar showed after six weeks a typical growth of bovine tubercle.

Progress of Case.—The abscess was evacuated with a trocar on three subsequent occasions, and disappeared. The patient left the hospital in two months, and I have not been able to trace him since.

CASE VI.

A man, aged 23, has had tuberculosis of neck glands for seven years, and involvement of axillary glands for two years. The glands had been excised extensively, but repeated recurrences had

involved further operations and scrapings. On admission he was thin and wasted, with all the appearances of lardaceous disease. His neck was one mass of ulcers and sinuses. The left axillary glands had suppurated and had been drained, leaving discharging sinuses. The left breast was enlarged, tender, but not suppurating.

Treatment.—The sinuses were treated antiseptically, but it was impossible to attempt anything of an operative character. The mastitis, however, gradually subsided, the pain on pressure disappeared, and the breast became normal in size. He gained weight, and left the hospital, but I have been unable to trace him since his discharge.

CASE VII.

A male, aged 22, seen in consultation, complaining of pain in the abdomen, right knee, and right breast. He had suffered from tuberculosis of the knee-joint, tibia, and ankle, with abscess formation two years previously. The abscesses had all been opened and drained, and when seen all the sinuses were discharging pus. Two months before his right breast became large and very tender. On examination there was general mastitis, the swelling was uniform, there were no nodules, and no fluctuation. The chest was apparently normal.

Treatment.—The breast was treated twice daily with iodoform inunctions, and a general outdoor life with rest and tonic prescribed. In addition he was given a course of twelve injections of human tubercle vaccine in increasing doses. He was seen again three months later, when a great improvement was apparent. The breast was normal in size, there was no tenderness, the sinuses (with the exception of one over the ankle) had quite healed, and he had gained 22 lb. in weight. As there was no suppuration of the breast, it was not possible to prove its tuberculous origin, but I have no doubt it was a case of early tuberculous mastitis, which was arrested. He is continuing to improve, and his doctor informs me that he is leading an ordinary life, and has returned to duty.

General Conclusions.

From a long clinical observation of tuberculosis in all its forms, embracing over 10,000 cases in hospital and private practice, I have formed the following conclusions:

(1) Tuberculosis of the breast is always caused by the bovine bacillus.

(2) It is never primary, but is the result of lymphatic extension from the glands of the neck or axilla (excepting in the rare cases of general miliary infection).

(3) In common with all other forms of surgical tuberculosis caused by the bovine bacillus, it responds readily to treatment by a tubercle vaccine prepared from an attenuated culture of human tubercle bacilli. The treatment must, however, be continued for a long period until an active immunity is established.

HAEMATEMESSES "WITHOUT LESIONS."*

BY

GUSTAVE MONOD, M.D. PARIS, M.R.C.P. LOND.,

PHYSICIAN TO VICHY THERMAL HOSPITAL.

I REPRESENT here the association of the Paris Faculty of Medicine for the furtherance of medical relations. It is our great desire to keep in touch with our British confreres. We will never forget the years we lived shoulder to shoulder.

International medicine has its organizations—the congresses—but times are hard, and we have been wondering if we might not find a simple bond of union by discussing simultaneously the same subject in the medical societies of the different countries. It is for this reason that I am bringing before you a question which is being discussed at the present time by the Gastro-enterological Society of Paris, and is shortly to be brought before the Cercle Médical of Antwerp. The best proof I can give of the opportunity of a working liaison between our countries is the fact that when I went through the literature of the subject I found that it had been very thoroughly investigated and discussed by our British colleagues.

I am seeking your criticisms and advice in answering the two questions: Is there such a thing as haematemeses without lesions? and, if so, how are such cases to be explained, and how should they be dealt with? In 1815 Latour published in Orleans a book—*Histoire philosophique et médicale des hémorragies*—in which he recognized gastric bleeding *sine materia*, and quoted cases from Portal, Franck, Hoffmann, van Swieten, Baldinger, and Warton. The question had been raised at an even earlier date (1803) by Dalché. In

* A paper read to the Medical Section of the Royal Society of Medicine.

our day, when presumptions give way to facts, Sir William Hale-White, in a lecture published in 1906, relied only on facts proved on cases where diagnosis was established either by operation or by autopsy. He collected twenty-nine cases published in this country. He pointed out that such cases were very rare in males, whereas deaths from haemorrhage due to ulcers are far more frequent in males than in females; he concluded either that "the outlook of gastric ulcer in women is much more favourable than it is in men, or many cases occurring in women and classified as gastric ulcer at the bedside are really instances of some other undescribed malady." To such a malady Sir William Hale-White gave the name "gastrostaxis." To his twenty-nine cases let us add a few recent ones observed in Paris hospitals. I will quote three reported to the Society of Gastro-enterology by Professor Pierre Duval and Dr. J. Ch. Roux, but not yet published. All three patients were women.

CASE I.

A woman, aged 24, complained of painful dyspepsia of nine years' duration, and for a year of haematemesis of 2 to 3 ounces.

First Operation.—She was operated on at the age of 16 and the appendix was removed (as is often the case with this type of patient). A blood count showed 4,900,000 red cells. Bleeding time, three minutes; coagulation time, twenty minutes. The screen showed an elongated stomach (mark this fact) and stagnation in the second portion of the duodenum.

Second Operation.—As medical treatment gave little result an operation was performed, and slight adhesions were found around the gall bladder.

Third Operation.—Three weeks later there was a new crisis—severe haemorrhage per anum. Red blood corpuscles 2,500,000. The surgeon, thinking that he must have overlooked a duodenal ulcer, operated again. The duodenum was opened from end to end, but no lesion was seen. Gastro-enterostomy was performed and the patient again made a good recovery.

Fourth Operation.—Three months later there was fresh haemorrhage (slight). Four months after this the patient was admitted again for severe pain. She was operated on for the fourth time. The surgeon opened the jejunum; no ulcer was found, but an unabsorbed stitch was removed. Since then the patient seems to be cured.

CASE II.

The patient complained of late pains after food. She had an acute attack of haematemesis. Professor Duval opened the stomach from cardia to pylorus. The duodenum was carefully inspected, but no lesion whatever was found.

CASE III.

A patient, aged 32, was operated on and the appendix removed seven years ago. She was admitted for haematemesis which appeared at the time of menstruation. An x-ray plate showed stenosis of the third portion of the duodenum. At the second operation a mesocolic band was found pressing on the duodenum, but no ulcer. Duodeno-jejunostomy was performed and she recovered.

Here, then, are three recent operations where ulcer was diagnosed but was not present. In two cases there was oozing or perhaps microscopical erosions. In the third case the bleeding was from congestion due to duodenal stenosis.

M. Basset, a well known Paris surgeon, has recorded three very similar instances on which he operated, and Professor Duval also has recently had two cases in which he thought to find ulcers. He operated and found nothing; both his patients recovered once for all.

M. Robineau, surgeon to the Necker Hospital in Paris, records the story of a patient, aged 39, who had haemorrhage so profuse that the blood count numbered 950,000, with 25 per cent. haemoglobin. The patient gave no actual history of syphilis, but stated that she had had hemiplegia which responded to intravenous injections. After a long and difficult examination under x rays a diverticulum of the fourth portion of the duodenum was found. M. Robineau dissected out the small pouch; the operation was prolonged owing to excessive bleeding from the gut. Neither macroscopically nor microscopically was any lesion to be found; a submucous angioma was not present. The patient recovered normally and has not since been troubled by haemorrhage.

Our cases support the demonstration that Sir William Hale-White and Lord Dawson have given, and the question of the causes of such haemorrhages arises. I think we may eliminate chlorosis—an ailment which, for obscure reasons, is losing ground every day. We hesitate, too, to call them vicarious. They do not follow the rhythm of normal periods, and we do not find that our cases report suppressed haemorrhoidal bleeding. Hind relates the case of a woman who suffered from haematemesis once a week for nine consecutive years.

I have to admit that the cases collected in France lack precision, the reason being that all were urgent, and that we do not admit a diagnosis of gastrostaxis unless direct proof be given. In cases not operated on the diagnosis of the ulcer is maintained.

Shall we find a possible cause in some nervous disorder? The well known French gastrologist Mathieu used to describe two varieties of nervous gastric haemorrhage. The first he called hysterical, the second emotional, and he recorded the case of a man who, after being startled by the report of a gun, had profuse gastric haemorrhage. But the lack of direct control does not allow any conclusion to be drawn from it.

The research work of Professor Latarjet of Lyons may perhaps throw some light on this question of etiology. We are indebted to this anatomist for the most accurate study of the gastric nervous supply. He has performed many resections of the nerves of the stomach in cases of gastric crisis due to tabes, ulcer, and various gastralgias. Experimenting on dogs, he found that the resection of the proper nerves of the stomach is followed by very marked vasodilatation of the gastric wall, of the omentum, and of the peritoneum (going as far as ecchymoses), but the mucous lining did not partake in the phenomenon. Vaso-dilatation of the mucous lining is perhaps under the control of local centres, but we do not know at present what may be the track followed by the afferent nerves. These facts open up the question of the pathogeny of haematemesis—a vast field yet to be explored. We may admit that gastrostaxis is due to some perturbation of the normal reaction of the general nervous system on the visceral system, and we may also suppose that there are local circulatory circuits regulated by and, maybe, disturbed by the neighbouring organs. Thus might be explained haematemesis in cases of appendicitis and cholecystitis. We trust that physiologists will give us a key to the meaning of at any rate some of these cases.

We owe to Müller the location of sympathetic vasomotor centres in the tractus intermedio-lateralis of the medulla. A short time ago Mariano Castex, at the Paris Academy of Medicine, showed that the same centres command vasotrophicity (as has been proved in cases of purpura). We may admit disturbance of these nuclei, or of the nerve ganglions, or nerve endings, as a cause of haematemesis. In the present poor condition of our knowledge we can but mention the problem.

I am a disciple of the late Frantz Glénard, and therefore must be expected to mention hepatism as a possible cause of gastrostaxis. This great Vichy clinician, a worthy follower of Murchison, taught us the part played in pathology by small defects in liver-cell adequacy. Hepatism accounts for many forms of haemorrhage due to slight alterations of the vessels.

Further, if we admit that the immense majority of patients suffering from Glénard's disease are in fact what he himself called "des hépatiques," then we cannot fail to be very struck by the figures given by Mr. Walton in his recent paper on chronic dyspepsia in women. Mr. Walton writes: "Haematemesis is a frequent symptom in the gastric type, especially in women. In my own experience 50 per cent. of the cases have had a sharp loss of blood at some time or other of the disease. The loss of blood is manifestly due to the supervention of small gastric erosions in a dropped stomach." How pleased my old master would have been to read these lines! Mr. Walton adds: "Are such patients suffering from an organic lesion as well as their chronic dyspepsia? and if it be certain that they have no organic lesion, should they be treated by operation?"

These small erosions recognized by Mr. Walton in ptosis were described by Professor Dieulafoy, who, however, contrary to the British surgeon, used to teach that these minute ulcerations ran a rapid and acute course. Haemorrhage is the first symptom.

In one of the cases recorded above the possibility of syphilis is mentioned. It is often a cause of bands and kinks, and consecutive stenosis might quite likely be a cause of gastrostaxis—a fact to be kept in mind.

Prognosis is good. Out of 7,500 *post-mortem* examinations at Guy's Hospital Sir William Hale-White found only four deaths due to gastrostaxis. I have not been able to find

any French statistical data, but the best proof of the benignity of the symptom is that none of the old authors, speaking of vicarious gastric haemorrhage, ever considered it to be a very serious matter.

As regards treatment we have to distinguish the treatment of the condition and the treatment of the symptom. The treatment is surgical or medical. Neither Sir Berkeley Moynihan nor Lord Dawson recognizes any marked improvement due to operation, but Professor Duval's patients grew very much better. A curious explanation was offered. It was said that his incision happened to cut across a small ulcer which was afterwards stitched up without being recognized! It is difficult to admit the explanation in one case, and in two cases it is impossible. It has also been suggested that a long cut through the stomach wall produces enervation of the stomach—the operation of Latarjet.

This might be admitted, and as a matter of fact in one of Latarjet's cases the patient, who suffered from haematemesis, was cured after gastric enervation. We might accept as a working rule in practice that if x-rays do not give any definite information no operation is justified, at any rate in our present state of knowledge. Furthermore, a patient who has had a severe haemorrhage is in poor condition to stand the shock of any operation.

The medical treatment, as has been shown by Sir William Willcox, consists primarily in complete rest. Dr. Hurst gives a striking statistical table showing the low mortality of the condition. As the question has been exhaustively discussed in this house this very month, it seems useless to insist. I am only sorry that hydrologists did not have a word to say as regards preventive treatment. This side of the question might be the object of some interesting remarks.

CONCURRENT VICIOUS CIRCLES.

BY

JAMIESON B. HURRY, M.A., M.D.,

READING.

THE morbid process known as the vicious circle, or *circulus vitiosus*, is widely recognized as playing a role of fundamental importance in pathology. It leads to the aggravation of disease, to the destruction of organs, and frequently to a lethal exitus. Recent medical literature abounds in references to this process, which frequently elucidates obscure phenomena in disease, and permits of more rational therapeutics.

Medical writers, however, usually confine their attention to some single and isolated operation of this circular

process in any disorder under consideration. Thus they may draw attention to the reciprocal interaction of anaemia and disordered haemopoiesis, of anoxaemia and exhaustion of the respiratory centre, of varicose veins and thinning of the vein wall, of gastroparesis and kinking of the duodenum, of retroversion of the gravid uterus and retention of urine, of glaucoma and blocking of the filtration angle, etc. As a rule, however, such succinct statements only represent one out of a series of concurrent reciprocally acting factors; the processes of disease are much more complex than such summary sentences might indicate. In truth there are usually numerous simultaneous circles which unite in perpetuating the evil and frequently convert what would otherwise be a simple and self-limiting disorder into a grave and complicated one. As a stream

Cardiac failure may involve—

1. Impaired coronary circulation	→	Anaemia of myocardium	→	Weakened systole	→	Cardiac failure.
2. Pulmonary congestion	→	Anoxaemia	→	Enfeebled myocardium	→	Cardiac failure.
3. Bronchitis	→	Paroxysms of coughing	→	Enfeebled myocardium	→	Cardiac failure.
4. Venous stasis	→	Hepatic congestion	→	Malnutrition of myocardium	→	Cardiac failure.
5. Gastritis	→	Fermentation and flatulence	→	Embarrassed cardiac action	→	Cardiac failure.
6. Nephritis	→	Ascites	→	Embarrassed cardiac action	→	Cardiac failure.
7. Arrhythmia	→	Retarded circulation	→	Malnutrition of myocardium	→	Cardiac failure.
8. Anoxaemia of bulb	→	Peripheral vaso-constriction	→	Increased cardiac effort	→	Cardiac failure.
9. Auricular fibrillation	→	Lowered blood pressure	→	Malnutrition of myocardium	→	Cardiac failure.
10. Tachycardia	→	Lowered blood pressure	→	Malnutrition of myocardium	→	Cardiac failure.
11. Persistent palpitation	→	Anxiety neurosis	→	Insomnia	→	Cardiac failure.
12. Restricted exercise	→	Obesity	→	Increased cardiac effort	→	Cardiac failure.

I. Cardiac Failure.

Pulmonary tuberculosis may involve—

1. Cough	→	Loss of sleep	→	Lowered resistance	→	Pulmonary tuberculosis.
2. Cough	→	Dispersal of bacilli	→	Infection of fresh foci	→	Pulmonary tuberculosis.
3. Haemoptysis	→	Blood pressure raised by anxiety	→	Renewal of haemoptysis	→	Pulmonary tuberculosis.
4. Haemoptysis	→	Anaemia	→	Lowered resistance	→	Pulmonary tuberculosis.
5. Pyrexia	→	Emaciation	→	Lowered resistance	→	Pulmonary tuberculosis.
6. Pyrexia	→	Dryness of mouth	→	Anorexia	→	Pulmonary tuberculosis.
7. Formation of cavities	→	Mixed infection	→	Suppuration	→	Pulmonary tuberculosis.
8. Absorption of toxins	→	Retarded haemopoiesis	→	Lowered resistance	→	Pulmonary tuberculosis.
9. Tuberculous laryngitis	→	Dysphagia	→	Emaciation	→	Pulmonary tuberculosis.
10. Hyperhidrosis	→	Physical discomfort	→	Restriction of exercise	→	Pulmonary tuberculosis.
11. Loss of strength	→	Unemployment	→	Malnutrition	→	Pulmonary tuberculosis.

II. Pulmonary Tuberculosis.

Obesity may involve—

1. Extra demands on heart	→	Cardiac hypertrophy and dilatation	→	Restricted exercise	→	Obesity.
2. Deposit of fat in mediastinum	→	Weakened respiratory activity	→	Retarded oxygenation	→	Obesity.
3. Deposit of fat in liver	→	Hepatic enlargement	→	Loss of active habits	→	Obesity.
4. Deposit of fat in mesentery	→	Sluggish peristalsis	→	Excessive absorption from ingesta	→	Obesity.
5. Hypothyroidism	→	Low basal metabolism	→	Diminished katabolism	→	Obesity.
6. Sense of exhaustion	→	Boulimia	→	Overnutrition	→	Obesity.
7. Muscular weakness	→	Restricted exercise	→	Saving of calories	→	Obesity.
8. Increased somnolence	→	Increased hours of sleep	→	Saving of calories	→	Obesity.
9. Tendency to hyperhidrosis	→	Physical discomfort	→	Restricted exercise	→	Obesity.
10. Increased body weight	→	Tendency to flat-foot	→	Restricted exercise	→	Obesity.
11. Diminished energy	→	Self-indulgent habits	→	Restricted exercise	→	Obesity.

III. Obesity.

Constipation may involve—

1. Absorption of water from faeces	→	Formation of scybala	→	Dyschezia	→	Constipation.
2. Dilatation of rectum	→	Weakness of muscles of rectum	→	Dyschezia	→	Constipation.
3. Anorexia	→	Diminished ingesta	→	Diminution of faecal matters	→	Constipation.
4. Haemorrhoids	→	Painful defaecation	→	Postponement of defaecation	→	Constipation.
5. Fissure of anus	→	Painful defaecation	→	Postponement of defaecation	→	Constipation.
6. Loading of colon	→	Coloptosis	→	Dyschezia	→	Constipation.
7. Loading of sigmoid	→	Kinking of sigmoid	→	Dyschezia	→	Constipation.
8. Formation of rectal diverticula	→	Obstruction of rectum	→	Dyschezia	→	Constipation.
9. Restricted exercise	→	Weakened peristalsis	→	Dyschezia	→	Constipation.
10. Irritation of intestinal mucosa	→	Spasm of intestinal walls	→	Arrest of faeces	→	Constipation.
11. Absorption of toxins	→	Weakening of intestinal muscles	→	Arrest of faeces	→	Constipation.
12. Coprolith	→	Defective action of levator ani	→	Dyschezia	→	Constipation.
13. Absorption of toxins	→	Hypothyroidism	→	Myxoedematous infiltration of intestine	→	Constipation.
14. Anxiety neurosis	→	Perversion of mechanism of defaecation	→	Dyschezia	→	Constipation.
15. Anxiety neurosis	→	Resort to quack remedy	→	Irritation of intestinal mucosa	→	Constipation.

IV. Habitual Constipation.