Observations ON THE PRACTICAL VALUE OF THE SLIT-LAMP.*

BY

T. HARRISON BUTLER, M.A., M.D. OXON.,
SURGEON TO THE BIRMINGHAM AND MIDLAND EYE HOSPITAL,
HONORARY OPHTHALMIC SURGEON TO THE COVENTRY
AND LEAMINGTON GENERAL HOSPITALS.

(With Special Plate.)

In the nineteenth century the discovery by Babbage of the ophthalmoscope ushered in the era of scientific ophthalmology. Alvar Gullstrand has performed a similar service for the twentieth century; his slit-lamp has opened, not perhaps an era, but most certainly an epoch in the surgery of the eye.

It is probable that with the aid of our present methods alone progress will be increasingly slow, that we have learned almost as much as we can about the fundus and the media with the ophthalmoscope and the simple lens. These instruments are now supplemented by the slit-lamp and the eye microscope, which attack problems afresh, not only by furnishing a vastly increased magnification and introducing stereoscopic binocular vision, but by allowing us to illuminate structures from behind, and, above all, by enabling us to see the anterior half of the eye in *optical section*.

The advantages conferred by the slit-lamp are not only those inherent in itself, but also greater efficiency in the use of the older methods. This is due to two factors: we can to some extent apply slit-lamp technique to the ordinary oblique illumination with simple lens and condensing lens; and secondly, when we have become accustomed to the aspect of the cornea, the anterior chamber, the lens, and the vitreous, under the high magnification of the eye microscope and the focal illumination of the slit-lamp beam, we are better able to interpret the same subject seen with the smaller magnification of the loup, which in general magnifies about six times.

If, instead of using for oblique illumination the ordinary frosted bulb, we employ a 40-candle-power gas-filled half-watt lamp, or even a more powerful one at a considerable distance, we can apply the principle of focal illumination. We can make a ray of light pass into the eye and note that the cornea is illuminated, the anterior chamber dark, and the lens bright, and in the case of lens sclerosis we can even make out that the lens shows dissociation areas—that is, is laminated. If we make the patient fix his gaze between the light and our lens we can see the lens shagreen, and can make out the sutures of the surface of the lens cortex just below the capsule. If we move the shaft of light from side to side and note when any given opacity lights up, we can locate its position along the shaft of light, and so form a fairly accurate estimation of its exact place in the lens. We can direct our pencil of light upon the iris, and see the cornea by reflected light, by retro-illumination; and by throwing the beam into the

lens we examine the iris in lens light and make evident atrophic areas in the pigment layer of the iris, especially in the region of the pupil margin, the *collerette* of the French.

In a case of interstitial keratitis I have been able to see that the blood column is segmented in the smaller vessels, and that the position of the segments slowly moves, whereas in a larger vein I could perceive the actual blood stream, the contents of the vessel becoming moniliform and streaming along rapidly. In fact, the slit-lamp picture is reproduced. To see the blood stream with a simple lens it is essential to secure full dark adaptation and so to direct a strong beam that the light is reflected back from the iris and traverses the vessel. Finally, using a point of light lamp, I have been able to see the actual circulation of the blood in the limbal vessels of a normal eye. In this case the beam was accurately focused upon the sclera in such wise that it was reflected back through the conjunctival vessels. Under suitable conditions it is not difficult to see the vitreous fibrils with the simple lens.

I mention these facts to illustrate my point that familiarity with the slit-lamp increases our efficiency with the older instruments.

The slit-lamp and corneal microscope are shown in illustrations 1 and 2. The light source is a nitra-lamp with a thick helical filament. The light is projected by a lens system through a slit which is capable of adjustment to furnish a wide or narrow beam of light. The image of the filament is projected upon an illuminating lens which moves along a radial arm by means of a rack-and-pinion motion. This allows the beam to be brought to an exact focus. The lamp can be raised or lowered by a spring-

balanced screw, and the radial arm moves freely from side to side. The whole is attached to a table by a double arm designed by Henker. The table has a polished glass top upon which the microscope slides freely. As an alternative the microscope can be mounted upon a double slide like a large mechanical stage. Probably the glass top is the best in the hands of an expert. The microscope, shown in Fig. 2, was invented by Czapski, and as now made by Zeiss is an adaptation of their well known prismatic dissecting microscope. It is capable of magnifications ranging from 9 to 108. A chin and forehead rest is attached to the table. Both table, chin rest, and microscope have a similar screw adjustment to the slit-lamp, which allows the patient to rest his head comfortably and the surgeon to direct the beam of light to any part of the eye at will. It is essential to have the correct instrument. I fear that some have been persuaded by the makers to buy the combined slit-lamp and simplified Gullstrand ophthalmoscope. Like most combination tools, this arrangement is inferior to the real slit-lamp.

The illuminating lens should, for ordinary clinical work, be the small Vogt lens with a focal length of 10 cm. The larger aspherical lens is not achromatic, and does not give quite the same definition of the corneal prism and of other structures. On the other hand, it collects more light, and in consequence is better for the examination of the vitreous.

I find that I practically use one magnification only—objective A2 and ocular 2. This combination enlarges 23 times. I do not find the magnification of 9 of great value, and I rarely use the objective A3, which with ocular 2 magnifies 35 times. With a magnification of 23 the apparatus costs £65, and this specification is enough to

[3309]

FIG. 1.—The corneal microscope and Gullstrand's slit-lamp.

FIG. 2.—Microscope head with Dr. Ulbrich's graduated drum attached.

* A paper read to the Ophthalmological Section of the Royal Society of Medicine, January, 1924.

begin with. As the observer becomes more proficient higher powers and adjuncts can be added from time to time. The resistance should be adjusted to furnish 8 volts, but the nitra-lamp will stand more for a time, and can be pushed when more light is needed to examine the vitreous.

The examination should be made in complete darkness, and the more difficult subjects will be appreciated only when the eye becomes dark-adapted. It is not desirable to take the patient to the dark-room, look at his eye with the slit-lamp, and then return to work which demands light adaptation. It is better to examine the patients in batches if the best observations are to be made. Dark adaptation is especially necessary in the search for cellular elements in the aqueous. It is important that the microscope be accurately adjusted for the observer's intrapupillary distance. The slightest error here will disturb the stereoscopic effect and produce eyestrain. Very often the merest touch will produce the desired effect. There is little or no discomfort involved in the examination; the majority of patients will tolerate prolonged sittings even when suffering from inflammatory states, but some have a definite time limit in this respect. If we are working with the narrow slit the patient is almost unconscious of the light. Quite young children make excellent subjects. The pupil does not contract much under the slit-lamp, and I make most of my observations without any mydriatic. A complete examination of the lens and vitreous calls for a dilated pupil, but in general we can learn all that we require with a normal pupil.

The slit-lamp and eye microscope are not easy to use; in fact, the instrument is as difficult as the ophthalmoscope, if not more so. We have all of us had careful and methodical instruction in the use of the ophthalmoscope, typical cases have been selected for us, and there are several excellent atlases dealing with the subject. It is even more necessary to receive definite instruction in the use of the slit-lamp with demonstration of typical conditions. Practice should begin upon pigs' eyes and continue upon the human eye. If, however, it is not possible to attend a suitable course, such as Professor Vogt's at Zürich, much may be done by reading up the subject, but unfortunately all the important literature is in German. There is a little in French, and a few articles will be found in American journals. Localization is ably dealt with by Vogt in a paper translated in the *British Journal of Ophthalmology* for March, 1924. Vogt's *Atlas of Slit-Lamp Microscopy* has been translated into English. Koeppe's valuable work on "*The Microscopy of the Living Eye*" has not been translated. This is a book which is absolutely necessary if we are interested in the microscopical side of the study, as it deals largely with appearances seen under magnifications of 60 and over. Gallimerts and Kleefeld have published an interesting series of articles in the *Annales d'Oculistique* for 1919 and 1920. They are well illustrated, but are somewhat out of date, and as regards the lens wholly so. There are many valuable and magnificently illustrated articles by Vogt in the *Archives of Ophthalmology* for the past eight years, but again all in German, and in somewhat involved language.

I wish to emphasize the fact that the slit-lamp in no wise supersedes our older technique. It supplements and amplifies it. When we have done all that we can with ophthalmoscope and simple lens we can carry our investigation much further, with the result that much that was problematical becomes clear: we can pass from what was often at best good guessing to certainty. I propose now to show that this is in fact the case and to indicate how the slit-lamp has helped me in my routine clinical work.

When the broad beam of the slit-lamp is focused upon the eye it first illuminates the cornea, producing a prism of light which will be described in detail later in this paper. The anterior chamber remains dark under normal conditions. The lens becomes illuminated, and behind it the vitreous fibrils can be brought into focus. When the slit is narrowed down the prism becomes a mere illuminated slice of the tissue under focus; we have in effect an optical section. The appearances presented are seen in Figs. 3 and 4. The first shows the effect with the broad ribbon of

light, the second with a narrow beam. It will be noted that in Fig. 2 there are two prisms, the corneal and the lental; the lental is not so sharp as the corneal.

We can now proceed to examine the structures of the eye in detail.

Cornea.

When the broad beam is focused upon the cornea we have, as already stated, a belt of illuminated tissue which may be regarded as a curved prism. Fig. 5 shows this prism magnified 35 times. It will be noted that it has two easily visible sides and three edges. In the young and normal cornea the fourth edge is difficult to see, but careful focusing will reveal it. It is faintly shown in the drawing. The light must be imagined to come from the left side. The left-hand surface is the direct reflection from the corneal surface; we can call it the epithelial surface. If a drop of cocaine be instilled, followed by another of fluorescein, the epithelial surface will be easily differentiated, for the fluid upon it will be stained green. The right-hand surface seen in the drawing is the section of the cornea. The prism has two more sides—the endothelial, and the lateral section which is not in view. It is quite obvious that observation of the section side will enable us to judge with reasonable accuracy the position of an opacity or structure in the cornea. We can say that it is on the epithelial surface, within the substance of the cornea, or on the endothelial surface. We can move our prism from side to side and trace out the configuration and position of the subject in question. The exact method of accurate localization is described by Vogt in the paper already mentioned.

If we now narrow down our beam of light we can almost abolish the epithelial and endothelial surfaces and work with a pure section. This section gives a clear indication of any irregularity in the corneal surface or marked change in its curvature. Fig. 6 shows the effect produced by a corneal ulcer; Fig. 7 the heaping up of tissue which has followed the healing of an ulcer. Fig. 8 is the section of a conical cornea; the thinning of the apex is well demonstrated. The white lines like threads of cotton crossing the prism in Fig. 5 are the corneal nerves. These never penetrate into the deeper lamellae.

If, when examining a case, we make the patient fix his gaze half-way between the axes of the lamp and microscope, we can examine structures close to the reflected light, in specular or mirror light. By this method we are able to see the endothelial cells of the cornea. They form a hexagonal mosaic of a yellowish tint. They are shown in Fig. 9.

I have found the slit-lamp of great service in the case of foreign bodies embedded in the cornea. It is valuable to know exactly how deep they have penetrated and how they lie. With this knowledge we can plan the method of extraction and remove them with the least damage to the cornea. When there is a history that an eye has been struck by a flying fragment of metal we can determine whether the cornea has been actually penetrated. In one case in which I suspected that a fragment might have been retained in the eye I was able definitely to decide that the wound had reached only to the centre of the corneal stroma and had not passed into the anterior chamber.

Many diseases cause profound changes in the appearances of the cornea. Figs. 10 and 11 are taken from a case of herpes ophthalmica. Fig. 10 is the view with the simple lens; it shows keratitic precipitates (K.P.) and a fine striation in the lower segment. Fig. 11 shows the appearance of these striations in the corneal prism, magnified 35 times. It will be noted that the white punctate deposits are quite superficial in the prism. Vogt says that this appearance is characteristic of herpes, and it is obvious that in this case the slit-lamp is of immense value in diagnosis. Fig. 12 shows the slit-lamp view of the surface of the cornea in a very severe example of corneal herpes which has ended with calcification of the cornea. An iridectomy had been performed by Mr. Maddox.

It is in inflammatory states of the eye that the slit-lamp is of special value. Slight irritation or rise of intra-ocular tension causes a bedewing of the corneal surface, either the epithelial, the endothelial, or both. Under the high magnification employed the appearance suggests that of

FIG. 3.

FIG. 4.

FIG. 5.

FIG. 6.

FIG. 7.

FIG. 8.

FIG. 9.

FIG. 10.

FIG. 11.

FIG. 12.

FIG. 12A.

FIG. 13.

FIG. 14.

FIG. 15.

FIG. 16.

FIG. 17.

FIG. 18.

FIG. 19.

a mirror in a steaming bathroom. Vogt calls the condition "*betanung*." The appearance is normal at the corneal periphery, but in abnormal circumstances it may cover the whole cornea. It is practically always found when the anterior chamber contains absorbing lens cortex. Vogt believes that the posterior bedewing is due to alterations in the endothelium, but Koeppe considers that it is due to cellular deposits. It is difficult to believe that an appearance which so frequently accompanies the slightest irritation of the eye can be caused by a deposit of cells. When the condition is universal it is almost impossible to decide whether it is epithelial or endothelial, but when it is confined to a definite area it can be localized by the ordinary methods described by Vogt.

Deposits on the endothelial surface, the so-called "K.P.," can be accurately localized and detected when the particles are too small to be seen with the simple lens (loup), but when they have been seen under the slit-lamp they can often be detected with the loup. These fine deposits can be seen both by reflected and incident light, and can be investigated by an alternation of both methods. This fine K.P. can often be found after operations on the globe which show no signs of inflammation when examined by the ordinary methods. The black variety is merely iris pigment and has no significance. Often (after a cataract extraction) there is a deposit which has the appearance of the dangerous "mutton-fat" K.P., but the eye microscope shows that it merely consists of flakes of absorbing lens cortex, and this again is harmless. Koeppe, working with a magnification of 60 and over, professes to be able to classify the varieties of keratitic deposits, and from his examination to determine the nature of the inflammation.

In most deep corneal inflammations Descemet's membrane is thrown into folds, and occasionally Bowman's membrane is folded. Fig. 12A shows folds in Descemet's membrane which were noted after a normal preliminary iridectomy. They had entirely disappeared in two days. Such folds are almost constant after any operation upon the cornea, but exactly similar ones are seen in keratitis. A central keratitis may leave permanent radial folds unaccompanied by any marked deterioration of visual acuity; an example is shown in Fig. 13. In Fig. 14 a central keratitis, probably tuberculous in nature, has left folds of both Descemet's and Bowman's membranes. The fine star is due to the folds in Bowman's membrane. In this case the slit-lamp cleared up a very doubtful diagnosis.

A case of interstitial keratitis cannot be completely interpreted without the corneal microscope and slit-lamp. The extent and position of the vascularization can be mapped out, and by examining them by retro-illumination we can at once decide whether the vessels are still functioning or not. Many a case appears to be cured, the vessels are regarded as absorbing and functionless, but the slit-lamp shows that they still carry a rapid blood stream. Folds of Descemet's membrane are frequent and varied in character.

Anterior Chamber.

Observation of the aqueous is difficult and calls for complete dark adaptation. It is best carried out by allowing the beam to pass through a pin-hole. We now obtain not a ribbon but a pencil of light. This pencil illuminates the cornea and the lens, but should be invisible in the anterior chamber. If, however, the aqueous contains albumin or is turbid the pencil of light is visible. If bright particles flash across the beam then the aqueous contains discrete particles, cellular or otherwise. The movement of the particles is due to the heat eddy in the aqueous. The difference in temperature between the cornea and the iris causes a constant stream of aqueous in the anterior chamber, rising in front of the iris and falling behind the cornea. The heat of the slit-lamp beam induces vortex currents, and these cause the particles to move across the beam of light and become visible. The presence of cells in the anterior chamber is often the first sign of a sympathizing ophthalmitis. If in a case in which we have reason to anticipate sympathetic ophthalmitis we detect cells in the aqueous, we have a timely

warning and can act accordingly. I have already saved an eye which I should not have dared to keep but for the fact that daily examination with the slit-lamp failed to detect any signs of inflammation in the other eye.

Iris.

The microscopy of the living iris is straightforward and is in the main merely an enlargement of what we see with the simple lens. We have, however, the great advantage that we can examine the iris in lens-light, by retro-illumination, and detect atrophy and ruptures of the pigment layer of the iris. Koeppe thinks that a deposit of fine granules of black pigment in the iris stroma is characteristic of glaucoma. It is true that it is very often seen in the glaucomatous eye, but it can sometimes be noted in a normal senile eye.

Lens.

The slit-lamp is of most value in the examination of the lens, and it is in this region that it has won its greatest triumphs. The optical section made by the narrow ribbon enables us to see the lens in profile exactly as though the section had been made with a knife. It is at once obvious, even to the naked eye, that it is not homogeneous, but shows what Vogt calls dissociation areas: it can be separated into definite lamellae. Starting from before backwards, we can recognize the following regions: The capsule shining as a thin bright band; behind it, separated by a dark area, is another line very similar to the capsule, except that it has a stronger curvature and is more opaque and generally slightly yellow in tinge: this we can call the reduplication line. The next area is the cortex, followed by the well marked surface of the adult nucleus, which again has a shorter radius of curvature than that of the reduplication line. The next layer is called by Vogt the outer embryonic nucleus, but, whereas it is present at birth, I propose to call it the infantile nucleus; this again has a greater curvature than the adult nucleus. I may remark that in the translation of Vogt's *Atlas* the word "*alterskern*" has been rendered as *senile* nucleus. This gives an entirely wrong impression, for this nucleus appears at about 10 years of age, and is in no sense senile. *Adult* nucleus is better. The centre of the lens is occupied by two plano-convex bodies always separated by a sharply defined, broad, dark line. This is the foetal or embryonic nucleus. The flat sides of this internal lens system face each other. On the anterior convex surface we find the anterior upright Y, the primary foetal suture of the lens, and on the posterior surface there is an inverted Y (λ). In 25 per cent. of all cases there is a small opacity on the flat surface of the anterior body. This can easily be mistaken for a cataract.

Not only are these nuclei sharply differentiated in optical section, but when focused under the illumination of the broad ribbon they give mirror effects, and show up the characteristic figures of the sutures, and often it is possible to see the actual lens fibres. Lamellar differentiation becomes more obvious in the old lens, and especially when there is sclerosis of the adult nucleus.

It is obvious that the power to separate the lens into definite anatomical regions greatly enhances our powers of diagnosis and prognosis, and has an influence upon treatment. I now make a practice of examining every case of cataract with the slit-lamp, and I have obtained great help in diagnosis, in determining whether the lens is fit for extraction, and in selecting the appropriate technique for the operation. Baracquer makes a careful slit-lamp examination of the lens before he uses his suction apparatus, and by so doing he is able to form an opinion as to the conditions he will meet, and to decide whether the case is suitable for the suction operation.

Fig. 15 shows the normal lens in optical section. The drawing represents the lens unduly flat, but in other respects it shows a common type. Figs. 16 and 17 portray the extraordinary appearance that is sometimes seen on the surface of the adult nucleus. The patient had fair visual acuity, and to ordinary examination was an example of subcortical cataract. The slit-lamp showed that the surface of the

nucleus was studded with elevations like a hobnail boot. In section it is seen that they stand out into the cortex, and that each throws a dense shadow back into the lens. The surface of the nucleus is strongly condensed, giving the appearance of an internal capsule. The cortex shows lacunae and opacities which are typical of subcapsular cataract. It will also be noted that the nucleus has a stronger curvature than the capsule, and being sclerosed it caused a lental myopia; it was yellow in colour. The myopia in this case was 5 dioptries. The nipple-like prominences on the adult nuclear surface are illustrated in Vogt's *Atlas* in a series of photographs from plastic models made by Dr. Lüssi, late chief assistant at the Basel Eye Clinic.

Lamellar cataract forms an interesting study and takes many forms. Fig. 18 shows a fairly typical example. It will be noted that there is a condensation within the adult nucleus, and that the infantile nucleus is of different texture from the rest of the lens. It shows a scintillating mass of dots and in some lights has a stony appearance. It is here that the chief opacity to light is found. Instead of passing on, light is reflected from the sparkling particles. It is obvious that in this case there have been two periods of modified nutrition of the lens—one at birth and the second during the first ten years of life. These changes are comparable to ridges on the nails and teeth after illness. In this drawing the central plano-convex embryonic nuclei are very evident. At the back of the lens it is sometimes possible to see a ring or concentric rings—the “*bogenlinie*” of Vogt. To this is attached the remnant of the hyaloid artery, which the slit-lamp shows to be present in about 25 per cent. of all cases. The arc-line is probably the spot to which Cloquet's canal was attached. These structures are never found exactly at the posterior pole, where they would form an impediment to accurate vision, but on the nasal side. An example is shown in Fig. 19; this was seen in a child of 10 and was symmetrical in each eye. The posterior inverted “Y” is seen, and radiating from it the lens fibrils. Just at the bifurcation of the nasal branch the arc-line is seen, and above it a second curved line. From the circle depends the corkscrew-like remnant of the hyaloid artery. Behind the lens capsule we see the normal vitreous fibrils.

Vitreous.

A complete examination of this structure calls for special apparatus. It is best to use an arc lamp instead of the nitra-lamp, and Koeppe's silver mirror; but much may be seen with the ordinary apparatus if the axis of the lamp is brought as close as possible to that of the microscope. The correct interpretation of what one sees in the vitreous calls for much experience. In some cases the vitreous seems to be almost optically homogeneous, in others it is full of floating fibrillae, of membranes, and particles, and yet is perfectly transparent.

Retina.

The retina can be examined with the slit-lamp, but it is necessary to place a contact-glass upon the cornea to abolish its curvature, and to use the micro-arc lamp. Such investigations have great value from the standpoint of research, but they hardly come within the scope of ordinary clinical work. In cases of obscure tumours deep in the eye this technique may render valuable aid in diagnosis, but it is exceedingly difficult and also unpleasant to the patient.

It is self-evident that the slit-lamp and corneal microscope have greatly enlarged our powers of diagnosis and research. The anatomy of the lens has been elaborated to such an extent that even the most modern textbooks are obsolete in this connexion. Our knowledge of the vitreous has greatly advanced; in fact, before the advent of the slit-lamp its minute structure was unknown.

The great drawback to the use of the slit-lamp is that it definitely increases the time taken by an examination. It is time well spent, for the investigation adds enormously to the interest and value of our work. It is lack of time and of an adequate staff in our ophthalmic clinics which makes it so difficult to develop clinical research work comparable in volume to the output of our Continental colleagues.

ALCOHOL AND LIFE DURATION.

BY

RAYMOND PEARL, PH.D., D.Sc., LL.D.,
PROFESSOR OF BIOMETRY AND VITAL STATISTICS IN THE SCHOOL OF
HYGIENE AND PUBLIC HEALTH, AND OF BIOLOGY IN THE
MEDICAL SCHOOL OF THE JOHNS HOPKINS UNIVERSITY,
BALTIMORE, MARYLAND, U.S.A.

THE problem of the effect upon duration of life of the use of ethyl alcohol as a beverage is still open. There is abundant evidence that *excessive* use of alcohol shortens life, but, because of the difficulty of getting critically adequate data, it has not been possible up to this time to determine whether the consumption of alcoholic beverages in *moderation* has any effect upon life expectancy, or, if it has, what the nature and sense of that effect is. The evidence which has been adduced as bearing on the point, chiefly from the records of insurance companies, can easily be shown, and repeatedly has been shown, to have no probative significance so far as the problem just stated is concerned. The case has been well put in the concluding paragraph of Chapter X of the second edition of the report¹ of the Medical Research Council's Alcohol Investigation Committee (p. 148), where it is said:

“We repeat that it does not follow from these conclusions that a consistently moderate use of alcoholic beverages shortens life, since we cannot sort out from the data brought into comparison those instances in which the habitual abuse of alcohol produces disease or death, or in which accidental abuse has a like consequence, nor can we exclude many other social factors operating for or against longevity. . . . What we have not proved, and, from the evidence before us, cannot prove, is that the moderate use of alcoholic beverages, employing the word ‘moderate’ as it is ordinarily used in everyday life, has any effect at all, one way or the other, upon longevity.”

In 1922 I prepared a preliminary report² upon a collection of data which may fairly be said, I think, to have unique value for the study of this problem. In brief this material consists of records of certain pertinent data in the life-histories of many thousands of persons belonging to the working class population of Baltimore. These records were collected by trained eugenic field workers who visited the families and by long-continued and patient investigation got the facts with a high degree of accuracy. Proceeding in the matter in an entirely objective way and without preconceptions, it was felt to be essential, if any critical results were to be obtained, to get the most elaborate and critically exact records possible about the environmental situation, the habits of life, the health history, the racial stock, anthropological characteristics, exposure to tuberculosis, etc., of all the members of the family group.

Every critical safeguard of the accuracy of the ultimate individual records that I have been able to think of was thrown around this work. No fact was entered in a history finally until it had been corroborated by the independent testimony of at least two persons acquainted with the individual in question. The material was all taken from one socially and economically homogeneous group of the population of the city of Baltimore—namely, what might inclusively be called working men's families. By the employment of field workers speaking a variety of foreign languages it was possible to get representation in the histories of the different foreign race stocks in about the proportion that they are represented in the total population of the city of Baltimore, and in this way there has been no racial discrimination in the study. The only exception to this is that the negro has not been included, owing to the difficulty—indeed, impossibility—of getting accurate genealogical information about negro families. These people do not know accurately about their own ancestors.

Special attention in these inquiries was devoted to the habits of each individual, throughout life, in respect of alcohol and tobacco. I have given elsewhere (loc. cit., pp. 237-246) a detailed account of the material, its mode of collection, the care taken to ensure its critical accuracy, etc., and these details need not be repeated here. It will suffice to state here that for a group of over 6,000 individuals we have accurate information covering the following points: