The left eyelids and conjunctiva were very oedematous. The left cheek, over the parotid region, was red and swollen. All the affected parts were somewhat tender. Temperature and pulse were normal; the tongue was very coated; the pupils were equal and active. All other reflexes were normal. On the chest and abdomen there was a rash, which had been present about twenty-four hours. The clinical features of this rash, throughout its duration, were those of varicella, but, in the light of subsequent events, the question arises whether the rash was due to varicella or to larval infection. Within forty-eight hours the swelling of eye and cheek had completely subsided, the tongue was clean, and the child was asking for food.

He remained quite well until March 14th, when the right eyelids and cheek became swollen and red. There were also several circumscribed swellings—one the size of a chestnut on the right side of the forehead just above the outer canthus, one the size of a hazel-nut behind the left ear, one the size of a walnut over the right parietal region of the scalp, and two small ones over the left parietal region. The swelling on the right side of the scalp was semifluctuating, the others were moderately firm. None of the swellings was red or painful. In a few days, with the exception of the scalp swellings, the parts returned to their normal condition and the child again seemed to be quite well.

On March 24th, after severe pain in the head, the oedema and redness returned, this time in both eyes and cheeks. Next day, from the right parietal swelling, which was now discharging pus, came a maggot about half an inch long. Within a few hours the swelling of eyes and cheeks had completely disappeared.

On March 27th a maggot was discharged from one of the two swellings on the left side of the scalp, and from the other swelling a maggot was removed on April 1st; both of these maggots were about a quarter of an inch long. There have been no further developments.

Curious features of the case are:

(a) The regular intervals between the appearance of the swellings. There were five distinct attacks with about ten days between each.

(b) Their symmetrical situation. The first swelling was in the left eye; the second in the left eye; the third in the right eye and cheek; the fourth in the left eye; the fifth in both eyes and cheeks. One maggot came from the right side of the scalp and two from the left side of the scalp.

(c) The rapidity with which the oedema subsided.

The swellings noticed on March 14th, one above the outer canthus of the right eye and the other behind the left ear, are suggestive of halts in the progress of the maggots either to or from the scalp.

Can the periodic swellings be accounted for by assuming that eggs were deposited and hatched in the scalp and that the larvae migrated, at intervals, to the orbital tissues, remaining there a few days only? The remarkably rapid onset and subsidence of the oedema seem to point to some such transient exciting cause.

I have ascertained that warble-flies are prevalent on the farm where the child lives, so that the source of infection is obvious.

Report by Major Austen.

The first maggot was sent to Major E. E. Austen, Natural History Section of the British Museum, who very kindly furnishes the attached report, which he permits me to use.

"The specimen is a penultimate stage larva of *Hypoderma lineatum*, Vill., which is one of the two British cattle warble-flies, the other being *Hypoderma bovis*, De G.

"This is the first British case of myiasis in man due to a warble-fly larva that has come to my personal knowledge, but a number of instances which have occurred on the continent of Europe and in the United States of America, where *Hypoderma lineatum* is also found, are on record.

"The preponderance of cases in which the victims of the fly's attack have been young children is noteworthy, but is perhaps to be correlated with the fact that in cattle young animals are usually most severely warbled.

"Another interesting point in your case arises in connexion with the way in which the patient became infected. The egg from which the larva developed must have been laid last summer, probably between July and September, and, if not actually deposited on a hair of the child's head or body, was in all probability laid in the ordinary way on a cow's hair, and then swallowed in milk.

"Under normal conditions, the eggs of both *Hypoderma lineatum* and *H. bovis* are attached by the parent flies to hairs on the legs and flanks of cattle, and the minute larvae, which hatch out a few days later, crawl down the hairs and enter the body, burrowing through the skin by way of the hair follicles. A period of many months of wandering follows, during which the larvae travel about the body through the connective tissue, and, before taking up their final position in 'warbles' beneath the skin of the back, are met with in situations so diverse as the wall of the oesophagus and the spinal canal.

"The full-grown larva is a grub of considerable size, and is many times larger than the specimen forwarded."

The following references to other published cases of this condition have been provided by Major Austen.

Riley, C. V.: "A grub supposed to have travelled in the human body." *Insect Life*, Washington, U.S.A., vol. ii, p. 238 (1890).

Condorelli, Mario: "Caso di 'myiasis' nell'uomo per larva cuticolare di *Hypoderma bovis* (De Geer)." *Boll. Soc. Zool. Ital.*, Anno xiii, pp. 171-181 (1904).

Topsent, E.: "Sur un cas de myase hypodermique chez l'homme." *Archives de Parasitologie*, T. iv, p. 609 (1901). Larva of *H. lineatum* beneath skin of upper part of thorax of girl of 8 years.

Topsent, E.: "Second cas de myase hypodermique dans le Morbihan." *Op. cit.*, T. xiii, pp. 438-489 (August 25th, 1903). Larva of *Hypoderma* beneath skin of face of girl of 6½ years.

Miller: "Larvae of *Hypoderma lineatum* in the human subject in U.S.A." *Journal of the American Medical Association*, December 3rd, 1910 (abstract in the *Journal of Tropical Medicine and Hygiene*, January 2nd, 1911, p. 13).

ETIOLOGY AND TREATMENT OF RHEUMATOID ARTHRITIS AND ALLIED DISEASES.

BY

S. G. BILLINGTON, M.B., B.S.LOND., F.R.C.S.E.,
PATHOLOGIST, BACTERIOLOGIST, AND HONORARY ASSISTANT
PHYSICIAN TO THE WOLVERHAMPTON AND
STAFFORDSHIRE HOSPITAL.

WITH ELECTRO-THERAPEUTIC NOTES

BY

SANDISON CRABBE, L.R.C.P. AND S.,
ELECTRO-THERAPIST TO THE BIRMINGHAM HOSPITAL FOR WOMEN.

THE term "rheumatism" is very casually applied to the signs and symptoms of inflammation of joint, nerve, muscle, or fibrous tissue. The first essential step is to determine the cause of the inflammation.

The causes of the so-called rheumatic conditions fall into line with the usual causes of inflammation—physical (mechanical or traumatic), chemical (metabolic, gouty, diabetic, alcohol, etc.), thermal (heat or cold), specific (syphilitic, neuropathic, haemophilic, and the specific bacterial such as typhoid, gonococcal, etc.) and non-specific bacterial.

The remarks here made will be confined to the non-specific bacterial group, though the others must be mentioned from the point of view of differential diagnosis.

Articular rheumatism, which is probably the most common variety, and certainly the most important from the point of view of disability caused, can be divided into two great groups:

1. *Rheumatic Arthritis*.—Usually polyarticular and associated with pyrexia, often accompanied by an endocardial lesion and generally controlled by salicylates. The joint lesions are characterized by painful and tender synovial and periarticular swelling which may appear and disappear comparatively rapidly.

2. *The Rheumatoid Group* (and in this I include the various types of rheumatoid arthritis and osteo-arthritis).—This variety may be polyarticular or monarticular, and acute, subacute, or chronic. The acute arthritis may develop as a primary disease or follow an acute febrile condition such as influenza, or result from a sudden exacerbation of the chronic. The chronic form may develop as such or supervene upon the acute.

The rheumatic affections of nerves, muscles, and fibrous tissue closely correspond, except that the acute varieties are less common if we exclude the nerves. In our experience the underlying causes are the same: which tissue is involved depends largely upon personal or regional susceptibility and various contributory causes.

A very high percentage of rheumatic affections may, for practical purposes, be regarded as metastatic manifestations of focal bacterial absorption. In a small proportion of cases exciting causes, such as nervous shock, worry, trauma, etc., appear to play an important part. In another small proportion the disease supervenes upon a febrile attack, though frequently in these cases a local septic focus has been lighted up.

The pathological varieties cannot be considered in this communication; but the fact may be emphasized that the

disease represents a definite sequence of events, and that the treatment usually required is composite, illustrating well the advantages of team work.

Three stages of rheumatoid arthritis can be recognized: (1) early, with slight or recently developed joint lesions; (2) active arthritis with definite arthritic changes; (3) old-standing arthritis, with or without the presence of the active disease, but with post-arthritic deformities, contractures, muscular atony, and wasting.

The treatment of the disease may be considered under the following headings:

- A. Removal of the infective focus.
- B. Neutralization of bacterial toxæmia in blood stream and affected tissue.
- C. Orthopaedic treatment of the results of the disease.
- D. Psychological treatment also would be advantageous in many cases.
- E. The general treatment of the patient by tonics, endocrine preparations, etc., must also be kept in mind.

In Stage 1 removal of the infective focus is usually all that is necessary, and the disease rapidly subsides when this has been done. In Stage 2 the toxæmia must be neutralized in the affected tissues after removal of the infective focus. In Stage 3 the treatment is more complex and falls under the headings of:

- (1) Removal of the infective focus if present.
- (2) The treatment of active arthritis if present.
- (3) The treatment of post-arthritic sequelae by electricity, massage, manipulation, exercises, or operation.
- (4) The use of a vaccine as a prophylactic against lighting up the disease by the above orthopaedic procedures.
- (5) General and psychological attention to the patient.

DIFFERENTIAL DIAGNOSIS.

The first essential in treatment is correct diagnosis, both as to the type of arthritis and as to the stage of the disease. In about 25 per cent. of the cases referred for antibacterial treatment the diseases have been non-bacterial in origin, the conditions ranging from flat-foot to congenital syphilitic synovitis.

The diagnosis must be made from diseases of metabolic or chemical origin, such as gout, diabetes, and alcoholism; from mechanical conditions resulting from trauma, occupation, pressure of new growth, etc.; and from displacements, such as flat-foot and coxa vara. Syphilitic, neuropathic, and the specific bacterial affections require separate consideration.

The diagnosis between active arthritis and post-arthritic sequelae is most important, the stage of the disease being often imperfectly realized. Patients have been subjected to courses of vaccines when the entire disability consisted of ankylosed joints with contractures, there having been no signs or symptoms of active arthritis for a number of years. The most difficult post-arthritic condition to distinguish is traumatic arthritis or synovitis resulting from apposition of roughened or cartilage-denuded joint surfaces. The distinction is that the pain and swelling in the mechanical forms occur during and after movement, and become more pronounced the more the joint is used, whereas within limits the pain and stiffness of active arthritis diminish with moderate use.

The three most important evidences of active arthritis are: pain apart from movement, the night pain being most characteristic; local tenderness; and general debility with often a toxæmic facial aspect.

TREATMENT.

In the *first stage* all that is required is to search for and remove the focus of origin. The commonest focus is undoubtedly dental, and if there is no obvious sepsis the teeth should be x rayed to exclude the presence of apical sepsis. The x ray is not infallible and several cases have been seen secondary to thickening of the periodontal membrane; but except in root-filled or extensively filled teeth the x ray is very reliable.

After the teeth the commonest sites of infection are the tonsils and accessory nasal sinuses. The gall bladder is a much commoner focus of infection in rheumatoid arthritis

and similar diseases than is generally supposed. The genito-urinary system may contain the focus, the uterus in the female and the prostate in the male probably being the commonest sites.

In a small percentage of cases of rheumatoid arthritis no focus can be found and the physician is forced to consider the indefinite intestinal source of origin. The appendix is a probable focus in some cases. As intestinal antiseptics dimor or salol and kaolin have been used, and endocrines when intestinal atony is definite.

In the *second stage* also the focus must be sought for and dealt with. The systemic bacterial toxins must then be neutralized by specific or non-specific remedies. Many drugs are alleged to exert some beneficial effect, such as guaiacol carbonate, iodides, collosol manganese, and alternating collosol iodine and collosol sulphur. In some cases drugs unquestionably do exert a beneficial effect, but their action is uncertain; it would appear to be due to their ability to neutralize the toxins of some bacteria in some individuals, while in blood or tissue fluids of slightly different composition the toxins of other bacteria are not neutralized.

Peptone or other foreign proteins may be used, and may produce very striking improvement. Generally speaking, however, foreign proteins are efficient only when given in sufficient amounts to produce severe febrile reactions; they make the treatment very unpleasant, and necessitate a good deal of incapacity and bed during its course.

The most reliable antibacterial agent is an autogenous vaccine, and if sensitized it can be pushed without producing any reaction or incapacitating the patient in any way. For over four years vaccines made from the blood have been used, either the auto-hæmo-vaccines,¹ or vaccines made from definite bacteria grown from the blood in ammonium carbonate, citrated glucose broth;² from those sources the correct antidote for the causal organism can more or less be guaranteed.

The antibacterial action of the vaccine can be greatly aided by electrical measures, which are used (1) to produce vaso-dilatation, in order to increase the antibody supply to the affected joint and hasten the absorption of effusion and oedema; (2) to exert direct antibacterial action; and (3) to help restoration of function by softening and aiding absorption of adhesions, and stimulating atonic muscles.

To produce surface vaso-dilatation the high candle-power lamp (1,000 c.p.) is the best; to promote deep hyperæmia, diathermy and the static brush effluve are the most suitable. To remove oedema and fluids the static wave is the best. It causes intermittent contraction of the arterial musculature and thus produces capillary massage. Diathermy and the galvanic current also aid the removal of effusions.

The tungsten arc lamp develops antiseptic rays, while high frequency, administered either with the glass vacuum tube or through the hand with a cotton glove on while giving massage, is useful.

For ankylosis and contractures the galvanic current, which has sclerolytic properties, is the most useful aid to massage and exercises. The static wave will contract the wasted and atonic muscles while the patient sits in a chair. Rhythmically broken galvanic current is a useful stimulant but liable to irritate the skin.

The operative measures that may be adopted are wrenching under an anaesthetic; or open operation for removal of osteophytic outgrowths and hypertrophied synovial membrane; or excision of the joint may be practised.

The psychological aspect in the third stage of rheumatoid arthritis is often very difficult to treat, and frequently patients are bedridden with perfectly quiescent arthritis. They think that they are unable to get out of bed, and they never will unless they can be persuaded that nothing but their own efforts will get them out, and that those efforts will be successful if sufficiently determined. At such a stage a vicious circle is apt to be established, the increasing incapacity leading to more pronounced metabolic and intestinal stasis, with consequent aggravation of the rheumatic disease. Deficiency of endocrine output and achlorhydria frequently accompany rheumatoid arthritis, probably representing other manifestations of the general toxæmia, and being aggravated by the continual pain. I have found pluriglandular preparations most useful

adjuncts in treatment. I think that the future will demonstrate that the essential chemical basis of the disease is a derangement of calcium metabolism.

In conclusion, I would emphasize that, from the practical standpoint, in rheumatoid arthritis (1) the etiological basis is a septic focus, and (2) the most important factors in treatment are the removal of the focus and the correct diagnosis of the stage of the disease.

REFERENCES.

¹ Billington: Auto-haemo-therapy, *Lancet*, March 1st, 1924. ² BRITISH MEDICAL JOURNAL, December 15th, 1923, p. 1157.

TREATMENT OF MIGRAINE.

BY

JANET LINDSAY GREIG, M.B., B.S. MELB.

RECENT authorities agree in stating that migraine is increasing in frequency and in accepting the nervous theory of its origin. The gastric theory that migraine is due to recurrent attacks of dilatation of the stomach has been ignored for many years, mainly because it does not seem to explain the many diverse symptoms.

It is possible that our knowledge of the stomach may not yet be complete. With the exception of the work done by Mr. Jefferson of Manchester, and reported at the Edinburgh meeting of the Anatomical Society of Great Britain, 1922, there has been no recent research work on stomach muscle. Its muscular mechanism is still more or less taken for granted.

What constitutes dilatation of the stomach and what is its clinical significance? We do not yet know the degree of dilatation which is the physiological normal of the stomach; it must vary with each individual. Radiography is only helpful when there are already gross pathological changes.

It is possible that in some individuals—the migrainous, for example—owing to muscular weakness, either inherited or acquired, the borderland between the two conditions may be a very narrow one, and the physiological may easily become the pathological. Accepting this as a possibility, the following scheme was drawn up for patients suffering from migraine, the endeavour being to maintain the stomach so far as possible in its normal physiological condition. The results have been most satisfactory. While patients are willing to submit to this routine they get no attacks of migraine and no sick headaches. The scheme is as follows:

I. Diet.—1. Exclude rigidly:

- (1) Cakes, scones, biscuits, pastry.
- (2) Steamed puddings, sponge puddings, trifles.
- (3) Green vegetables, including peas and beans.
- (4) Aerated drinks, including whisky and soda, and champagne; also fruit salts, sal hepatica, etc.

2. Allow any of the following:

Breakfast: Porridge, fish, eggs, bacon, chops, bread-and-butter or toast, marmalade, jam, tea or coffee.

Luncheon: Soup, fish, meat (of all kinds), potatoes, marrow, pumpkin, tomatoes, milk puddings, custards, junket, stewed fruits, baked apples, fresh fruit.

Afternoon Tea: Bread-and-butter, toast and butter, sandwiches, tea or coffee.

Dinner: Soup, fish, meat, poultry, potatoes, marrow, pumpkin, tomatoes, custards, jellies, fruit salads, ice-cream, fresh fruit, coffee.

II. *Medicine.*—During the first two weeks of treatment the following mixture is to be given three times a day, half an hour after meals, and then discontinued altogether:

R. Acid. hydrochlor. dil. m v to m x
Glycerin. m x
Aqua ad jss

As an aperient, cascara evacuant (P. D. and Co.) 1/2 to 1 drachm, water to 1/2 oz., should be given at night time when required, but not as a routine.

III. *Posture.*—The patient must sleep without a pillow, so that the head, neck, shoulders, trunk, and limbs are all on one level. Patients very quickly become accustomed to this position.

IV. *Exercise* in the open air and gradually increasing in amount. Walking is probably one of the best exercises and is available for all, whatever the condition or circumstances of the patient.

This method at least has this advantage, that it makes it possible to eliminate those cases of migraine which are due to dilatation of the stomach. It leaves, in my experience, very few cases for any other form of treatment. How long it will be necessary for those subject to migraine

to continue this routine I cannot say. While continuing the routine every effort should be made by means of tonics, good food, fresh air, but above all by exercise, to build the patient up physically.

It is probable that, with increased knowledge of the mechanism of the stomach muscle, we will understand more clearly the pathology of migraine, and understand better also how to maintain the stomach muscle in that stage of contraction which is its physiological normal.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

OPIUM POISONING AND PONTINE LESIONS.

THE record of Dr. Sydney Smith's interesting case of simulation of opium poisoning by an aneurysm of the basilar artery (BRITISH MEDICAL JOURNAL, June 7th, p. 994) may perhaps allow as a footnote a somewhat similar story.

An elderly woman, who lived alone, was found in bed and apparently unconscious by the charwoman who called in the morning by daily custom. The medical practitioner who was immediately summoned recognized a condition of profound coma with extremely contracted pupils, laboured respiration, and normal temperature. In addition, on a table at the side of the bed stood a phial of some two ounces capacity; it was empty and was labelled "laudanum." In these circumstances the practitioner not unnaturally decided to wash out the stomach, for the evidence, both clinical and circumstantial, clearly carried a suggestion of opium poisoning. The stomach contents, however, yielded no smell of laudanum, and later in the day an aggravation of the general condition and a temperature of 105° F. were judged to throw doubt on the original diagnosis. In the evening the patient died, and a *post-mortem* examination revealed a haemorrhage in the pons and fourth ventricle. Apart from the empty and labelled bottle there was no proof that the woman had taken laudanum, and certain other facts rendered this course unlikely.

Such an experience illustrates the ease with which a diagnosis may sometimes be handicapped, not helped, by external circumstances, and it emphasizes Dr. Smith's two points—namely: (1) that the clinical picture of pontine haemorrhage may closely imitate that of opium poisoning, and (2) that a high temperature as indicating a pontine lesion is of much value in the differential diagnosis. But, as the present record shows, and as seems also to have been the case in Dr. Smith's patient, the high temperature cannot be expected at the outset, and therefore there may well be a stage when a confident distinction between the two conditions may be difficult or even impossible. Add to this possibility an empty bottle labelled "laudanum" standing at the bedside, and most voices would surely be for the stomach-pump. Yet, as now appears, even here the majority might be wrong—perhaps not in practice but certainly in opinion.

London, W.

C. O. HAWTHORNE.

SURGICAL EMPHYSEMA DUE TO PARTURITION.

"SPONTANEOUS emphysema from rupture of a vomica, or even from rupture of the healthy lung in violent efforts, such as those of parturition, is a rare and curious affection." These words, from a footnote in the fourth edition of Holmes's *Surgery*, are my excuse for reporting the following case.

On May 20th I was called to attend a young and healthy primipara who was having frequent strong pains. By 10 a.m., the os being fully dilated but progress very slow, I decided to apply forceps. Chloroform was administered by my partner, and a large child delivered without undue difficulty. It weighed 9 lb. and was strong and healthy. At 7.30 p.m. the husband came and asked if I could give him a lotion as his wife's face and neck were swollen and painful. The following morning I found the neck and adjacent parts over the clavicles gave the characteristic crackling feeling of emphysema. By the end of a week the air had been absorbed with no bad symptoms. The patient could not remember feeling any pain in the chest during the labour.

As the quotation that heads this note is the only reference I can find in the literature at my disposal, I conclude that the condition is rare.

Goudhurst.

D'ARCY HARVEY, M.B.