and satisfactory. The thoroughness with which bone surgery is discussed will appear from the fact that at least nine methods are mentioned for the closure of cavities in bone.

The third volume deals with the surgery of the face, neck, breast, and chest, including the special surgery of the ear, nose, throat, and eye—an unusual combination for a book on operative surgery. The section on plastic surgery of the face is profusely illustrated, but we cannot find any allusion to the methods of Gillies and his co-workers in this country—a rather strange omission. In describing the operation for diverticulum of the oesophagus only the method of excision of the sac is discussed, and no mention is made of the operation in two stages or marsupialization of the sac which is popular with many surgeons. The chapters dealing with the surgery of the breast, thyroid, tongue, and ear are concise. Perhaps the best chapters are those on chest surgery. The operations for allowing the chest wall to fall in and of decortication of the visceral pleura are more clearly set forth than in most textbooks on the subject. In the treatment of unilateral pulmonary tuberculosis the author advises the subperiosteal resection of the second to eighth ribs inclusive from their angles to their cartilages. No mention is made of the recently discussed method of extrapleural thoracoplasty, where small sections of the posterior portions of the ribs, first to tenth or eleventh, are excised to achieve the same result.

The illustrations all through the volumes are not only numerous, but so well designed and arranged that in many instances the operation can be followed without reference to the text. The descriptions of the operations are written on a uniform plan. Dr. Bickham has collected and set out a great amount of information not easily to be obtained elsewhere.

NOTES ON BOOKS.

A *Pocket Oxford Dictionary of Current English*⁸ has been issued this summer by the Clarendon Press. It is the joint work of the late F. G. FOWLER and his brother, H. W. FOWLER, and is nominally an abridgement of their invaluable *Concise Oxford Dictionary*, first issued in 1911. Both works are therefore based on the great *New English Dictionary* (otherwise known as the *Oxford Dictionary*, or *N.E.D.*), of which the tenth and last volume is now appearing in sections. The present book, though it contains nearly as many pages, is narrower and less bulky than the *Concise Oxford Dictionary*, and in the ordinary paper edition can be carried easily in an overcoat pocket; the thinner India paper edition could no doubt be made to fit a coat pocket if the tailor were not looking. In a prefatory note the publishers express their appreciation of the late F. G. Fowler's services to literature and of his skill in planning this dictionary, which they have good reason to describe as a triumph of condensation. The authors' preface mentions the difficulty of squeezing their earlier work, and indicates the changes in method that have been adopted apart from mere reduction of quantity. In the first place, a pronunciation system has been devised in order to make clear how the compilers recommend that every word should be pronounced; secondly, the alphabetical arrangement of phrases within all long articles has been carried as far as possible; thirdly, etymologies have been cut down to little more than bare statements of ultimate origin; and, fourthly, a large number of words and senses not in the former work (some of them by-products of the war) have been included. Whether carried about in the literary workman's pocket or kept in a handy place on the writing table, this admirable little reference book should prove a boon to countless users of the English language.

The second or Summer Number of the sixth volume of the *Annals of Medical History*⁹ has Sir William Bowman's portrait on the cover; his life is sketched by Dr. Burton Chance, and is the first of his three biographies of the famous ophthalmologists of the nineteenth century, the other two being Donders and von Graefe. This article is generously illustrated by four full-page plates of microscopic drawings from Bowman's work, *The Physiological Anatomy of Man*. The frontispiece of this instalment represents Carl Weigert, Paul Ehrlich's cousin, the author of a monograph on the

lesions of small-pox, which he wrote out six times; in it he first described coagulation-necrosis, and in his "Schlwa" theory, so called after the Hindu god of destruction, opposed Virchow's view that all inflammatory growth is due to the stimulus of external irritants. Dr. Burget writes pleasantly of Lazzaro Spallanzani and his important physiological discoveries; while visiting Bonnet at Geneva Spallanzani met Jean Senebier, of whom Professor C. Greene Cumston elsewhere in this instalment provides a note and a portrait. In his account of Sir William Buttes of Norfolk, physician to Henry VIII, Dr. T. N. Toomey has made considerable additions to our knowledge of the only English physician mentioned by Shakespeare, for Dr. Toomey considers that the John Kaye (Calus) in *The Merry Wives of Windsor* was not an historical person or the real Calus. The editor, Dr. F. R. PACKARD, collects the references to syphilis in Shakespeare's plays, and Dr. M. Charlton of Montreal provides the third part of his outlines of medicine in Lower Canada. James Atkinson of York, who wrote a medical bibliography as far as the end of the letter B, is described by Dr. John Rahrh of Baltimore as an unconscious imitator of Sterne, and as the author of a work which should, after some judicious trimmings, repay republication. Dr. Henry J. John's translation of notes on the surgical clinic of the Charles University, Prague, Bohemia, a hundred years ago, makes the reader feel thankful that he was not one of the patients.

MISS ELIZABETH S. HALDANE's work on *The British Nurse in Peace and War*¹⁰ is divided into two parts. The first, which consists of five chapters, contains an account of nursing from the earliest times, including the history of St. Bartholomew's Hospital, the London Hospital, and the Edinburgh Royal Infirmary, the teaching work of Miss Nightingale, and the evolution of modern nursing. It is said that the darkest period of nursing was from the later part of the seventeenth century to near the middle of the nineteenth century, when, owing to the influence of Miss Nightingale, a great and momentous change took place. A special chapter devoted to modern nursing contains a short account of the nursing profession in the United States and Canada, and of special branches of nursing, such as district nursing, Poor Law work, public health nursing, private nursing, maternity work, etc. The second part deals with the organization of the nursing profession in the great war both at home and abroad.

¹⁰ *The British Nurse in Peace and War*. By Elizabeth S. Haldane, C.H., LL.D. London: John Murray. 1923. (Cr. 8vo, pp. x + 232. 7s. 6d. net.)

A TABLE OF STANDARD DIETS FOR USE IN DIABETES.

BY

R. PYBUS, AND D. MURRAY LYON, M.D.,

SISTER DIETITIAN, ROYAL CHRISTISON PROFESSOR OF THERAPEUTICS,
INFIRMARY, EDINBURGH, UNIVERSITY OF EDINBURGH.

A LARGE number of papers have appeared recently dealing with the insulin treatment of diabetes. All the writers agree that the use of insulin, far from getting rid of the need to regulate the diet, has made a careful control of the food intake more important than ever. The reason for this is that the employment of insulin has introduced the new potential danger of hypoglycaemia from overdosage. Besides this, there seems to be no doubt that the patient stands the best chance of improving his condition if the insulin and the diet are so balanced that the blood sugar level is kept approximately normal. This can only be managed by strict control.

Vast improvements have recently been made in our knowledge of how to diet a diabetic most efficiently. The object to be attained in treating a patient is to give a diet which will yield sufficient calories, will avoid acidosis and glycosuria (or rather hyperglycaemia), and will at the same time rest the pancreas and metabolism generally. Certain basic principles underlie the construction of such a diet.

Even while a patient is resting quietly in bed all day he must burn up a certain number of calories in order to maintain life, and every activity, muscular or mental, must be paid for in additional calories. The basal metabolic weight of a person can be predicted from his size and weight by reference to such tables as those of Du Bois. In addition to the allowance to be made for exercise, etc., a further surplus must be provided to allow for storage in the case of greatly emaciated patients. For these reasons the total

⁸ *The Pocket Oxford Dictionary of Current English*. Compiled by F. G. Fowler and H. W. Fowler. 1924. Oxford: The Clarendon Press. (3½ x 6½; pp. vi + 1,000. 3s. 6d. net; India paper 6s. net.)

⁹ *Annals of Medical History* (Summer Number, 1924), vol. vi, No. 2. Edited by Francis R. Packard, M.D. New York: Paul R. Hoeber Inc.; London: Baillière, Tinsall and Cox. (8½ x 12½; pp. 143-244; illustrated. Subscription in Great Britain, 22 2s. per volume.)

number of calories required can only be guessed at approximately, and no useful purpose is to be served by employing more detailed formulas or tables in calculating the total daily rations.

There are, however, certain other principles which can be applied with advantage in the construction of the diet. The first limit imposed is the patient's difficulty in dealing with more than a small quantity of carbohydrate. Unrestricted use of protein is also to be deprecated since protein tends to hasten the metabolic rate, and this is to be avoided. The minimum supply of protein necessary to make good nitrogen wastage is variously estimated at from 0.5 to 1.5 grams per kilo body weight. This is really a fairly wide difference. Taking the weight of a boy 5 ft. high and 15 years old as 50 kilos, and that of a man 55 years old and 6 ft. high as 75 kilos, the lower values would be 25 to 37.5 grams of protein and the higher estimates 75 to 112.5. The figures at best are therefore arbitrary, and are so divergent that we feel justified in choosing a medium range for the protein and neglecting individual calculations for each patient.

Since fat at 9 calories to the gram is a much more valuable food than protein or carbohydrate at 4 calories per gram, it is desirable for the diabetic to take as much of it as possible. This allows him a diet of higher calorific value and makes up for the deficiency of carbohydrate. But in the case of fat there is also a limit to the amount which can be taken. It is known that in order to permit complete combustion of a unit of fat a certain quantity of carbohydrate must be burnt up at the same time. When an excess of fat is broken up, some is only partially combusted, and in consequence dangerous ketone bodies appear in the blood and urine and give rise to acidosis. The correct ratio of carbohydrate to fat is shown by the formulas of Woodyatt. In calculating the glucose and fatty substances in the diet, it must not be forgotten that under certain conditions some of the protein, instead of being built up into the tissues, may become transformed into either fat or carbohydrate. Woodyatt has shown that in estimating the total available glucose ("G.") there must be included 100 per cent. of the carbohydrate, 58 per cent. of the protein, and 10 per cent. of the fat (glycerin fraction), while in addition to the remaining 90 per cent. of the fat, 46 per cent. of the protein may yield fatty substances ("F.A."). Every gram of "G." in the diet will allow of the complete combustion of $1\frac{1}{2}$ grams of "F.A." The maximum amount of fat that can be included in a diet without the appearance of acid bodies is shown approximately by the formula $F.A. = 2C. + \frac{1}{2}P$. Practical experience of the use of these formulas has taught us their immense value.

If consideration has to be given to all these points in the construction of a diet the process becomes complicated enough, and when new diets have to be drawn up daily during treatment the process is so time-consuming as to be impracticable for the busy man. The difficulty can be got over in hospitals where large numbers of diabetics are under treatment at the same time by delegating the duties to a competent dietitian or by the use of standard diet tables.

Appended are the tables which we employ. A great deal has been written about the necessity for individualization of the patient, and with that we agree. But the use of such a table as that given does not make this impossible. The daily summaries serve as a basis, and modifications can readily be made as required to suit the patient. The diets are not really so monotonous as they appear, since the range of vegetables is considerable, the heading "meat" includes chicken, rabbit, etc., which are approximately the same value as beef, and the eggs, etc., can be served in various ways. The real limit to variety is the relatively small quantity of protein present.

The advantages of a graduated series of standard diets seem obvious enough. They benefit both the patient and his attendants. The tables are easily used, and their employment greatly simplifies the dieting of diabetics. The diets have been drawn up so as to yield a certain number of calories. A step of 100 calories between each summary allows of a sufficiently gradual advance. Since the G.:F.A. ratio has been carefully balanced, the amount

of the protein and the quantity of carbohydrate also increase very gradually. The diets are divided into four classes, constructed for different purposes.

Group "A" diets are for cases who show clinical signs of acidosis, dyspnoea, acetone in the breath, etc. They contain a relatively large proportion of readily available carbohydrate, and have a G.:F.A. ratio of about 2:1. This excess of glucose is sufficient to aid the complete combustion of a considerable amount of fat. Another result of the use of this series is that glycosuria may be increased, but removal of the ketones is so important that the glycosuria can be neglected for the time being. Insulin should always be employed with the "A" diets.

The remaining diet summaries—B, C, and D—can equally well be employed with or without insulin, and have also been used for mild or ambulant cases.

Group "B" diets are intended to "desugarize" the patient. The G.:F.A. ratio in this series is 1:1, which still leaves a considerable margin for dealing with extra fat. This is necessary because the diets in this group are below the amount required to maintain the basal metabolism and the body stores of fat or protein have to be called upon to make up the deficiency. All cases, if free from clinical acidosis when first seen, are put on B 15 at once. Mild cases may become sugar-free on three or four days of B 15 or after a further three days on B 12. Lower diets are only given to such ambulant cases as are in good physical condition and are practically free from acetone. More severe cases are usually put to bed, while the diets are dropped below basal levels. The rate of lowering the diets varies. Sometimes the patient remains a day or two on each level, often he drops rapidly from B 15 to B 7 and B 3. Usually the urine becomes sugar-free without the subject actually requiring to starve. When this stage is reached the patient is transferred to the dietary ladder (Group "C" diets), starting with a diet having about the same G. value as the last "B" diet or slightly less.

Group "C" diets form an increasing series which are used to test the limit of the patient's tolerance and which serve for a time as low maintenance diets. When time permits the increase in diets should be very slow. So long as the urine remains sugar-free the patient advances each day or each second day in most severe cases. Should sugar reappear he drops for one day to a diet of about half the value, then comes back to the highest diet on which he was sugar-free, and thereafter attempts to advance again. Often this is successful, but eventually a point is reached which the patient cannot pass. This is the limit of his tolerance at the moment. Where it is at all feasible the patient should advance to this point without the use of insulin, so that his unaided capacity may be gauged. Insulin need only be given in those cases where the limit is too restricted to permit of a useful life being led; it is not required when the patient's tolerance is fairly good. Ten units of insulin should allow the addition of about 200 or 300 calories. The patient may be kept on a diet of this class for some months or indefinitely. The small amount of protein, however, restricts the choice of foodstuffs, and for those patients who wish a change or who require more protein the Group "D" or maintenance diets are provided.

A weekly low day on which about half rations are consumed is essential for cases who take no insulin, and even where insulin is being administered it is well to observe the same custom. It appears to be quite satisfactory to control the progress of the case by examining a sample of the collected urine for the twenty-four hours, blood sugar sampling being unnecessary.

The figures given in each horizontal line of Table I represent the daily total, which has to be divided into suitable meals. This offers little difficulty, and often can be left to the patient. When no insulin is being given the food is divided up as equally as possible. If insulin is being used the carbohydrate should be so distributed as to be "covered" by the drug.

It will be noticed that the greatest amount of carbohydrate is in the form of unsweetened oatcake or porridge. This is because we believe that the diabetic can tolerate oatmeal better than other forms of starchy food, owing to its being more slowly absorbed. An ounce of dry oatmeal

TABLE I.—STANDARD DIETS.

DIETS.	Milk.	20% Cream.	5% Vegetable.	Eggs.	Lean Meat.	Clear Soup.	Bacon.	Butter.	Oatmeal.	Dry Meal (as porridge).	White Bread.	Potato.	Cheese.	Orange Pulp.	Olive Oil.	White Fish.	GRAMS.						TOTAL CALORIES.
																	Carbo- hydrates.	Protein.	Fat.	G.	Ratio.	F.A.	
GROUP A.																							
A 24	20			2	3	16		1		3	6½	4		16½		4	271.5	127.8	91.2	354	2.5 : 1	142	2419
A 20	20			2	2	16		2		2	6½	4		8½		2½	226	106	76	294	2.5 : 1	117	2014
A 17	20		6½	2	3	10		2		1½	5	3					164.25	89	76	224	2 : 1	109	1700
GROUP B.																							
B 18	5½		11½	2	2½	8	2	2½	3	1½							94.25	76.3	130.5	152	1 : 1	152	1856
B 15	2		16½	2	2		1½	2½	4								78	60	106.5	124	1 : 1	124	1510
B 12	3		16½	2	1		1	2	3								64.5	46.75	89.5	102	1 : 1	102	1244
B 9	2		15	2	1		1	1	2								46.5	41.25	62	76	1 : 1	76	909
B 7	2	1	11½	1	1		1	2	1½								37	32.85	52.85	61	1 : 1	63	755
B 5		2½	13½	1				½	½							2	22	28.4	34.25	43	1 : 1	44	510
B 3	2	2	11½	1													15	15	20	26	1 : 1	25	300
Starve day.*	2																3	2	2				38

* Tea, coffee, and bovril *ad lib.*

If not sugar-free give coffee only every two hours till urine has been free of sugar for 24 hours.

GROUP C.																	7.5	3.75		9	1 : 0.2	2	45
Starve day.*			8½														15	7.5		19	1 : 0.2	3	90
C 1			16½														15	15	20	26	1 : 1	25	300
C 3	2	2	11	1													19	29.5	34	39	1 : 1.1	44	500
C 5	2	1	16½	1	1	8		½									25	37.75	50.5	51	1 : 1.2	62	705
C 7	2	2	16½	2	1	8		2	½								31.5	39.75	72	62	1 : 1.3	83	900
C 9	2	2	18½	2	1	8		1½	2								37	42.25	88.5	70	1 : 1.4	98	1114
C 11	2	4	16½	2	1		1	1½	1								47.5	45.6	104	85	1 : 1.4	115	1308
C 13	2	4	20	2	1		1	1½	1½								50	50.25	122.5	91	1 : 1.5	133	1500
C 15	2	4	20	2	1	7	1	2½	1½								55	57.5	139.5	101	1 : 1.5	151	1701
C 17	2	4	20	3	1	7	1	2½	2								62.75	60.5	157.25	113	1 : 1.5	169	1908
C 19	2	4	20	3	1	7	1	3½	2	½							68	60.6	166.25	119	1 : 1.5	177	2010
C 20	2	4	20	3	1	7	1	3½	1½	1							70.5	64	172.5	124	1 : 1.5	185	2100
C 21	2	3	20	3	1½	7	1	4½	1½	1		1					73.5	68.5	181.5	130	1 : 1.5	195	2201
C 22	2	3	20	3	2	7	1	4	1½	1		1½			1		77.6	69.4	190.5	136	1 : 1.5	203	2302
C 23	2	3	19	3	2	7	1	4	1½	1½		1½			1½		80.75	69.5	200	141	1 : 1.5	211	2401
C 24	2	3	16½	3	2	7	1	4	1½	1½		1½			2		85.75	70.75	210.5	148	1 : 1.5	222	2520
C 25	2	3	16½	3	2	7	1	4	1½	1½		1½			2½		91.25	72	220.75	155	1 : 1.5	231	2639
C 26	2	3	16½	3	2	7	1	4	2	1½		2			2½		98.75	74	238	165	1 : 1.5	248	2837
C 28	2	3	16½	3	2	7	1	4	2½	1½		2			4½								

* Tea, coffee, and bovril *ad lib.*

High Maintenance																							
GROUP D.																							
D 20	4	4	15	3	2	8	2	2½	2½				1				61	80.75	164.25	124	1 : 1.5	185	2045
D 21	1	4	15	3	2½	8	2	2½	2½			1	1				62.5	82.75	170	127	1 : 1.5	191	2112
D 22	1	4	15	3	3	8	2	3	2½			1½	1				65.5	87.25	177.75	134	1 : 1.5	200	2211
D 23	4	4½	15	3	3	8	2	3	2½			1½	1				70.5	90.75	187.25	142	1 : 1.5	210	2330
D 24	4	4½	15	3	3	8	2	3½	2½			2	1				73.5	91.25	195.25	146	1 : 1.5	218	2416
D 25	4	4½	15	3	3	8	2	3½	2½	½		2	1				79	92.85	204	153	1 : 1.5	225	2513
D 26	4	4½	15	3	3	8	2½	3½	2½	½		2	1				81.5	96	211.5	158	1 : 1.5	234	2613
D 27	4	4½	15	3	3	8	2½	3½	2½	¾		2	1½				84	99.75	211	163	1 : 1.5	245	2723
D 28	4	4½	13	2	2½	8	2½	4	2½	1		2	1½		1	2	87.5	103	228	170	1 : 1.5	252	2816
D 29	4	4½	13	2	2½	8	2½	4	2½	1½		2	1½		1½	2	92.5	104	236.5	176	1 : 1.5	260	2912
D 30	4	4½	13	2	2½	8	2½	4	2	1½		2	1½		2	2	95.25	105	247	180	1 : 1.5	270	3020
D 31	4	4½	13	2	2½	8	2½	4	2½	1½		2	1½		2½	2	99	106	256	185	1 : 1.5	279	3124
D 32	4	4½	13	2	2½	8	2½	4	2½	1½		2	1½		2½	2	102.75	107	262	191	1 : 1.5	285	3203
D 33	4	4½	13	2	2½	8	2½	4	2½	1½		2	1½		3½	2½	105.25	108	271	195	1 : 1.5	295	3301

N.B.—The measure for olive oil is a tablespoonful. The other figures indicate ounces except in the case of eggs. Oatmeal is shown in ounces (dry weight) before being made into porridge.

makes about four tablespoonfuls of porridge. Oatcake and porridge are interchangeable in the proportion of 1 oz. oatcake=3 tablespoonfuls porridge. White bread has a calorific value very similar to that of oatcake, and a small quantity of it may replace the oatcake from time to time. We find, however, that large amounts of oatcake can be tolerated by patients who at once show sugar when a small quantity of bread is taken.

One of the difficulties in following out the régime of balanced dieting is the presence of large quantities of fat in the higher diets. Four ounces of butter is difficult to dispose of unless sufficient solid food is provided to carry it. It is therefore convenient to serve most of the starchy food as oatcake rather than porridge. If salad oil cannot be taken butter should replace it, and the excess of this can be served with the eggs or as sauce with the meat, fish, or potato. Tea, coffee, and bovril have no calorific value, and can be taken within reason with all these diets. They are required in considerable amount on those days when the diet is extremely low.

Vegetables containing 5 per cent. carbohydrate include: Lettuce, cucumber, spinach, asparagus, rhubarb, endive, marrow, celery, mushroom, tomato, brussels sprouts, watercress, sea-kale, cauliflower, cabbage, radishes, leeks, broccoli.

Vegetables containing 10 per cent. carbohydrate include: French beans (young), turnip, carrots, onions, green peas (canned).

After the patient has been for some time on a maintenance diet he may be taught to vary it by substituting articles selected from a suitable list, or may even build up a complete menu to a prescription of so many grams of carbohydrate, protein, and fat yielding the desired number of calories. Food values for this purpose are to be found in the food tables of Harrison and Lawrence, and many other publications.

TABLE II.—Quantity of Carbohydrate, Protein, and Fat in 30 grams, or cubic centimetres, or in 1 oz.

	Carbo- hydrates.	Protein.	Fat.	Calories.
Milk	1.5	1	1	19
Skim milk	1.5	1	0.09	11
Buttermilk	1.5	1	0.15	11
Whey	1.5	0.25	0.09	8
20 per cent. cream	1	1	6	62
5 per cent. vegetable	1	0.5	0	6
10 per cent. vegetable	2	0.5	0	10
Eggs, each	0	6	6	78
Lean meat	0	8	5	77
Chicken	0	8	5	77
Boiled chicken without gravy	0	8	3	59
Bacon	0	5	15	155
Lean ham	0	6	6.5	83
Butter	0	0	25	225
Oatmeal	20	5	2	118
Oatcake	15	3.75	1.5	89
Porridge	10	2.5	1	60
White bread	16	3	0.5	81
Potato	6	1	0	28
Orange pulp	3	0	0	12
Cheese (average)	0	8	11	131
Olive oil (two tablespoonfuls)	0	0	30	270
Boiled fish	0	6	0	24
Clear soup	0	0.6	0.25	5

Summary.

It is pointed out that the directions usually given for drawing up individual diets are too complicated to be universally serviceable. A carefully balanced dietary scale is submitted to serve as a basis for treatment. The methods to be followed to clear up acidosis, to get rid of glycosuria, and to test the patient's carbohydrate food tolerance are also described.

THE STATE OF THE PUBLIC HEALTH.

SIR GEORGE NEWMAN'S REPORT FOR 1923.

[FIRST NOTICE.]

THE annual report of the Chief Medical Officer of the Ministry of Health for 1923¹ is the fifth made under the Ministry, but is the sixty-sixth of the series begun by Sir John Simon under the Privy Council and carried on by his successors under the Local Government Board.

Sir George Newman begins by pointing out that as there is no system of registration of sickness the sources of information available in considering the state of the public health are the returns made by the Registrar-General relating to births, deaths, and infectious disease, and the data and records in respect of invalidity under the National Health Insurance Acts. The information so available is, however, supplemented by annual and special reports of medical officers of health, by the results of investigations by medical officers of the Ministry and by other Government departments interested in the physical, industrial, and social welfare of the people.

AGE AND SEX DISTRIBUTION OF THE POPULATION.

The following table is given showing the age distribution of the population of England and Wales according to the censuses of 1841, 1911, and 1921.

Age.	1841.			1911.			1921.		
	M.	F.	Persons.	M.	F.	Persons.	M.	F.	Persons.
0-5	6,587	6,649	13,236	5,368	5,319	10,687	4,438	4,329	8,767
5-10	5,989	5,981	11,970	5,121	5,128	10,249	4,663	4,625	9,288
10-15	5,533	5,352	10,885	4,845	4,858	9,703	4,849	4,811	9,660
15-20	4,912	5,060	9,972	4,588	4,662	9,250	4,561	4,686	9,247
20-25	4,546	5,196	9,742	4,166	4,638	8,804	3,823	4,495	8,318
25-35	7,388	8,006	15,394	7,850	8,662	16,512	6,919	8,288	15,207
35-45	5,471	5,667	11,138	6,477	6,957	13,434	6,589	7,522	14,111
45-55	3,900	4,099	7,999	4,698	5,084	9,782	5,630	6,036	11,666
55-65	2,506	2,720	5,226	3,009	3,364	6,373	3,650	4,038	7,688
65-75	1,412	1,628	3,040	1,671	2,100	3,771	1,926	2,410	4,335
75-85	545	651	1,196	509	749	1,258	595	916	1,511
85-	83	119	202	63	114	177	66	135	201
	48,872	51,128	100,000	48,365	51,635	100,000	47,709	52,291	100,000

M. = Males, F. = Females.

Upon this table Sir George Newman makes the following observations:

"It will be seen that the indirect consequence of the war, in accelerating the decline of the birth rate, has been much greater than the direct effect, the destruction of young male adults. This latter consequence is brought about in (a) the decreased proportion of males of 15-35, who formed 16.8 per cent. of the population in 1841 and almost as many (16.6 per cent.) in 1911, but only 15.3 per cent. in 1921; (b) the increased proportion of females to males, 51.1 per cent. of the population in 1841, 51.6 per cent. in 1911, and 52.3 per cent. in 1921. But owing to the declining birth rate, a decline temporarily accelerated by the war, the change in the proportion of children has been much greater. In 1841, children under 5 constituted 13.2 per cent. of the population; by 1911—after rather more than thirty years of a declining birth rate—the proportion had declined to 10.7 per cent., a decrement of 2.5 per cent. In 1921 only 8.8 per cent. of the population were under 5 years of age; the decrease in these ten years was more than 78 per cent. of the whole decrease over the previous seventy years and two and a half times as great as between 1901 and 1911. The proportion of young children was only 66 per cent. of what it had been eighty years ago. On the other hand, the proportion of persons over 65 was 36 per cent. greater than in 1841. In 1921, for the first time, the first quinquennium of life did not contain a larger number of living persons than any other quinquennial group of ages. Even

¹ *The State of the Public Health*. Annual Report of the Chief Medical Officer of the Ministry of Health for the year 1923. H.M. Stationery Office. (Med. 8vo, pp. 212. Price 3s. net.)