It must always be remembered, however, that manipulations of any kind can introduce virulent organisms into the uterus. It is impossible to sterilize absolutely a vagina, however thoroughly it may be attempted, and many virulent organisms can and do exist and multiply there with no harmful results. But if those same organisms gain access to the inside of the uterus they may do untold damage.

Two of my cases were of that most fatal kind, the acute aplastic type of streptococcal infection. Both had been infected with gonorrhoea, and one had subacute gonorrhoea at the time of delivery. She was also hypermature, had a flat pelvis, and required craniotomy into the bargain.

SUMMARY.

1. When fever appears in the puerperium, having excluded maladies outside the pelvis which may cause it, it is far safer to give an anaesthetic and carefully examine the patient's pelvis than to "wait and see." If intra-uterine sepsis is present, explore the uterus and take steps to empty it or clean it, or both. To wait and see if the fever will subside may mean waste of precious time.

2. A so-called sapraemia is quite often the precursor of the most virulent septicaemia, and anyhow a sapraemia will not be improved by leaving even small and removable fragments of placenta or membrane *in situ*, if they are poisoning the patient.

3. The septic uterus will not involute, but the modified Carrel-Dakin treatment, assisted perhaps by drugs, will help involution better (in my opinion) than anything we have at our disposal.

4. If antistreptococcal serum is to be used, use it frequently until the temperature falls.

5. Further adjuncts are "peptone shock" and intravenous eusol.

6. Take a culture from the endometrium and from the blood, as a vaccine later may be of value, especially if prepared from the blood.

7. Pay careful attention to posture in the nursing of these patients. Fowler's position will aid drainage, whereas the supine position will tend to dam back secretions. Similarly micturition and defaecation in a complete sitting position are useful aids.

8. In early infection of the adnexa, the modified Carrel-Dakin method may possibly prevent the necessity of abdominal section, and will certainly speed up recovery.

9. Where anaemia or debility is caused by haemorrhage or toxæmia keep a special look-out for sepsis.

10. Perform all manipulations or instrumentation with extreme gentleness whenever possible, and practise asepsis most punctiliously.

11. Sew up all tears of soft parts immediately.

Puerperal sepsis is a type of wound infection which will not improve by watching and waiting. Operative measures and certain forms of medication have scope, though still limited, in its treatment. There is no panacea, though we hope it may appear. Until it does, each case will require careful study with a view not only to its own cure, but also to the discovery of a real antidote to this scourge.

OCCLUSION OF THE OS FOLLOWING SURGICAL TREATMENT OF UTERINE AND VAGINAL PROLAPSE.

BY

J. NIGEL STARK, M.D., F.R.F.P.S.,

SURGEON, ROYAL SAMARITAN HOSPITAL FOR WOMEN, GLASGOW.

MANY women are compelled to seek relief for the disability and discomfort arising from varying degrees of prolapse of the vaginal walls and uterus. A pessary is the usual means of initial treatment, but often, after a while, it is found necessary to discard it because either it fails to afford sufficient support, or, if it is composed of vulcanite, its pressure upon the soft parts causes pain and irritation. Not infrequently the patient herself becomes impatient and demands more radical treatment, having learned from experience that a pessary, especially a rubber ring, demands

personal attention which becomes irksome as well as several visits annually to or from her medical adviser.

In the event of any of these circumstances occurring surgical treatment can with confidence be advised in the majority of cases. It is now almost universally realized that plastic operations upon the perineum, the vaginal walls, and the cervix of the uterus are sufficient in the great majority of women to afford relief and ensure comfort. The strenuous advocacy for many years of Fothergill and others has resulted in their theories having been accepted and their method of treatment adopted by the great majority of gynaecological surgeons.

As a routine step in the operations, amputation of the cervix is nearly always performed, as it is followed by great benefit in more ways than one. The cervix in conditions of bad prolapse is so often extensively lacerated, eroded, and ulcerated that it is a constant standing source of disease, and, at middle life, a menace as a likely seat of cancer. The purulent discharge which has its origin in such a cervix is a fruitful source of local irritation and discomfort, while the absorption of the toxins is responsible, in my opinion, for much ill health, the prime cause of which is frequently unsuspected. Removal brings about cessation of the discharge, and, in addition, tends to relieve the congestion and to lessen the weight of the generally enlarged and heavy uterus.

Caution, however, must be exercised in the performance of this simple operation, should the patient be of parous age, or her last condition may be worse than her first. I have seen several examples of repeated abortions subsequent to operation, caused without doubt by too high amputation of the cervix, and the following history of the yet more serious consequences which may result from an operation for prolapse in a woman before her menopause seems to be worth recording as a reminder that zeal must always be tempered with discretion.

A multipara, aged 42, who had suffered for several years from a moderate but gradually increasing degree of prolapse, chiefly of the vaginal walls, consulted a well known and esteemed gynaecologist during the absence from home of her usual medical attendant. She was ordered into a nursing home, where, apparently, the operations of anterior and posterior colporrhaphy, amputation of the cervix, and perineorrhaphy were performed, and in two to three weeks she was again at home.

Shortly afterwards attacks of abdominal pain began, at first mild and soon passing away, but gradually becoming more severe until at last they were agonizing enough to call for a hypodermic injection of morphine. Between the attacks the patient was fairly well and was able to go about, but, six months after the operation, she was seized with an attack of such unusual severity that it was considered necessary to have a consultation.

The foregoing history was told to me, and I learned further that menstruation had not appeared since the operation, although it had previously been regular. During the attacks there had never been a rise of temperature. The patient was well coloured and stout. On making a vaginal examination it was found that the operations had successfully effected the formation of a restored perineum and of a normal vaginal canal. The vaginal vault was devoid of any trace of a cervix, its place being taken by a firm cicatrix. Abdominal examination revealed the presence of a tense pyriform swelling, evidently the body of the uterus, rising mesially almost to the umbilical level. Upon questioning the patient the fact was elicited that the pain from which she suffered came spasmodically and reminded her of labour pains.

The diagnosis was apparent, and a day or two later I performed hysterectomy and double salpingectomy. The walls of the uterus were greatly thickened and its cavity contained large quantities of thick blood of a dark brown colour. No trace of the internal os could be found—the cavity was entirely shut off; stenosis was complete. The Fallopian tubes were greatly distended by the same thick chocolate-coloured fluid as was in the interior of the uterus.

No other treatment could have been attempted with any hope of success. Obviously, serious obstacles and dangers which readily occur to one's mind and need not be mentioned would have militated against a good result from any attempt to open up the uterus from below. The patient having children and being 42 years of age, there was all the less objection to hysterectomy being performed. She has remained in good health and has had no pain eight months after the operation.

A too thoroughly radical removal of the cervix in a woman of child-bearing age may therefore be, as this case proves, a dangerous proceeding in the surgical treatment of prolapse, leading to stenosis which may occlude completely the canal at the internal os and so cause retention of menstrual blood.