sound-conducting apparatus, enable sonorous vibrations more easily to reach the defective, but not impotent, auditory nerve.

Our next step is ocular inspection of the drum-membrane. It is only in rare cases that changes in the substance of the membrane, sufficient to account for marked impairment in hearing, are met with. Sometimes, it is true, we can see it to be thickened and altered that good hearing-power is out of the question. Again, it may be atrophied and flaccid, moving with every respiration, or perfectly transparent. Patches of opacity may exist as the remnants of an acute process in years gone by, which has left perfect hearing-power. The irregularity of the malleus, described by von Troeltsch as common in dry catarrh, is by no means always absent in health. The bright spot may be altered in a variety of ways without causing any symptoms, and I believe that its significance is much overrated. There is, however, an appearance which is often met with, and which, I do not doubt, is familiar to all present, although I have not seen it exactly described in works on otology. I allude to a streaky—one might almost say stringy—opacity in a membrane little altered in transparency except where these opaque fibrous-looking lines occur. This condition is common in those insidious cases in which the hearing goes from bad to worse, or at best remains bad.

What, then, are we to consider as distinguishable conditions of the membrane which render the prognosis favourable? So far as my little experience goes, I should say that in cases in which the drumhead is distinctly indrawn, but not fixed in this position by adhesions, we may have fair hopes of effecting improvement, although, of course, a positive prognosis can only be given after using further methods of examination. Again, the appearance of fluid behind the drum-membrane renders the prognosis very hopeful; for exudation can be removed, and its very presence shows that the mucous membrane lining the cavity has not altogether passed from the stage of cellular infiltration to interstitial thickening.

To sum up; the knowledge we gain by inspection of the drum-membrane is in a large proportion of cases only to be utilised in conjunction with other methods of examination. Thus, if it be opaque in certain parts from interstitial thickening, and if there be deafness, we assume that the opacity is a visible sign of thickening in other parts essential for the act of hearing; while, if it be opaque without the presence of deafness, we make no such assumption.

Turning next to the examination of the naso-pharynx, I need hardly allude to adenoid vegetations and nasal polypi, both of which conditions are often causes of curable deafness. Chronic naso-pharyngeal inflammation, however, requires some further remarks. It is universally admitted that, in a large proportion of cases, tympanic disease is the result of extension of a catarrhal process along the Eustachian tubes. In such cases we may, I believe—other things being equal—consider that the condition of the pharyngeal mucous membrane may be taken as approximately representing that of the tympanum. If, then, that be a pharyngitis characterised by unorganised moist swelling and excessive secretion, we may, *a priori*, assume a corresponding condition of the middle ear as part factor of deafness; while, if we find an inflammation of the naso-pharynx which has gone on to the deposit of fibrous tissue, then we are justified in suspecting the existence of thickening or adhesions within the tympanic cavity. The absence of all sign of pharyngeal mischief in a case of chronic non-suppurative middle ear-disease is also an unfavourable prognostic sign.

I shall next call your attention to what I believe to be the most important aid, alike to diagnosis, prognosis, and treatment. I allude, of course, to the air-douche. Auscultation during this process will enable us to decide whether the air passes through a wide or narrow tube, and also whether the latter contains any free exudation. The absence of secretion, as denoted by a dry sound, naturally leads one to fear that the pathological condition has passed from the stage of swelling and excessive secretion to that characterised by the deposit of fibrous tissue. Inspection of the membrane while compressed air is driven through the Eustachian tubes may also furnish valuable information as to the presence of adhesions. The same knowledge may, of course, also be gained by using the pneumatic speculum.

It is of great importance to note accurately the effect of the air-douche upon the hearing-power of the patient. The factors of deafness which are liable to be influenced by its agency are:

1. Obstruction of the Eustachian tube; so much of the deafness as is due to this cause alone being at once completely, though temporarily, banished.

2. Excessive secretion, causing the presence of exudation within

the tympanum. The action of the air-douche will probably scatter the secretion, thus relieving the clogging of the ossicles and membrane. We must remember, however, that this process of excessive secretion must necessarily—except, perhaps, in cases of hydrops *ex vacuo*—be associated with swelling of the mucous membrane, due to hyperæmia and infiltration of lymphoid cells—a condition amenable to treatment.

3. Adhesions delicate enough to be influenced by the current of air, and so situated as to be stretched or ruptured by its action.

4. Perhaps I should here mention another possible factor of deafness, which is only, however, a secondary result of disease or abuse of treatment, namely, a flaccid condition of the drum-membrane. In such cases, temporary improvement—which may sometimes, by suitable treatment, be maintained—is produced by the use of the air-bag.

If inflation of the middle ear produce improvement, however temporary, we are entitled to hold out hope to our patient, because we have a curable lesion to deal with—always remembering that the prospect of benefit is directly proportionate to the duration of such temporary improvement. If, on the other hand, there be, after several injections of air, no change, then we must suspect the existence of an incurable factor of deafness, viz., the presence of adhesions or thickening in situations where they cannot be acted on by the therapeutic means at present recognised as safe and efficient. In arriving at this conclusion, however, we must take into consideration the various other points to which I have alluded, more especially the condition of the naso-pharyngeal mucous membrane, and the phenomena yielded by the Eustachian catheter and auscultating tube. In calling this factor of deafness incurable, I am aware that the way to criticism is opened, for cases of interstitial thickening are just those for which fluid-injections *per tubam* are advocated. While not prepared to deny altogether the efficacy of this plan of treatment, I cannot help thinking that it has been much overrated. How we can hope to cause disappearance of organised tissue by such highly diluted solutions as alone it is safe to inject into the middle ear, I fail to understand. A solution of iodide of potassium, of the strength of ten grains to the ounce, would not be expected, even by the most sanguine believer in its deobstruent action, to cause the disappearance of an urethral stricture. Why, then, should it remove a similar pathological condition from the tympanum? If those who obtain good results from tympanic injections would record a few cases in which repeated injections of air altogether failed to relieve deafness, but in which tympanic injections succeeded, then I think there would be fewer sceptics as to their therapeutic value, while their *modus operandi* would still remain a mystery.

In this paper, inspissated mucus has not been mentioned as a factor of deafness in chronic non-suppurative inflammation of the middle ear, because most of the recorded cases seem to me to have been more or less subacute. It is difficult to conceive, in the first place, why this tendency to drying up should exist in a part whose air-supply must first pass through a portion of the respiratory tract; and, in the second place, how inspissated mucus can tend to remain long in a cavity where all the physical conditions seem in favour of absorption.

I have endeavoured to allude, shortly to the various points to be considered in forming a prognosis. It will, however, be understood that the various factors of deafness which have been discussed are often associated, and the different points of value in prognosis can only be utilised in combination.

CASE OF ACUTE BILIARY CIRRHOSIS,

CLINICALLY SIMULATING ACUTE YELLOW ATROPHY OF LIVER,
FATAL IN TWENTY-FOUR DAYS.

Read in the Section of Pathology.

By R. SHINGLETON SMITH, M.D. Lond., B.Sc., M.R.C.P.,
Physician to the Bristol Royal Infirmary.

JAMES P., aged 33; shoemaker, living in Clifton, married, having two children, was admitted to the Bristol Royal Infirmary on March 20th, 1883, with jaundice.

He stated that he had been quite well up to thirteen days previously. He was attacked in the night with diarrhoea, which was followed by pain across the epigastric, and right and left hypochondriac regions. The pain was severe, but it did not give rise to vomiting. He was not able to go to work on the day after, as the pain still continued; but two days afterwards the pain abated,

although he still felt some uneasiness in the region of the liver. The next day, he became jaundiced; he then remained in bed, and obtained medical advice. He admitted that he had been a hard drinker; but there was no definite history of syphilis.

For two weeks before admission, the patient thought he had a cold, and expectorated some phlegm, occasionally streaked with blood. There had also been some diarrhoea, and a difficulty in taking food, because of a constant feeling of sickness.

On admission, March 20th, the patient was seen to be very deeply jaundiced everywhere, and he complained of weakness. The heart-sounds were good; pulse 80, small. The lung-sounds were normal. There was no increase of splenic dulness. The liver-dulness was slightly increased; there was marked tenderness over the region of the liver, but not much complaint of pain otherwise. The gall-bladder could not be felt through the abdominal wall, but there appeared to be an increase of dulness under the right rectus muscle, as if from distension of the gall-bladder. The urine was of specific gravity 1030, acid, no albumen; bile was abundant. The patient's temperature was normal. The tongue moist, slightly coated.

March 26th. The skin was rather less yellow. There was no complaint of pain, but there had been slight epistaxis three times.

March 27th. There was more epistaxis, about eighteen ounces; and the anterior nares were plugged. The faecal evacuation was darker, possibly from intermixture with swallowed blood. Pulse 100, small and weak. The tongue was notched and coated. The liver-dulness was normal.

March 28th. He had more epistaxis, to about twenty ounces, which was only stopped by plugging. He was delirious all the night, jumping out of bed, and making a noise. A hypodermic injection of a fourth of a grain of morphia was given at 4 A.M. At noon he was found to be comatose, with pupils contracted. Pulse 88, regular, compressible, but moderately full; respiration slow, deep, and laboured. The temperature was normal; the tongue dry. There was slight oedema of the legs. There was no marked blanching of the lips, nor any other result of hæmorrhage. The urine was drawn off with the catheter.

March 29th. The patient had slept almost continuously, but could be roused, and would sometimes speak. Pulse 96; respiration 12. The pupils were of medium size, equal. The urine was drawn off; it was of specific gravity 1013, acid, and contained no albumen, but much bile. There had been no more epistaxis. A soft systolic murmur was heard, of greatest intensity at the third left cartilage, and conducted along the vessels, inaudible at the apex. The oedema of the legs was increasing. There were no petechial spots. The patient took milk and soda-water abundantly; he had also six ounces of brandy.

March 31st. A colocynth enema brought away a light coloured stool. Pulse 92; respiration 20. The patient was more comatose, and could not be roused. The pupils were dilated, but sensitive. The liver-dulness was less than before; three inches vertically in nipple-line. The oedema of the legs was increasing. The murmur at the base was as before. Death took place on the same day.

At the necropsy it was found that the liver weighed forty-nine ounces. The gall-bladder contained a thin yellow fluid, but it was not distended; there were no calculi within it. A catheter was passed along the duct into the duodenum, but no obstruction or stricture could be found. The substance of the liver was hard and tough; it was of a dark yellow copper-colour, mottled with lighter parts between; there was no roughness of surface, and no abnormal flattening of the organ. In section, all the lobules were distinctly visible, the individual lobules being isolated from each other by a grey intervening substance; the lobules were of a dark yellow colour, affording a marked contrast to the grey substance between. There was no dilatation of the larger bile-ducts within the liver, and no bile-staining of the sections of the ducts in the portal canals. The stomach contained some altered blood. The duodenum presented no appearance of inflammation or ulceration. The urine drawn from the bladder contained much epithelium of various kinds, spermatozoa, mucus-corpules, etc., but on evaporation it did not yield crystals of either leucine or tyrosine. Uric acid and cubes of chloride of sodium were seen, but no globes or needles. No renal casts or renal cells were observed.

On microscopic examination, the individual lobules of the liver were found to be separated by a connective-tissue growth giving the sections the well known appearance of the normal liver of the pig; and were of a bright yellow colour, the intervening substance colourless. They had not diminished in size; the cells were quite distinct and well defined. No distinct crystals of hæmatoidin or of tyrosine could be discovered in any of the sections. On stain-

ing with gentian-violet, the fibrous new growth became deeply stained, numerous small nuclei now becoming visible, and the nuclei of the cells in the lobules could be seen through the surrounding pigment and fatty granules. Minute dark-coloured spots like micrococci were visible, but could not be distinguished from the multitudinous pigment-granules of the liver cells. The sections of the liver showed numerous branches of the hepatic ducts running through the new tissue between the lobules; in some parts they were so numerous as to form quite a plexus of tubes, all containing a cubical-celled epithelium with large nuclei.

REMARKS.—This case, from the clinical aspect, presented considerable difficulty as regards diagnosis. It was at first thought the cause of the jaundice must be catarrhal obstruction; and this view was borne out by the completeness or intensity of the jaundice, by the presence of diarrhoea, the suddenness of the attack, and the presence of some little pain. The pain was not of sufficient intensity to justify the diagnosis of gall-stone impacted in the duct, and there was no decided evidence of distension of the gall-bladder. The disease was too acute in character for chronic atrophy; the jaundice was much deeper than is usually found in the ordinary forms of cirrhosis; and this hypothesis appeared to be untenable, in consequence of the absence of ascites and of enlargement of the abdominal veins. The symptoms were more in accord with the clinical features of acute yellow atrophy than with any other recognised pathological condition; the early occurrence of head-symptoms, the profuse epistaxis, and speedy death from gradually increasing coma, were, at least, suggestive of acute yellow atrophy; but, on the other hand the liver dulness was not less than normal, and was not observed to diminish from day to day, there were no petechial spots or bruises; no leucine or tyrosine could be obtained from the urine, and the jaundice was deeper than is commonly observed.

It was only at the *post mortem* examination that a definite opinion as to the pathology of the case could be given; and even then, although it was at once evident that a condition of cirrhosis was present, the appearances differed markedly from those commonly observed in the usual forms of chronic interstitial inflammation of the liver.

Cases of jaundice appear to be not uncommon, which cannot be classified under the ordinary typical forms of liver-disease, and these "residual phenomena" are variously explained and described by different observers. These are the cases described by Goodeve (Reynolds's *System of Medicine*) as "malignant jaundice." Thierfelder (Ziemssen's *Cyclopædia*) describes them as rarer forms of interstitial inflammation, and alludes to three varieties, the unilobular, or insular, cirrhosis, as distinguished from the multilobular, or ordinary, cirrhosis; interstitial hepatitis arising from affections of the gall-ducts; and the hypertrophic cirrhosis of Olivier, Hayem, and others.

This last class is the one which most nearly adapts itself to the clinical and pathological condition of the case now reported. The characteristic feature of this class is that the diminution in volume of the organ, resulting from the shrivelling of the newly formed connective tissue, such as characterises the later stage of ordinary cirrhosis, does not supervene. Decided icterus makes its appearance early, and persists, with fluctuations, during the entire course. No obstruction of secretion or inflammation can be demonstrated in the larger bile-ducts, but the interlobular canals become dilated and ramify into finer ducts, forming, immediately in front of the margin of the lobule, a network, from which still finer branches proceed, to disappear between the hepatic cells. All these canals are lined with cubical epithelium, and, in the ultimate divisions, the calibre often appears to be plugged with this epithelium. These tubes appear to be none other than the network of bile-capillaries, which, in some unexplained way, have acquired an epithelial lining.

The first symptoms are usually those of derangement of the digestive function; pain, icterus, and hepatic swelling follow. Progressive emaciation, cachexia, hæmorrhages, and death after severe cerebral symptoms, occur. Ascites is not usually present; it may be absent even in patients already very far reduced; more frequently it appears as a concomitant of general dropsy, induced by cachexia. There is usually enlargement of the spleen. The duration of the disease seldom amounts to less than a year; generally two years and over; in individual cases, it is stated to be five and seven years.

The case differs from the above description in some important respects; its acute onset, the absence of decided enlargement of the liver or of the spleen, and its speedily fatal termination; but *post mortem* examination disclosed the existence of a chronic interstitial hepatitis, in spite of the absence of symptoms till twenty-

four days before death, and also proved that the conditions present in acute yellow atrophy and various other well known pathological conditions were absent. The acuteness of the attack, and the speedy death of the patient, therefore stamp the case as one of cirrhosis of an almost unique character, and suggests a caution as regards the prognosis in other cases of apparent catarrhal jaundice.

Dr. Saundby (*Pathological Transactions*, vol. xxx, 1879) has endeavoured to show that the new formation of canaliculi is by no means essential or pathognomonic of the cases in which the three clinical features—*a*, jaundice; *b*, enlargement of liver; and *c*, absence of ascites—are associated. He shows that there may be very little alteration of ducts in cases of this type; and that, on the other hand, in cases of undoubted alcoholic cirrhosis, there may be a multiplication of biliary canaliculi; as also in the cirrhotic livers of tubercular patients, and in some forms of atrophic nutmeg-liver.

On the other hand, Bamberger reports a case of biliary cirrhosis of the liver, as follows (*Allgemeine Wiener Med. Zeit.*, No. 49, 1882). The patient had jaundice two years, with general wasting, but no pain. The urine was deeply coloured; the stools were always light, and at times colourless. Distinct enlargement of the liver could be felt; and, at one point, distinct puckering of the right lobe. The spleen was also enlarged. There was no ascites whatever. The occasional appearance of bile in the motions proved that the obstruction was not complete. In regard to the diagnosis of this case, he considers that a form of cirrhosis exists which is unaccompanied by ascites, viz., biliary cirrhosis—a form which has its origin in the biliary passages alone. It may be induced by catarrh, followed by concretions, but consists essentially of an overgrowth of connective tissue about the ducts. Hence the important and indicative symptom is the jaundice, ascites being only a secondary phenomenon. (*London Med. Record*, June 15th, 1883.)

The case I have described quite supports the view adopted by Thierfelder, Bamberger, and others, that a form of cirrhosis exists in which atrophy of liver-tissue is not present; jaundice occurs early and is very intense, as if from complete obstruction of the main duct; and ascites does not occur. The most prominent pathological condition is the presence of a great development of new bile-ducts along the interlobular passages, around which there is an overgrowth of connective tissue, resulting in a general unilobular cirrhosis. The term "biliary cirrhosis" indicates the distinguishing feature, and points to the bile-ducts as the starting-point of the lesion; this term may well be retained to distinguish the disease from the ordinary multilobular cirrhosis, in which the growth of connective tissue may be much greater, but the development of new bile-ducts is not a striking feature.

CLINICAL MEMORANDA.

PICRIC ACID AS A TEST FOR ALBUMEN.

IN the JOURNAL of January 5th (p. 10) there appeared a brief abstract from the *Maryland Medical Journal*, to the effect that Drs. Cook and Watkins had found that the urine of patients taking cinchonidia "gives the same reaction, with picric acid, as if they had albuminuria." If this is really the conclusion at which these observers have arrived, they must be unaware of the facts mentioned in my lecture, published in the JOURNAL, December 8th, 1883 (p. 1106), that "most of the vegetable alkaloids, such as morphia, atropine, etc., are precipitated by picric acid, and by the potassio-mercuric iodide; but quinine is the only one which is likely to be taken in sufficient quantity to render the urine opalescent with either of these tests, and then the complete clearance by heat at once distinguishes it from an albuminous precipitate." One of my pupils, a healthy young man, volunteered to swallow ten grains of sulphate of cinchonidia in water. The urine passed two hours afterwards was rendered opaque and milky by picric acid, but the transparency was completely restored by boiling, which is the exact converse of what happens when albuminous urine, which has been rendered opaque by picric acid, is exposed to heat. I repeat here a statement which I made in the lecture before referred to: "There is no known substance occurring in either normal or abnormal urine, except albumen, which gives a precipitate with picric acid insoluble by the subsequent application of heat."

The statement of the American observers that "the urine of patients taking cinchonidia gives the same reaction with picric acid as if they had albuminuria," is not less misleading and absurd than would be the assertion that "the urine of patients taking alkalies, or the alkaline salts with vegetable acids, e.g., citrate or tartrate of

potash, gives the same reaction with heat as if they had albuminuria," such an assertion implying ignorance of the fact that the turbidity produced by boiling neutral, or alkaline urine, is due to a deposit of phosphates, which is distinguished from one of albumen by its ready solubility on the addition of a minute quantity of acid, whether vegetable or mineral.

GEORGE JOHNSON, M.D., F.R.S.

PERSISTENT HICCUGH, TREATED BY CHLORAL.

ON July 4th, 1883, I was called to see Mr. W. H., aged 50. Two days previously, whilst walking in the street, he suddenly became giddy, reeled, and had to cling to a railing for support. With assistance, he got into a cab and was brought home. During the night he began to hiccough, at first only occasionally, but soon very frequently.

When first seen, it was about forty-eight hours after his attack in the street, and about forty hours since the hiccough began. He had tried all sorts of remedies, suggested by his wife and friends, also some medicine supplied by a neighbouring druggist; he had purged himself, produced vomiting, and blistered his abdomen, but was in no way relieved.

Bromide of potassium, belladonna, camphor, acetic acid, spirits of chloroform, hydrocyanic acid, fetid spirits of ammonia, morphia, pressure on the phrenic nerve, and over the epigastrium, were all given a trial. None of these had any effect, even large doses of morphia, hypodermically, only producing sleep, during which the hiccough continued as before.

At last, after twelve days' incessant suffering, thirty grains of chloral-hydrate induced sound and undisturbed sleep for six hours, and the patient awakened, free from his tormentor.

I have since had several opportunities of trying chloral, under similar, though less aggravated circumstances, and have always found that a liberal administration of this drug produced prompt relief.

Leavitt recommended chloral for hiccough in the *Medical and Chirurgical Review* of 1871.

GEO. C. KINGSBURY, M.A., M.D., Blackpool.

OBSTETRIC MEMORANDA.

INJECTION OF HOT WATER IN POST PARTUM HÆMORRHAGE.

AS every case bearing upon the treatment of *post partum* hæmorrhage is of consequence, I venture to record the following. On January 1st (this year), I was summoned to a young lady, a primipara, who had been six hours in labour; everything appeared to have gone on satisfactorily, the head (first position) being, at my arrival, almost on the perineum; the latter, however, was somewhat rigid. After upwards of two hours had elapsed, during which time I dilated the perineum, as little or no progress had been made, and the pains were gradually diminishing in frequency and force, I administered chloroform (a little had previously been given), and applied the forceps; before extracting, however, I gave a subcutaneous injection of ergotine to guard against the occurrence of *post partum* hæmorrhage. The extraction occupied about ten minutes, and was followed by the customary amount of discharge; the nurse, by my direction, carefully following up the child during the birth, and afterwards maintaining firm pressure on the abdomen. As the child did not breathe, I was some four or five minutes in exciting respiration; just as this had been satisfactorily accomplished, I was alarmed by the sound of a tremendous gush of fluid in the bed. Hurrying to the patient, I took the nurse's place, and made firm pressure on the uterus, which was quite relaxed, and at once removed the placenta; this was followed by another torrent of blood. I immediately introduced my hand into the uterus, and cleared it of clots, making counter-pressure outside; and, although there was a feeble attempt at contraction, the hæmorrhage continued. Fortunately, I had my Higginson's syringe at hand, and a long uterine tube ready attached; there was also a can of very hot water in the room; so, without a moment's loss of time, I introduced the tube, and injected some of the water (the hand could just be borne in it), with the effect of producing almost immediate contraction and cessation of hæmorrhage. At the same time, I administered some beef-tea, as hot as the patient could drink it, along with brandy and champagne. Although there was no further hæmorrhage, I gave a second injection of ergotine, and also kept up firm pressure on the abdomen, with the occasional application of an iced cloth. Under the influence of