prepare evidence. The greatest objection to an examination forerunning the trial is that it would be almost impossible to prevent it from eliciting confession of the deed, which would often be embarrassing and contrary to the spirit of our law (although, in France, as you may know, confession is encouraged or provoked). A solicitor for the defence would decide whether this danger existed or not, and would have a mental examination instituted or not, as he thought best for his client. An official examination, forerunning the trial, which had the misfortune to elicit a confession fatal to the prisoner would, I think, be condemned by English opinion. I do not know what legal right the prosecution or the executive has to order the examination of a prisoner committed for trial.

The examination of a man who has already been condemned to death for murder, and the formation and expression of an opinion as to whether he is insane in such a manner or degree that he ought to be reprieved, or not so, and therefore ought to be executed, imposes such a burthen of responsibility, that it ought not to be laid upon any man without those easements in the discharge of duty which lighten the responsibility of all who take part in judicial investigations.

From every consideration, therefore, of judicial method and consistency, the present mode of dealing with condemned men suspected of insanity, ought to be abolished, and a systematic investigation by sworn examiners, who should give public evidence subject to cross-examination, under the control of a judge, should be substituted.

NOTES OF A SUCCESSFUL CASE OF OPERATION FOR STRANGULATED UMBILICAL HERNIA, IN A PATIENT SIXTY-EIGHT YEARS OLD.

By TREVOR FOWLER, L.K.Q.C.P.I., and L.R.C.S.I.,
District Medical Officer Epping Union, etc.

On the morning of October 20th, 1883, I was called in to see Mrs. B., a very stout lady, aged 68, whom I found suffering from an old irreducible umbilical hernia, which had become strangulated from the previous evening. Pain had been considerable during the night, and there had been some vomiting. Upon the first accession of pain, the patient noticed an increase in size of the hernia, accompanied by an unusual hardness.

In the months of January and February previously, I had attended her for an attack of typhlitis; and in the month of May I was sent for in consequence of the umbilical hernia having become strangulated. The taxis, however, was successful upon this occasion, and a Moemain truss, with a concave pad, was advised to be worn. All went on well until October 20th, when, as already stated, the hernia again became strangulated; and, upon the taxis failing, the patient was put upon opium, an enema administered, and steps were taken to obtain further assistance, with a view to an operation.

On Sunday, October 21st, Mr. Walsham kindly came down from London, and upon consultation it was decided to give chloroform, and failing then to reduce the hernia, to cut down upon the seat of strangulation. The hernia was about the size of the double fist, and lobulated in shape, a fresh protrusion having apparently occurred at its upper part. It was tense, non-resonant, and painful on handling. There was no impulse on cough, and the skin over it at one part was mottled and dusky. The constipation was complete, no flatus passed, and vomiting occurred after any attempt at taking food. At 8 P.M., the patient being under chloroform and the taxis having again failed, Mr. Walsham made a longitudinal incision over the hernia about three inches in length, in the mesial line, and after cutting cautiously down upon the sac and opening it, the seat of stricture was reached and cut through with a hernia-knife, and the bowel was returned. The latter presented a dark purple or claret colour, but as it was still smooth and glistening, there was no question about the propriety of returning it. The greater portion of the contents of the original tumour, consisting entirely of omentum, was removed in separate pieces, each piece being first ligatured with carbolic catgut. The edges of the wound were brought together with eight silver wire sutures; pieces of gutta-percha tissue were inserted for drainage, and then the whole was covered up with eucalyptus gauze, pads, and bandages. The operation, and the subsequent dressings, were carried out antiseptically. At 9.30, the patient having recovered from the chloroform, thirty minims of tincture of opium were given, and five minims ordered to be repeated every two hours if awake; no food to be taken.

On the 22nd, 9.30 A.M., the patient had passed a good night. Pulse, 112; temperature, 100.4°. There was no abdominal tenderness. At 6 P.M. the temperature was 101.4°; at 7.30 P.M., pulse, 112, tempera-

ture, 102.1°. The pupils were contracted, and the patient was drowsy. The opium was continued as before. Ice was allowed.

On the 23rd, at 8 A.M., she had passed a good night. Pulse, 100; temperature, 100.3°. There was free diaphragmatic breathing. At 8 P.M. the pulse was 104, and temperature 101°. The pupils were contracted and the patient drowsy. The wound was now dressed for the first time. The lower portion of the remains of the hernial tumour looked decidedly dark; otherwise the wound was going on well. An extra dose of opium was given after dressing, and ordered to be continued as heretofore. Ice was allowed, also soda-water and milk in small quantities.

On the 24th, at 8 A.M., she had passed a good night. Pulse, 96; temperature, 99.2°. Flatus had been passed several times during the night. The wound was dressed again to-day. The pupils were still contracted, and the patient drowsy. Opium, ice, milk, and soda-water were given as before.

On the 25th she was still doing well. Flatus continued to be passed. The wound was dressed again. A considerable portion of old hernial tumour was beginning to slough. Charcoal poultices were applied. The opium, etc., were continued.

On the next three days the pulse and temperature were normal. The wound was dressed daily, and the opium, ice, milk, and soda-water were continued.

On the 29th I began to reduce the opium. Beef-tea was allowed in small quantities.

On the 30th, the bowels acted freely three times, and continued to act almost daily afterwards. From this time the patient steadily improved. The sloughs were slow in separating, and there was some burrowing of pus about the wound, which did not finally heal until November 26th. Recovery was also retarded by the formation of several very painful boils on the back. The patient, however, was able to leave her room on December 2nd, exactly six weeks from the date of the operation. She is now in perfect health, and the operation may be said to have been more than successful, inasmuch as it has effected a radical cure of the hernia by the complete obliteration of the sac.

The case presents several points of interest: 1. The recovery of a patient, at the advanced age of 68, from so serious an operation; 2. The radical cure of the hernia, which appears to have been accomplished by inflammatory adhesions about the neck of the sac sealing up the ring, which then became consolidated by granulation and contraction of the wound; 3. The illustration which the case offers of the value of pushing opium, and of the absolute safety with which this may be done, provided it be only given, as here, in small doses at a time, at frequent intervals, and with strict injunctions to the nurse that the patient is never to be awakened for his dose.

THE LIVERPOOL POISONINGS: CASE OF MARGARET JENNINGS.

By FREDERICK W. LOWNDES, M.R.C.S.,
Surgeon to the Liverpool Police.

In addition to the three cases fully and ably detailed by Dr. Whitford in the JOURNAL of the 16th instant, there was a fourth case, in which the whole responsibility of the exhumation and examination was left to me. The following are details condensed from my report to the authorities and my evidence at the recent trial.

On November 16th, 1883, I attended at Ford Cemetery, near Liverpool, at the request of the police-authorities, to witness the exhumation and to examine the body of Margaret Jennings, who had died ten months previously, under circumstances suggestive of her having been poisoned by some irritant, most probably arsenic. The cemetery is situated four miles from Liverpool, and is for the exclusive use of Roman Catholics. On my arrival, shortly before 9 A.M., I was conducted to the public portion, which stands higher than the other part of the cemetery, is formed of sand mixed with clay, and appeared dry and well suited for its purpose. I was shown a grave which had been just opened, and from which two bodies had been already removed. A coffin was lying at the bottom, from which a plate had been removed; it bore the inscription, "Margaret Jennings, died 25th January, 1883, aged 18 years." The coffin was a pine one, and perfectly intact; it and the coffin-plate were identified by the man who made them and conducted the funeral. The death had taken place within the jurisdiction of the coroner of Liverpool (Mr. Clarke Aspinall); and, as the cemetery was in the district of the county coroner, the late Mr. Barker, it was arranged by both gentlemen that permission should be obtained from the Home Secretary to remove the body to Liverpool. The coffin w