The lecturer then stated his opinions as to Localisation, limiting consideration to movements. For illustration, he imagined a centre for the arm and the leg. The "Universaliser" would say that every part of the centre represented both the arm and the leg in the same way; the "Localiser" would say that one part of it represented the arm only, and another part the leg only—would, indeed, make two centres. This is the current hypothesis. The lecturer did not take either of these views; he restated the opinion which he had held as to localisation, both before and since the recent experiments of Hitzig, Ferrier, and others on the mid-cerebral cortex.

"It may be said that one convulsion will represent only the movements of the arm, another only those of speech, another only those of the leg, and so on. The facts above stated show that this is not the plan of structure of the nervous system. Thus, to take an illustration, the external parts, x , y , and z , are each represented by units of the corpus striatum. But the plan of representation is not that some units contain x largely only, as x_s , others y largely only, as y_s , but that each unit contains x , y , and z —some, let us say, as x_s , y_s , z_s ; others as x_s , y_s , z_s , etc. When we come to the still higher evolution of the cerebrum, we can easily understand that, if the same plan be carried out, a square inch of convulsion may be wanting, without palsy of the face, arm, and leg, as x , y , and z are represented in other convulsions; and we can also easily understand that discharge of a square inch of convulsion must put in excessive movement the whole region; for it contains processes representing x , y , and z , with grey matter in exact proportion to the degree of complexity." (*St. Andrew's Medical Grad. Trans.*, 1870.) This is only another way of stating the doctrine of compensation and its "inversion," co-operation in excess (see second lecture). This double doctrine, he thinks, applies to representation of movements in the mid-cerebral region of dogs. Destruction of a part of the dog's mid-cortex entails no permanent paralysis, but discharge of any part of it produces convulsion. Ferrier, Charcot, and others have shown, since the part quoted was written, that destructive mid-cortical lesions in monkeys and men entail some permanent local paralysis; in these animals, it would seem that there is less compensation than in dogs. The important question is—Is there some recovery, and thus some compensation, in the case of man? The lecturer did not now suppose compensation for any destructive lesion to be ever absolute. His hypothesis was that the so-called centre for the arm represents that part very specially, but that it also represents the leg and face more generally, and the parts on the other side of the body still more generally; similarly, *mutatis mutandis* for the other centres. According to this view, destruction of the arm-centre should entail some decided paralysis of that limb, and very little, if any, of the leg and face, which two parts would have also a large representation in, so to speak, their own more particular centres. On the other hand, excessive discharge limited to the arm-centre should produce convulsion, beginning in that limb, and yet spreading beyond it to the leg, face, etc. The currents developed would, however, not be limited to one centre; they would, no doubt, spread to collateral centres, and discharge them; thus a convulsion beginning in the arm and becoming universal, was no evidence that a single centre—the arm-centre—represents all four limbs.* Yet certain experiments by Franck and Pitres on dogs tended, the lecturer thought, to show that his view of localisation—that every centre represented all parts, but that each centre represented some one very specially—held in the case of these animals. They conclude from these experiments: "Elles démontrent, en effet, que l'épilepsie peut se généraliser bien que l'excitation provocatrice ait été limitée à un seul centre cortical [they isolated the arm-centre, and by stimulation of it produced convulsion of all four limbs] et que la destruction préalable d'un centre cortical [they cut out the arm-centre in another experiment] n'empêche pas l'extension des convulsions aux muscles correspondant au centre préalablement détruit."

It must be said that these distinguished physicians believe, as stated in the first lecture, that, "Le cerveau commence l'attaque, la protubérance, le bulbe et la moelle la généralisent." Ablation of the cortical centre, "primitivement excité" does not (Albertoni, Franck, and Pitres) stop the convulsions. Franck and Pitres' argument is very able and powerful; their contention deserves most respectful consideration. No doubt centres in the pons, medulla, and spinal cord (lowest centres), are discharged by the prior cortical discharge. At any rate, the experiments mentioned go towards showing that all the lowest centres

are under the command of—if they are not all, as the lecturer thinks, re-represented in—each part of the mid-cortex. The order of the discharge of all the lowest centres is determined by the particular part of the mid-cortex primarily discharged.

Admitting difficulties in the way of acceptance of his hypothesis of localisation, the lecturer thought the current hypothesis did not account for all the facts. Any adequate hypothesis had to account for the following: (1) that from a destructive lesion of motor centres the paralysis was especially of the more voluntary, etc.; (2) that from a discharging lesion, the development of movements (convulsion) was the same: an epileptiform seizure starting in the arm began, first of all, in the hand; one starting in the leg began in the foot; one starting in the face, in the side of the mouth; (3) that the progress in each (1 and 2) was in compound order; (4) that a part might be permanently imperfectly paralysed, and yet the whole of it might be occasionally convulsed; (5) that recovery, or some recovery, follows on permanent destructive lesions producing paralysis; there is often at least some compensation; (6) that a small destructive lesion of a motor centre may entail little or no obvious paralysis, whilst a sudden and excessive discharge beginning in such part produces (indirectly) very great convulsion; (7) that from discharges beginning in different parts of the mid-cortex, we have fits affecting the same regions, but their parts in different order ("isomeric seizures"); (8) that a patient, who has no paralysis before a convulsion, has much (temporary) paralysis after it. (Cases of monoplegia [the paralysis only existing after a convulsion] must not be adduced as evidence to show that any lesion found *post mortem*, say a tumour, had produced paralysis by its direct, its destructive, action.)

THE LUMLEIAN LECTURES ON THE ETIOLOGY OF PHTHISIS.

Delivered at the Royal College of Physicians.

By J. ANDREW, M.D., F.R.C.P.,
Physician to St. Bartholomew's Hospital.

LECTURE II.

THE causes of phthisis which come next under consideration are those connected with modes of life and industrial occupations. A mere enumeration will be almost sufficient in the case of the majority of the members of this group. It is all but impossible to separate the influences of the day-workshops and of the occupations from that of the houses, and especially of the sleeping-rooms, of the operatives, so that whatever arguments on either side might be drawn by a confident statistician could be of little value. The conclusions could scarcely be of greater weight than the premisses. Still, as the influence of the home-life is probably about the same as all occupations which are paid at the same rate, except, of course, those which must, from their nature, be carried on in specially unhealthy localities, it would be reasonable to attribute, in part at least, to the occupation any great variation from the phthisis-mortality in either direction; although it might be impossible to determine the exact proportion in which the conditions of labour and of home-life contribute to it. Still more justifiable would it be to do so in cases where general rules, such, for example, as that phthisis in country districts is more frequent among women, are interfered with. With these limitations, I would divide this group into two subgroups, viz.:

1. Those occupations and modes of life in which there is the possibility, the likelihood if you will, that direct contagion may come into play;
2. Those in which there is no greater danger of such contagion than all of us are exposed to in the ordinary intercourse of life.

1. All modes of life, all occupations which are carried on indoors, contrast unfavourably with outdoor pursuits. The naked savage, whatever ills he may have to bear, rarely finds phthisis among them; but with every addition to his clothing, and to the comfort of his tree or cave, his proneness to it increases. In this respect, in an advanced civilisation, the effeminacy or luxury of the rich and the necessities of the poor bring about the same result. Sometimes, perhaps, even members of our own profession are forgetful, in the advice they give, of the advantages of an open-air life. I remember more than one medical student with incipient phthisis, compelled by circumstances to undertake country practice in a bleak district, who, instead of being

* At that time, I arranged centres morphologically, and did not then make a division into highest and middle motor centres. I used to speak of convulsions being the corpus striatum "raised to a higher power." As stated in the second lecture, I now arrange centres on an anatomico-physiological basis.

* Of course, the term "centre" is an arbitrary expression; but using the term "centre for the arm," there would in it, one must suppose (judging from the mode of spreading of convulsion in that limb from discharge beginning in part of the centre), be subcentres.

injured by the constant exposure to all sorts of weather, has regained seemingly perfect health. In the case of trades, it is impossible to ascertain the true proportion of cases of tubercular phthisis included in the totals of deaths from pulmonary diseases; but the facts, such as they are, contained in Dr. Greenhow's "Local Inquiries into Excessive Mortality from Lung-diseases," printed in the Reports of the Medical Officer of the Privy Council for the year 1860-61, make it probable that here too, as in general statistics, when the death-rate from diseases of the lungs is large, that of phthisis is also excessive. Wherever imperfect ventilation of workshops, with or without overcrowding, exists, the possibility of direct contagion must be admitted, whatever may be the nature of the occupation. It would be superfluous to enumerate the well-known industries and localities in which this is the case. A very long, if not complete list, will be found in the paper just referred to; but, at the same time, it must be remembered that, apart from any special infection, exposure to an atmosphere vitiated in this way is invariably recognised as a most potent cause of ill-health, and as a predisposing cause of many other forms of disease besides phthisis.

2. Certain occupations, however, in which there is no overcrowding or want of ventilation, give an abnormally high death-rate from pulmonary diseases, and these constitute my second subgroup. Instead of enumerating the occupations themselves, it will be sufficient to mention the chief causes of their unhealthiness, and these appear to be the following:

- a. The presence in the air of fine dust of any kind, of carbonic or of other noxious gases.
- b. Exposure to great vicissitudes of temperature.
- c. Inhaling hot air, either very dry or very moist.
- d. The workman being obliged to carry on his labours in some cramped posture, interfering with the freedom of the respiratory movements.

It is clear that the excess of phthisis which Dr. Greenhow's facts prove to exist under these conditions cannot be due to direct contagion. But the complexity of the factors to be dealt with in considering the influence of occupation is so great, that its discussion would require a length and mode of treatment impossible in these lectures. I trust, however, that the majority of my hearers will agree in the probability of the conclusion just stated, viz., that the excess of phthisis in some trades cannot be altogether accounted for by direct contagion only. The causes of phthisis, so far briefly reviewed, have all been external to man's organism. Some of them are physical conditions more or less beyond his control; others artificial, due to his own mismanagement, and blots upon his so-called civilisation, yet one and all acting upon him from without. I pass now to a different order of facts, those connected with some special, though unknown modification of bodily constitution; which renders those in whom it exists more or less liable to be attacked by phthisis. Sometimes it is found in large numbers of men, either of one common stock, or, if unrelated, yet living under similar physical and social conditions, sometimes it may occur only in a few isolated individuals. It may be either inherited or congenital or acquired. It may be labelled, according to fancy, temperament, or diathesis, or idiosyncrasy, or constitution, or habit. But it is met with in so many different forms, and under such widely differing circumstances, that I find it difficult to believe that its nature is always the same, although its result, viz., susceptibility to phthisis, is so. It must be one of the more remote causes, not the proximate cause of the disease, and its production is one of the modes in which external agents exercise their injurious influence. It falls naturally under the heads of race, inheritance, congenital origin, and acquired diathesis.

It would be an unwarrantable trespass on your patience to treat any of these in detail, as if they still required proof; but it is necessary for my argument to give one or two illustrations in connection with their bearing upon the question of direct contagion.

In Peru, according to Dr. Alphonse Guilbert, in his well-known paper on pulmonary phthisis in Peru and Bolivia, the influence of race comes out in the clearest manner. He states (*op. cit.*, p. 20) that the different races now found in the country suffer from phthisis in the following order of severity:

1. Negroes;
 2. The descendants of the Spaniards who took part in the conquest;
 3. The half-breeds, in the various crosses between Spanish, Negro, and Indian strains of blood;
 4. Persons born in Europe, and in their case the disease is not only less frequent, but its average duration is longer;
 5. Indians of pure descent, among whom it is all but unknown.
- There are one or two points worth noticing here.

The well known proclivity of the negro race asserts itself—a proclivity which makes it difficult to believe that the interior of Africa could have escaped if phthisis were spread by direct contagion, and which also brings out into stronger relief the anomalous immunity of Senegambia and of the Gold Coast.

Again, it is interesting to note that, in little more than three hundred years, there has been developed among the Spaniards born in Peru a liability to phthisis far in excess of that prevailing among immigrants from the mother country.

The social condition of the half-breeds is much lower than that of the Spaniards. Their habits and modes of life are much more favourable to the spread of infectious diseases, and yet they suffer less from phthisis. Again, the native Indian race has not yet been attacked by this imported contagion, any more than the North American tribes had been in the time of Dr. Rush. Making every allowance for the influence of race and modes of life, etc., in the causation of phthisis, this presents a strange contrast to the history of the introduction of small-pox, measles, whooping-cough, and other infectious diseases, among savage peoples. Thus, several millions of Indians are said to have died from small-pox on the Pacific coast of North America within three or four years of the first appearance of that disease among them.

Directly, infectious diseases are ordinarily more destructive on their first introduction into a new country; but in Peru, in the South Sea Islands, and in Australia, according to Dr. Thompson, this does not hold good in the case of phthisis; its virulence increases as time goes on.

Again, a race may escape phthisis in one country, and yet suffer from it in another. Thus, the West Indian negro, not specially prone to it in his native island, becomes so when transferred to the Gold Coast; and this too, although his near relatives there escape almost entirely, and do not even suffer from the contagion which he may have brought with him. This long known liability of new-comers to suffer more than natives from certain diseases is a matter of great importance in reference to the properties of the contagion of those diseases. In the common case of a stranger visiting a country infected by a disease—*e.g.*, yellow fever—unknown in his own home, and at once falling a victim to it, there is certainly nothing unreasonable in the supposition that the native escapes in consequence of having been exposed to minute doses of the poison, and thus, as by an artificial inoculation, obtaining protection against its fatal effects. But it must be remembered that sometimes the native suffers, whilst the stranger escapes, although exposed for the first time to a virulent contagion. Thus, some regiments raised in the Soudan, and employed in Mexico in behalf of the cause of the unfortunate Emperor Maximilian, were proof against an outbreak of yellow fever which decimated the Mexicans themselves. Here the inoculation-theory fails utterly, and we are driven to believe that some peculiar racial constitution must be, in part, at least, the cause of the immunity.

There is, however, a third combination, of which the case of the West Indian soldiers on the Gold Coast is an instance; and in this neither of these explanations is admissible. Here we have two very closely allied races, or rather, we might say, members of the same race, living together, and yet the new-comers suffer excessively from a disease to which they are at least as much exposed in their own country as the natives of the Gold Coast, who comparatively escape it, are in theirs. Under these circumstances, the theory of race and that of inoculation seem to me alike insufficient. A better explanation is to be found in the supposition that, although we speak of phthisis in the West Indies and on the Gold Coast as one and the same disease, there is some difference in its exciting cause in the two regions; that on the Gold Coast it has some property unknown to it in the West Indies, which gives it increased power over the West Indian negro, but does not render it dangerous to the native of the Coast. Granted, for argument's sake, that the exciting cause is some micro-organism, then in the West Indies and on the Gold Coast we have, if not different species, at any rate varieties, of that micro-organism. Allow me to remind you that this supposition is supported by the existence of numerous parallel instances; *e.g.*, by the greater or less activity of many medicinal plants, according to the soil and climate in which they are grown; by the fact that the same species of fungus is wholesome food in one country and poisonous in another; and, above all, by the greater or less activity of some of the septicæmic bacilli, according to the manner in which they are cultivated. I shall again refer to this, in attempting to determine the nature of the contagion of phthisis.

The influence of inheritance is the same in kind as that of race, in the production of which it is indeed a principal factor. It is less widely distributed, but within its area of distribution is more intense.

It is necessary to reassert the fact that a predisposition to disease,

like any other personal peculiarity, such as the colour of the hair, may be congenital and extremely powerful, although there is no record of its previous occurrence in the family; and, further, that the congenital predisposition is frequently shown by several, or even by all the children of the same parents. Thus, when several cases of the same disease occur among brothers and sisters, it is undoubtedly quite right to note the fact, whether it is or is not inherited, but not to do so in such a way as to imply that, in the absence of inheritance, there can be no common constitutional predisposition.

Acquired Diathesis.—Its reality is attested by the frequently observed relation between the development of phthisis and attacks of other forms of disease, such as syphilis, the continued fevers, or inflammatory affections of the lungs; and also by the increased liability which follows upon deterioration of general health by exposure to hardships, privations, or insanitary modes of life.

Sex.—The difference in mortality seems to depend rather upon the difference in the occupations of the two sexes, and on their relative sobriety and morality, than on any physiological factor. In towns, especially in London, the male death-rate is the higher; in the more purely agricultural districts, the female; but, for the whole of England, in the year 1880, the Registrar-General's returns give a large excess on the male side, viz., males, 1,946, females, 1,807, per 1,000,000 living. It is, however, noteworthy that many writers, e.g., Dr. Walsh and Dr. Jas. Pollock, state that inherited phthisis is more common in women than in men.

Annual Report of Registrar-General for 1880, pp. 150-1.

	Total under 1 year.	1	2	3	4	Total under 5 years.
Males	521	433	191	108	93	1346
Females	443	408	173	111	98	1223

	5	10	15	20	25	35	45	55	65	75	85	95	All Ages.
Males ..	546	668	1683	2720	5214	4816	3387	1031	652	75	3	..	22,981
Females	624	1137	2648	3148	5904	4905	2517	1207	424	57	1	1	26,196

Age.—Most fatal in adult life; i.e., from the age of 15 to 45. In the above table, the actual number of deaths, not the percentage, on those living at the second ages, is given; so that the true mortality at the more advanced ages, is much higher than is shown by these figures.

The causes so far enumerated by no means exhaust the detail of the etiology of phthisis, but they readily fall into three main classes, which may be roughly termed climatic, social, and personal; and to one or other of these, if phthisis be a diathetic or developmental disease, every one of its possible causes may be referred. If, on the contrary, it depends upon some external exciting cause, of the nature of a morbid poison or parasite, then they would stand to this in the relation of predisposing causes. I say of a morbid poison or parasite, for the time has not yet come in which it is possible, in all cases, to draw a distinct line between the two. And yet, even now, diseases in which the existence of a micro-organism as their exciting cause has been established, ought to be no longer classified as depending upon morbid poisons. For, surely, a bacillus or a bacterium in the blood, or tissues is neither more nor less a parasite than a filaria or spirochete in the blood, than sarcina in the stomach, than a tapeworm in the intestines, than a trichina spiralis in the muscles, or a fluke in the liver. The chief difference among these unwelcome guests, being that some are certainly animal, whilst others are probably vegetable parasites; but the nature of their relation to their "host" is the same in all. However much they may vary in the region or tissue they infest, or in the effects produced by their vital activities, they all agree in this, that they live, for the time, at the expense of the sheltering organism, and, in so doing, bring upon it more or less damage or discomfort.

Is it possible, then, from the examination of these three groups of causes, to draw any practical conclusion as to the nature of phthisis? I believe it to be so, and that it is a necessary step on the road to any certain knowledge of a most important subject. For such an examination will make it plain that these causes, singly or in combination, are insufficient to account for the phenomena of phthisis. That it is not, cannot be, developmental or diathetic, but must be ranked among the specific febrile diseases, and that, therefore, some micro-organism, or other must be a necessary condition, and not a mere accident of its presence.

Unless we can find among these causes some one which is never absent, the very number and well-attested rights of the claimants be fatal to the claims of any one of them to be regarded as the sufficient cause. Were they each and all always present, and in definite proportions, then indeed there would be nothing paradoxical in the ascription of the morbid changes following their action. But that causes so diverse, so distinct, as dampness of soil, exposure to a hot/dry atmosphere, imperfect ventilation, an attack of pneumonia, or the being descended from a phthisical ancestor, or any other causes of the same sort, should give rise singly, or in combination, in all climates, under all known forms of barbarism or civilisation, to singular pathological changes, associated with a similar train of clinical symptoms, and this in spite of manifold differences in the personal factor—this cannot be accepted without the production of the most cogent proof, and in default of any simpler explanation. Indeed, the longer the complete list is made under these heads, the more signal does its insufficiency become. Each addition to the roll increases instead of diminishing our difficulty. Many of these conditions are the exciting causes of ordinary forms of disease; e.g., an ordinary bronchial catarrh is frequently induced by exposure to vicissitudes of temperature, or by breathing an irritating atmosphere, whether it be artificial, or supplied liberally by nature herself in the form of the spring east winds. That such causes should do this appears natural enough (unless, indeed, we try to realise the mode in which the effect is produced), but that they should also, from time to time, give rise, instead, to a disease with such special constitutional and local characters as phthisis, is hard to understand, except by giving what seems to me to be undue weight to the influence of constitutional predisposition. In point of fact, the only general developmental disease with which phthisis can be fairly compared in respect of its wide geographical distribution, and its practical uniformity under an infinite variety of conditions, is old age.

Except for the sake of maintaining a paradox, we must look for some one dominant factor, always and everywhere present, which can be shown, at any rate by analogy, to be sufficient to explain not only the general type of the disease, but also the anomalies in its distribution, and the variations presented by individual cases; and such a factor can only be found in some such a micro-organism as the bacillus of Koch. It cannot be found in any of the three groups of causes hitherto considered.

Not many discoveries of equal importance have received such speedy and strong confirmation, as that of the tubercle bacillus. I shall not reproduce here, much less attempt a variation upon, the report by Mr. W. Watson Cheyne to the Association for the Advancement of Medicine by Research, on the relation of micro-organisms to tuberculosis. Until it be disproved, I am well content to accept the bacillus of Koch as the essential cause of phthisis, and that, too, in the extreme form in which the doctrine has been stated to me by my colleague Dr. Klein, viz., "no tubercle without bacillus, no bacillus without tubercle." Even should it be proved that this particular bacillus does not hold this intimate relation to phthisis (although this becomes every day more unlikely), I shall still maintain the same view of the essential nature of phthisis, and look forward confidently to the eventual discovery of its true micro-organism.

Let us see how the hypothesis of a specific organic agent, present in all cases of phthisis as its proximate exciting cause, clears up and reconciles the obscurities and contradictions which exist in every branch of the subject.

I have already mentioned the great difficulty which, on the developmental or diathetic theory, is caused by the practical uniformity of the new growth, and the consequent improbability that it can be due, in different individuals, to so many and such widely different causes. And be it remembered that we are not dealing with a rare disease, of which so few instances have been observed that doubts might reasonably be entertained as to the existence of certain definite pathological changes, and their relation to clinical symptoms, but with one of the most common and most fatal of diseases, and one which has been known from the earliest times. This difficulty, if it did not consciously suggest, was met by the attempt to disprove the specific nature of pulmonary phthisis, by restricting the term "tubercle" to the miliary form only, and asserting that, in the great majority of cases of the disease, there was no tubercle present at all, but only the effects of non-specific irritations or inflammations; and that the similarity of the symptoms in so many cases was due to the fact that the same organs, the lungs, were attacked in each and not to any identity of the morbid agent or process. A still more unintelligible assertion was made regarding caseous matter, viz., that, although the product of processes having no connection with tubercular disease, it might, from time to time, the cause of a new growth, with such well marked special characters as those of miliary tubercle. The attempt

failed to produce universal conviction, and its partial success was in great part due to concentrating attention upon the results of what would now, perhaps, be regarded as very inadequate microscopic examinations, to the exclusion of the more general and less easily misunderstood anatomical and clinical phenomena. But surely these last are not the least important factors in determining the nature and relations of any new growth or morbid change. Until such time as our modes and instruments of research are so far improved that all discrepancy between the two ceases to exist; a difference in properties, however slight, must be held to prove a difference more or less complete in kind, in spite of apparent histological identity under the microscope. And, on the other hand, similarity in properties must be held to prove more or less close relationship, if not identity, in spite of histological diversities. All through the dispute, it was virtually admitted, by almost every one engaged in it, that there is something special in the lung of phthisis. The appearances which one man proved to his own entire satisfaction not to be "tubercle," he yet never doubted would be held to be such by some other observer. It must have seemed a little odd, now and then, that the results of so many different processes should be so readily mistaken for one and the same new growth. Those who, like myself, believed throughout in the similar and specific nature of the vast majority of cases of phthisis, and in the essential identity of the various forms of new growth found in their lungs, welcomed the discovery of the bacillus as a demonstration of the truth of that opinion. I have neither wish nor intention to undervalue the results or to depreciate the labours of the great histologists who endeavoured to solve the question of the nature of tubercle by what, with our imperfect resources, was an impossible method. We owe to them all that is exact in our knowledge of the external characters of the various forms of new growth produced by the same internal agent in different individuals and in different organs and tissues; forms which vary, of course, with variations in the personal factor in each case. But it is well to recognise the impossibility of success in any attempt to determine the nature of a pathological process or product by its histological characters alone.

For argument's sake, I have admitted the possibility of this being, at some future time, accomplished; but the prospect must be a very remote one, if the present views of physicists as to the constitution of matter are even approximately true. Thus Sir John Lubbock, in his address to the British Association at York, in 1881, stated that it has been calculated that the smallest sphere of organic matter which could be distinctly defined, when associated with others, by our most powerful microscopes, is about $\frac{1}{1000000}$ inch in diameter; that a particle of this size would contain many million molecules of albumen and water, and that, consequently, there may be an almost infinite number of structural characters in organic tissues, which we can at present foresee no mode of examining.

Compare the doubt and uncertainty of a few years ago with the certainty of to-day. Many of us must remember the dismay with which, if not himself the parent of some pathological bantling, he heard from the lips of an inquiring student the simple common question, "Pray, sir, what is tubercle?" For my own part, when placed in this predicament, having given in reply the latest, or what seemed to me the best, theory, I was generally careful to add that my answer was only provisional; that tubercle, next year, would probably be something quite different; but that phthisis remained the same, and that its macroscopical appearances would not materially change. Now, we feel no hesitation in answering this question. The presence of the bacillus is the essential condition, the others vary within certain limits.

Again, there is now no difficulty in our accepting the close relationship which the clinical observations of our predecessors undoubtedly established between the strumous diseases of childhood, especially those affecting the lymphatic glands, pulmonary phthisis and acute tuberculosis. The true statement that caseous matter may sometimes generate tubercle can no longer be made ridiculous by denying to that matter the possession of specific properties. For in all these instances the presence of the same material agent has been demonstrated. And thus it is no longer necessary to consign classical writings, like those of Sir Thomas Watson on the connection between scrofula and tubercle, to the limbo of medical mythology. Again, the anomalies in the geographical and local distribution of phthisis, and in its incidence upon different races, families, and individuals, the undoubted existence not only of general laws governing its manifestations in both these directions, but also of startling exceptions to those laws, admit of no rational explanation on the developmental theory, or, let me add by anticipation, on that of direct contagion. Given the bacillus, or some other micro-organism, and both sets of difficulties, the geographical and the anthropological, wide apart as they appear to be, receive the self-

same easy solution. The bacillus, whether inside or outside the body of man or of any other of its "hosts," cannot thrive except certain conditions be present. The analogy of other micro-organisms (e.g., that of the woolsorters' disease), and also of the higher classes of plants, proves how very slight are the changes which suffice to prevent or to favour the existence of vegetables, and probably also of the lowest members of the animal kingdom; thus the distribution even of many phanerogamous plants is extremely capricious. The exemption, then, of certain localities and of certain races from phthisis may be, is, due to the fact that the bacillus has been unable to contend with some adverse circumstance or other with which it has met, in the first case during the independent, in the second during the parasitical, phases of its life. Further illustration of these points will necessarily have to be given in discussing the question of direct contagion, and of the analogy between phthisis and ague. Notwithstanding, however, the great force of the arguments just stated in support of the theory that there exists some definite exciting cause of phthisis, and this cause is a "contagium vivum;" notwithstanding the presumption in the same direction, arising from the rapidly increasing number of instances in which changes taking place in organised and even in unorganised matter are being proved to be, if not entirely dependent, at any rate greatly promoted by, the presence of micro-organisms—e.g., the formation of nitric and nitrous acid from ammonia and the organic matters contained in the soil (Robert Warrington, *Journal of Chemical Society*, January 1878, July 1879); notwithstanding this, it might still be fairly objected that the alleged proof was inconclusive in this particular case, that the bacillus, or any similar thing, must be regarded as an attendant, and not a cause, of the morbid changes with which it is found associated, if it could be shown that the clinical characters of phthisis are inconsistent with this view of its origin. Now, with our present very imperfect knowledge of the limits within which the presence and activity of so-called "germs" are restricted, it would be rash to deny the possibility of any germ-disease being due to such a cause merely because its symptoms happen to differ widely from others whose dependence upon it has been established. But, in point of fact, the same theory which furnishes a ready explanation of one set of difficulties will be found not less effectual in accounting for the symptoms of tubercular phthisis in its regular and in its irregular forms.

For my present purpose, phthisis may be conveniently divided into two main classes, viz.:

1. Those in which the onset is insidious or gradual; and
2. Those in which it begins abruptly.

These latter fall naturally into two subgroups, viz.:

- a. Cases in which the symptoms of illness from the very first are those of phthisis, e.g., in so-called hæmorrhagic phthisis;
- b. Cases in which the disease follows sharply upon, or is even continuous with, some acute attack, it may be a pleurisy or a pneumonia, or some general fever, or syphilis.

The true typical features of the malady are exhibited most distinctively by the first or insidious form. Here, as in the case of other infectious diseases, certain symptoms precede its full development, and these it is useless to pretend to divide into two well marked groups, as belonging, the one to the period of incubation, the other to that of "primary fever." For, the line which theoretically separates them cannot be drawn with any certainty even in all the continued fevers; and in phthisis we have not yet obtained the data necessary for attempting to do so. Whatever view we adopt of the nature of its exciting cause and of its contagiousness, the period of incubation is absolutely unknown. There are, at least within my knowledge, no recorded instances of the disease following upon a single exposure, or even one of a few days only, to some source of infection; but, judging by the results of inoculation-experiments, the interval between the actual reception of the poison into the system and the production of its first effects, is probably a short one, and to be measured, if not by days, at any rate by weeks, rather than, as some seem disposed to assume, by months. Before there is any evidence to be gathered, either from physical signs or from symptoms of the existence of the pulmonary lesion, sometimes even for months previously, the patients are conscious of a general failure of strength and vigour, for which no adequate reason can be detected. Anæmia and slight loss of weight are the prominent features of this, the earliest, stage, and to these is not unfrequently added some special functional derangement which is naturally regarded as the cause of the *malaise*. Some form of indigestion is very commonly present, which may, however, amount to little more than loss of appetite, from inability to assimilate the usual quantity of food. Of course, phthisis by no means necessarily follows upon every slight indigestion, neither does pneumonia upon frontal headache, or some serious illness upon a rigor. But, in presence of phthisical antecedents, a state of debility, persistent yet seemingly

causeless, or manifestly out of proportion to some trivial functional derangement, e.g., gastric or uterine, ought always to be regarded with suspicious alarm.

Further, I have met with several cases in which, during this period, the urine was albuminous; the albumen sometimes disappearing before the development of the chest-symptoms, but without any corresponding impairment in health. Sometimes the albuminuria has persisted throughout the whole course of the disease. A close parallel to this is furnished by the relations of albuminuria to some forms of specific febrile diseases, i.e., scarlatina and typhus. I should have added rheumatic fever and pneumonia, but that their place in our nosologies is not yet determined.

These earlier symptoms are very often neglected, for the specific phthisis begins and ends only with the disease; and, by the time the lungs have been seriously involved and medical aid is sought, their very existence may have been forgotten. But careful inquiry almost always leads to the acknowledgment by the patient or his friends that for some time before the date at which he fixes the beginning of his illness, it may be by the occurrence of slight hæmoptysis, or by the access of cough, he was not quite so well as usual. Again, the night-sweats, or rather sleep-sweats, of phthisis are very often most profuse in the earlier parts of its course, when new growth in, rather than destruction of, lung is taking place, and gradually diminish or even cease, although the rapidity of tissue-disintegration and the quantity of muco-purulent expectoration is increasing. In fact, these sweats belong to the acute general stage of the disease, and are not merely part of the heat due to suppuration. This sequence of events is closely analogous to that of some of the common symptoms of the continued fevers, e.g., the headache and the gastric catarrh, so common during the first five or six days, and which then frequently come to an end. Add to these phenomena the rise of temperature which takes place at an early date, and the resemblance between gradually fully-developed forms of phthisis and the specific febrile diseases, especially those attended by some local lesion, become very complete, and supplies an additional reason for removing it from the class of developmental or diathetic diseases.

That phthisis should begin with greater violence and abruptness, after exposure to infection, in cases where the general health has been impaired or the lung itself injured by some previous disease, is only what might be expected, even from our short survey of its predisposing causes. Whilst the sudden explosion of acute tuberculosis during the course of ordinary pulmonary phthisis finds a ready explanation in the sudden entry of large numbers of bacilli into the blood, e.g., in consequence of the bursting of some softening tubercular mass into a vein. With regard to hæmorrhagic phthisis, the cases in which hæmorrhage, not due to already existing pulmonary disease, leads to consumption are probably exceedingly rare.

Thus, the clinical characters of phthisis, no less than the facts connected with its geographical distribution, and with its predisposing causes, harmonise so well with the hypothesis that some micro-organism or other is its exciting cause, and not a mere accidental attendant, that it seems to me all but impossible to come to any other conclusion, even apart from the results of inoculation-experiments. I have dwelt on this at what seems to be unnecessary length, because no one who denies it, or who fails to recognise the complete change in the manner in which the problem must now be stated, has any right to claim a hearing on the question of contagion. He would be no better fitted for such a task than the blind to discern a painting, or the deaf to hear music. But having stated my unreserved adhesion to the belief that phthisis is a member of the group of specific febrile diseases, and that therefore, like its congeners, it is dependent upon some morbid germ introduced from without, am I then obliged to admit that it is propagated by direct contagion from the sick to the healthy; or even that its transmission, through indirect channels, as in the case of typhoid fever or of cholera, is a matter of frequent occurrence? Most certainly not. The grounds of my disbelief in what is, once again, becoming the popular theory, I shall lay before you in the concluding lecture. Some of them have been already incidentally mentioned; but their full force only comes out when it is seen how some of the very facts which establish conclusively the specific nature of phthisis, place it also unmistakably among the non-contagious members of the group to which it belongs.

LORD BELPER has been appointed President of the Nottingham General Hospital for the ensuing year.

DR. HENRY COLEBROOKE, of Southborough, has been presented with an illuminated address, and a purse containing £500, by the subscription of 586 friends and patients.

A CLINICAL NOTE ON KAIRIN.¹

By ROBERT S. ARCHER, B.A., M.B., M.Ch.DUBL.,

Physician to the Netherfield Fever Hospital, Liverpool.

KAIRIN (or kairine) being a comparatively new drug, the action of which has not yet been thoroughly investigated in this country, and there being but few observations published on it in British medical literature, it is hoped that the clinical record of a case illustrating its antipyretic properties may prove of some interest to members of this Institution. Our experience of the medicine is limited to the case to be quoted below and another one; therefore, it is not intended to formulate any opinion as to the general use of the drug, whether with reference to its dose, or as to the class of cases it should be employed in. We simply lay before you a record which it will be yours to criticise, and to come to any conclusion each individual may see best.

Kairin is described in chemical language as oxy-chinolin-methyl-hydrate, the formula for it being $C_{10}H_{13}NO$, and is an artificial alkaloid, prepared from chinolin (C_9H_7O). It was discovered by Fischer and König, and its therapeutical action investigated first by Filehne and other German physicians. It is a yellowish white crystalline powder, with a rather acrid disagreeable bitter taste. It is soluble in water; but, on account of its taste, is recommended to be prescribed in gelatine capsules. (We used it in mucilage and water.) It has no perceptible odour.

I have here to express my sense of indebtedness to Dr. Oldham, resident medical officer at Netherfield Fever Hospital, for the care and precision with which he conducted the majority of the following observations.

CASE.—Wm. H., aged 19, was admitted to hospital on Wednesday November 14th, 1883, at 11.30 A.M., suffering from enteric fever. We could not procure a history of the case, and were therefore unable to judge accurately of the day of the fever.

November 14th. On admission, his pulse was 124; temperature 104.6° ; and respiration 36. The tongue was dry in the centre and moist at the edges. There were rose-coloured spots scattered over the abdomen. He was very heavy and stupid looking, aspect rather livid; no abdominal distension nor apparent tenderness; sordes on teeth; almost comatose, and quite helpless.

November 15th. He had passed a restless night; in other respects, he was much the same as yesterday, with the addition of a good deal of congestion of the left conjunctiva and some bleeding from the gums. At my visit at 11 A.M., the pulse being 128 and dicrotous, respiration 42, and temperature having risen to 104.6° , it was determined to try kairin in $7\frac{1}{2}$ grain doses. The first dose was not given till 2.30 P.M., when the pulse was 124, respiration 40, and temperature 103.3° . One hour after the kairin was given, the pulse was found to have fallen ten (10) beats, the respiration was unaltered, and the temperature was less by more than a degree and a half (1.5°). The patient was dozing; and the skin moist.—4.30 P.M. The pulse was 108, dicrotism gone; respiration 32; temperature 100.5° . The patient was sleeping, his skin perspiring freely. Another dose of kairin was given. 5.30 P.M. Pulse 108; respiration 30; a further fall of temperature to 99.9° ; sleeping; skin soft, but not perspiring. He had half an ounce of brandy at 5 P.M. No kairin was given. 7 P.M. Rigors. Half an ounce of brandy hot. 7.15 P.M. Temperature had run up to 105.6° ; pulse could not be counted at the wrist, but at the heart 130 beats were registered in the minute; respiration 40; skin dry; extremities livid; $7\frac{1}{2}$ grains of kairin were given. It will be noticed that, on the omission of kairin for an hour and three-quarters, the temperature had run up 5.7° , the pulse had become imperceptible at the wrist, and, on the whole, the patient may be said to have been almost moribund. At 7.30 P.M.; that is, a quarter of an hour after the kairin was given, the temperature had fallen close on a degree; the pulse was 126, could be counted at the wrist, but was slightly intermittent; respiration continued the same (40); skin soft, lividity had passed away; patient apparently dozing. 8 P.M. There was a further fall of temperature of nearly a degree; pulse 126, regular, not intermittent nor dicrotic; respiration 36; perspiring freely; dozing; slight cough; a few bronchial rales. 8.30 P.M. A slight rise in temperature, the medicine having been omitted for an hour and a quarter; therefore, a dose was given, with the effect that at 9.45 P.M. the temperature had fallen a little over a degree; pulse remaining the same, and respiration having risen slightly; skin moist; patient sleeping. Half a dose ($3\frac{1}{2}$ grains) was given, hoping that it might keep the temperature under control. 11 P.M. There was a slight rise of temperature to 103.1° ; pulse 119; respiration 32. Half a dose was given. 12 midnight. The temperature having risen half a degree, there also being a slight increase in the pulse and respiration rates, it became evident that the half dose,

¹ Read at a meeting of the Liverpool Medical Institution.