enforced about three months; the maximum of daily temperature having been noted, 98.5° on August 9th, when yellow fever had begun in Brownsville; and it had fallen to 65.5° on November 14th, when it had altogether ceased at Brownsville.

At Pensacola the disease raged between September 8th. and October 26th; the greatest number of attacks in any one day having been 77, and of deaths 11. The last attack occurred on November 28th, and the last death on the 29th. The general opinion held in the United States is, that no infected place can be considered safe until a frost has set in, which was not seen in those instances.

It is important to know that at Brownsville the majority of inhabitants were of Mexican, and at Pensacola, of Negro race; which accounts in a degree for the mildness of the disease.

This concludes the general information combined in the report of the supervising surgeon-general. The history of the epidemics is very instructive in a pathological sense, but that may lie beyond the immediate scope of this notice; which has been merely to show, at this moment, when a Bill is before Parliament to devise means to diminish sacrifice of life amongst our merchant-seamen, how much it is possible to effect in the direction of curing their diseases.

SUCCESSFUL REMOVAL OF TWO SOLID CIRCUMRENAL TUMOURS.

By SIR SPENCER WELLS, BART.,

Surgeon to the Queen's Household.

ANY student of abdominal surgery in general, and especially of what is now termed the "surgery of the kidney," is certain to be interested in noticing two specimens which have lately been added to the Museum of the Royal College of Surgeons. The following description of these portions of two solid tumours which I removed last November from around the right and left kidneys of a lady, with part of the left kidney attached to one of the tumours, has been prepared by Mr. Eve. One tumour weighed 16½ lb., and the other 14½ lb.

"The portion of kidney torn away comprises more than one-third of the lower end of the organ. A calyx and papilla may be seen upon the torn surface; they afford conclusive evidence that the pelvis was opened. Within the calyx is a small mass of fat. A section of the kidney has a normal appearance. Surrounding the surface of the kidney is a tolerably firm, greasy, pale yellowish white substance, evidently consisting in great part of adipose and fibrous tissues. It is in close contact with the capsule of the kidney, but is only adherent to it over a surface about half an inch in length. The capsule can be easily separated from the kidney at this point, leaving the subjacent parenchyma perfectly smooth.

"The portion of the tumour from the opposite side was slightly more fibrous in texture, but otherwise similar in its naked-eye characters.

"In minute structure, both tumours consisted chiefly of adipose tissue. The fat-cells were large, and closely approximated. There were also many large tracts and bands of fibrillar connective tissue, consisting of a close interlacement of delicate fibrils, which gave it somewhat the appearance of mucous connective tissue, but stellate cells were absent. The blood-vessels were large and numerous, and around them were masses of small round, so-called 'indifferent' cells, which could be traced through transitions, by elongation, to the formation of fibres.

"The formation of fat was evidently secondary to the growth of fibrous tissue, the advance of the fat-cells into it being apparent; while the various stages in the transformation of the fixed connective tissue-corpules into fat-cells were distinctly seen. Sections, cut through the point where the tumour was adherent to the kidney, showed that numerous parallel fasciculi of fibrous tissue intervened between the kidney-substance (which was healthy) and the tumour.

"The anatomical relations of the tumours to the kidneys, and the microscopic examination, support the view that the tumours originated neither in the pelvis nor capsule of the kidney, but in the circumrenal connective tissue.

"Histologically, they were fibro-lipomata, and no grounds exist for believing them to be in any part sarcomatous, since the young cell-element exhibited a distinct tendency to form mature tissues, either fibrous or fatty. A complete immunity from further disease may, therefore, be anticipated, provided that the growths were completely removed."

I removed the two tumours at Tooting on November 13th, 1883, assisted by Mr. Meredith and Dr. Ridley Hilder. Bichloride of

methylene was given by Dr. Day. I had been very doubtful as to the precise character and origin of the tumours before operation, simply saying that there were two solid tumours, so movable that I did not think there could be any unusual difficulty in their removal. On making the incision usual in ovariectomy, a growth resembling an ordinary fatty tumour was seen to be covered by layers of loose serous membrane, just as in tumours covered by the broad ligaments of the uterus. This membranous capsule was divided, and the tumour easily shelled out from a large deep cavity in the left side of the pelvis and left loin. Attached to the deepest part of the tumour was a piece of dark red tissue, which I at first thought was a part of the spleen; but which afterwards proved to be about a third of the left kidney. The remaining portion of the kidney and the ureter were then seen at the bottom of the capsule. There was very little bleeding. The capsular cavity was carefully sponged, and the edges of the divided capsule were placed in apposition, but no sutures were used.

Another large tumour was then found similarly situated on the right side, and was removed in the same way; but the right kidney was neither injured nor seen. The only other difference in the proceeding on the two sides was, that the ascending colon led to no difficulty; while, on the left side, the descending colon was attached to the tumour anteriorly, and was pushed aside after dividing the serous capsule of the tumour. There was not much blood lost, only a few small vessels requiring ligature.

The patient recovered quite as well as after an ordinary ovariectomy. In quantity and appearance, the urine was quite normal. Dr. Hilder once or twice found some small clots of blood in the deposit, but, he said, "none as large as a pin's head." I saw her last month quite convalescent.

I need only add that the lady was forty-eight years of age; was married when twenty-one; had never been pregnant; had begun to suffer from abdominal trouble in 1874; and in 1878 was sufficiently large for pregnancy to be suspected. The enlargement had gradually increased. Ever since 1881, owing to prolapsus of the uterus, a ring pessary had been worn until the day of operation, and numbness and stiffness of the right leg had been troublesome. The catamenia had been regular, and the urine was always normal when examined.

MIRYACHIT: A NEWLY DESCRIBED DISEASE OF THE NERVOUS SYSTEM AND ITS ANALOGUES.

Read before the New York Neurological Society.

By WILLIAM A. HAMMOND, M.D.,

Surgeon-General of the United States Army, Retired List; Professor of Diseases of the Mind and Nervous System in the New York Post-Graduate Medical School and Hospital.

In a very interesting account of a journey from the Pacific Ocean through Asia to the United States, by Lieutenant B. H. Buckingham and Ensigns George C. Foulk and Walter McLean,¹ United States Navy, I find an affection of the nervous system described, which, on account of its remarkable characteristics, as well as by reason of certain known analogies, I think should be brought to the special notice of the medical profession. I quote from the work referred to the following account of this disease. The party is on the Ussuri River, not far from its junction with the Amur, in Eastern Siberia.

"While we were walking on the bank here, we observed our messmate, the captain of the general staff (of the Russian army) approach the steward of the boat suddenly, and, without any apparent reason or remark, clap his hands before his face; instantly the steward clapped his hands in the same manner, put on an angry look, and passed on. The incident was somewhat curious, as it involved a degree of familiarity with the steward hardly to have been expected. After this we observed a number of queer performances of the steward, and finally comprehended the situation. It seemed that he was afflicted with a peculiar mental or nervous disease, which forced him to imitate everything suddenly presented to his senses. Thus, when the captain slapped the paddle-box suddenly in the presence of the steward, the latter instantly gave it a similar thump; or if any noise were made suddenly, he seemed compelled against his will to imitate it instantly, and with remarkable accuracy. To annoy him, some of the passengers imitated pigs grunting, or called out absurd names; others clapped their hands and shouted, jumped, or threw their hats on the deck suddenly, and

¹ "Observations upon the Korean Coast Japanese, Japanese Korean Ports and Siberia, made during a journey from the Asiatic station to the United States through Siberia to Europe, June 3rd to September 8th, 1882." Published by the United States Navy Department, Washington, 1883, pp. 51.