TWO LECTURES ON SCLEROSIS OF THE SPINAL CORD.

By JULIUS ALTHAUS, M.D., M.R.C.P.,

Senior Physician to the Hospital for Epilepsy and Paralysis, Regent's Park.

LECTURE I.

EVOLUTION OF STRANDS; MORBID ANATOMY; PATHOGENESIS.

ALTHOUGH the term sclerosis, or induration, is now-a-days indiscriminately applied to a variety of diseases of the spinal cord, both primary and secondary, it should be understood that we do not always find a very pronounced hardening of the affected parts. In some few instances, indeed, where the disease is comparatively fresh, the affected structures may even appear softened; yet where the morbid process has been going on for a considerable time (and such cases constitute the immense majority of those which are submitted to our inspection) the parts are decidedly tougher and harder than the normal ones, and may therefore be properly called sclerosed.

By sclerosis of the spinal cord we mean an irritant morbid process which invades certain well-defined, and evolutionally, anatomically, and physiologically distinct areas or systems of that organ; and which leads, in course of time, to disintegration and wasting of the nerve-tubes, often to destruction of the axis-cylinders, and always to overgrowth of intertubular connective tissue. The best known of these affections is one with which we have long been familiar as *tabes dorsalis*, more recent synonyms being *progressive locomotor ataxy* (Duchenne), *posterior leuco-myelitis* (Vulpian), and *sclerosis of the posterior columns* (Erb). The name *locomotor ataxy* does not appear to be happily chosen, as the ataxy of gait which constitutes the most prominent symptom of the fully developed malady is in many cases absent for a number of years; while the terms proposed by Vulpian and Erb refer only to the pathological anatomy of the disease. The old Hippocratic term "*tabes*," therefore appears to be the best, and is now again gradually displacing the others; while the adjective "*spinalis*" might be advantageously substituted for "*dorsalis*," which latter appears to refer to one portion of the spinal cord only.

An analogous affection of the lateral columns of the spinal cord has only recently been introduced into our system of nomenclature by Erb, who has called it *spastic spinal paralysis*, while Charcot has subsequently termed it *spasmodic tabes dorsalis*. Synonyms are *primary lateral sclerosis* (Berger), and *sclerosis of the pyramidal strands* (Dreschfeld). The existence of this affection as a separate pathological entity, however, is still very doubtful, for there is as yet not a single case on record in which primary sclerosis existed in the lateral columns of the cord without being complicated with disease of other important portions of that organ; and there is every reason to believe that the affection which is now generally designated as lateral sclerosis, will eventually have to be split up into a number of different maladies which, although having some points in common, yet diverge most widely in others. This point will be fully discussed later on.

No doubt exists about the occurrence of a combined affection of the white lateral columns and the grey anterior cornua, which was first distinguished by Charcot, and termed by him *amyotrophic lateral sclerosis*; the chief symptoms being motor paresis, and muscular rigidity followed by atrophy of the muscular fibre.

Another primary affection of this kind is multiple, insular, or disseminated sclerosis, which is also called *sclerosis in patches*, or *cerebro-spinal sclerosis*, and which invades chiefly the antero-lateral columns of the cord; but likewise the medulla oblongata, pons Varoli, the central white substance of the cerebellum and of the cerebral hemispheres. In this disease the morbid process is, as a rule, not so severe as in *tabes*; for while in the latter the nerve-tube eventually perishes altogether, that important portion of it which is known as the axis-cylinder appears to be habitually spared in insular sclerosis.

All forms of primary sclerosis are symmetrical, either from the beginning, or at least in the later stages of the disease.

Secondary sclerosis, on the other hand, is mostly unilateral, and appears as the consequence of some other primary disease, such as hemorrhage into the brain, softening of the cerebral substance from embolism or thrombosis of blood-vessels, or of hemorrhage into the substance of the cord, and acute myelitis. When it occurs after, de-

structive lesions of the motor area of the brain, it affects only that portion of the cord which corresponds physiologically to the diseased hemisphere, that is, the crossed pyramidal column.

Clinical observation, as well as morbid anatomy, seem to point to the conclusion that all these several diseases are closely allied; and that the differences in the symptoms which they present during life, are in the closest possible connection with the different physiological functions of the several portions of the cord which may be affected. It is a fact which stands out more clearly as our knowledge advances, that fibres which have certain functions in common are liable to become diseased together, at exceedingly different levels of the cord, while others in their immediate neighbourhood which have different functions are spared. Moreover, such fibres are liable to be affected symmetrically in the two lateral halves of the organ, the degree of the morbid change being often exactly parallel in both sides. The differences in the physiological function are naturally correlated to evolutionary, anatomical, and possibly also to chemical differences in these several parts. Indeed, the cord consists, like the brain, of a number of different areas or systems, each of which, although histologically identical with others, yet follows a special type of evolution in the fetus, has different connections with more peripheral and more central parts; may possibly be chemically quite distinct from its surroundings; and lastly, possesses peculiar pathological predispositions or proclivities, which are not shared by contiguous parts, however closely connected with it in anatomical position, and however similar in histological elements.

An apt illustration of these peculiarities is afforded by the manner in which different poisons act on different portions of the cord. Bread contaminated with ergot of rye will, when habitually taken for some time, cause well marked disease of the posterior columns; while bread containing an admixture of the *Lathyrus cicera*, which is eaten by the lower classes in certain parts of Italy, will lead to equally striking symptoms of disease of the lateral columns. Strychnia again has a special effect in exalting the excitability of the grey centre of the cord, and bromide of potassium in diminishing it. Finally, lead will, when absorbed for some time consecutively, cause granular disintegration of the large ganglionic cells of the grey anterior cornua, and thus lead to a peculiar form of muscular atrophy.

In order, therefore, to render the more minute pathology of the diseases under consideration thoroughly intelligible, it will be necessary for me to enter first into some explanation of the manner in which the highly complex structure of the spinal cord is built up and arranged. Histology and experimental physiology have done very little for the elucidation of this subject, while the study of evolution on the one hand, and of the finer pathological anatomy of the degenerations to which the organ is liable on the other hand, have been of the greatest service for unravelling the intimate structure of the cord. Either of these methods of inquiry may be used to control the results given by the other, thus affording a greater degree of reliability than each one singly could do.

With regard to evolution, Flechsig has shown that the central nerve-tube is at first laid down simply as a naked axis-cylinder, and that this latter is only subsequently furnished with a sheath of myeline. The latter is, therefore, a secondary formation, and is, in the several portions of the cord, developed at totally different periods of fetal life, just as well as the axis-cylinder itself. This mode of evolution follows a definite law with regard to the period of embryonic existence when it takes place; and it may thus be understood why tracts of fibres, which are, anatomically speaking, at a considerable distance from each other, are far more nearly allied than others which may be much nearer neighbours.

The succession in which the evolution of the different parts of the cord takes place is as follows:

1. The fundamental strands of the anterior columns (A c, Fig. 1).
2. The postero-external or Burdach's columns (B, Fig. 1).
3. The anterior mixed zone of the lateral columns (A L c, Fig. 1).

These three strands become developed at the earliest period of fetal life; the naked axis-cylinders being laid down at the end of the fourth week; while the myeline sheaths are fully formed at the end of the fifth month.

4. The lateral terminal layer of grey matter. No definite views have as yet been advanced concerning the period of evolution of this portion of the cord.

5. The postero-internal or Goll's columns (c) are laid down at the end of the second month, and the formation of their myeline sheaths appear to be finished at the end of the sixth month, giving again a period of four months for the entire process.

6. The direct cerebellar strands (c, Fig. 2), which establish a connection between the grey matter of the cord and cerebellar centres, are

first observed at the commencement of the third month, and finished at the beginning of the seventh.

7. The crossed pyramidal strands (CP) are the latest of all, being laid down about the middle of the fifth month, while their myeline sheath is finished at the end of the ninth. They establish a connection between the grey matter of the cord, on the one hand, and the central ganglia and convolutions of the hemispheres, on the other hand.

From the foregoing description, it appears that the different conducting paths in the spinal cord are laid down and formed in an absolutely systematic manner; and that, at the same period of fetal life, only

FIG. 1.—Section of the spinal cord from the region of the fifth cervical nerve, from a fetus 12 centimètres long (after Flechsig). P = anterior pyramidal column; AC = fundamental tracts of the anterior columns; ALC = anterior mixed lateral column; PC = processus reticulares; CP = crossed pyramidal columns; B = Burdach's columns; G = Goll's columns.

one, or several allied systems are formed. Those which are most important for the life of the fetus, and which are reflexory in nature, are early developed, their faculty of incessant function being an indispensable condition for viability altogether. On the other hand, those strands which place the spinal cord under the influence of the psychomotor centres, or the will, become evolved at a later period. These latter are more abundantly developed in man than in the lower animals, and do not become formed at all in congenital absence of the brain.

Anomalies of evolution have unquestionably an influence in the causation of diseases which occur later in life, and may eventually be found to be the chief material base of what is now called the "neurotic constitution." Congenital fissures or cavities may become the starting-point of glioma, glio-sarcoma, and chronic interstitial degeneration. Anomalies in the distribution of the white and grey matter are not

FIG. 2.—Section of the spinal cord from the middle portion of the cervical enlargement, of a fetus 35 centimètres long (after Flechsig). P1 = anterior pyramidal strand; AR = anterior roots; C = direct cerebellar columns; CP = crossed pyramidal columns; B = Burdach's columns; G = Goll's columns. Magnifying power, 6.4 lin.

rarely met with in persons who have died of certain diseases of the nervous system, such as general paralysis of the insane. Kahler and Pick found, in a person who had died of tabes, abnormal smallness and irregular shape of the central grey matter, and extreme smallness of the posterior columns, which was not wholly due to atrophy. As there are "born poets" and "born criminals," so there are born sclerosers. The gradual formation of the myeline sheath of the central nerve-tube may, in some cases, be carried out in a deficient manner; and such a condition may, later on, lead to actual disease of imperfectly formed or protected parts. Where undue claims are made upon a small or badly built medulla oblongata or spinal cord, there will be greater tendency to the occurrence of asthma, diabetes, tabes, and paralysis, than there

would be in cases where these organs have been more perfectly developed.

In tabes spinalis, more or less striking morbid changes are discovered after death, not only in the posterior columns, but also in the spinal pia mater; the posterior nerve-roots; some parts of the central grey matter, more especially Clarke's vesicular columns; the direct cerebellar strand of the lateral column; the sympathetic system of nerves; almost all cranial nerves; and the terminations of the spinal nerves in the skin, the joints, and the muscles. The question therefore arises, which of these numerous alterations is the primary and essential one, and what is the exact pathological nature of the change?

Posterior spinal lepto-meningitis is an habitual concomitant of tabes, and Arndt, Waldmann, and others consider it the primary event, and the actual cause of the wasting of the posterior columns, this being, according to them, a secondary degeneration. This opinion, however, is contradicted by several well observed cases of tabes in its initial stage, in which no meningitis existed (Strümpell, Tuzek). As tabes has in general no tendency to shorten life, cases in the first stage of the disease come only rarely on the *post mortem* table, and then mostly in consequence of complications, such as typhoid fever, etc. It may now be looked upon as settled, that meningitis is not the primary event, but only occurs in the later stages, when there are not simply slight changes in the structure of the central nerve-tubes, but when gross alterations in the size and consistency of the posterior columns have already become perceptible to the naked eye. The thickening seen in the pia mater is most likely owing to mechanical conditions, as it helps to fill up the vacuum which is produced by the shrinking of the posterior columns. The statement that the peculiar lightning-pains of tabes are caused by inflammation of the pia mater is certainly wrong, as these have been observed in certain cases where, eventually, no alteration of the membrane was discovered.

Another view which has been put forward is, that atrophy of the posterior nerve-roots is the initial lesion in tabes, and that the disease creeps from them upwards into the substance of the cord itself, like an ascending neuritis. It is quite true that, at a somewhat advanced stage of the malady, the posterior roots are habitually found wasted; but the roots never suffer to the exclusion of the columns; and, in early cases, the roots have been found healthy when there was already unmistakable disease in the columns. This latter fact deals the death-blow to the posterior root-theory.

Déjerine, to whom we owe some useful observations on the atrophy of peripheral nerves which is seen sometimes to accompany tabes, has recently started the extraordinary notion that tabes is really a peripheral affection, and should be called *neuro-tabes périphérique*. This he concludes from two cases in which he found signs of old neuritis, while the posterior columns were healthy. This simply shows that the cases were not such of tabes, but of neuritis. Yet, such is the feverish desire for novelty nowadays, that Déjerine's most immature idea has already been hailed in the medical journals as a wonderful discovery, and one likely to lead to a system of treatment widely different from that based on the supposed lesion of the cord.

Another original theory is that recently brought forward by Page, who sees the starting-point of this formidable malady in a corn! Here we seem to have arrived at the infinitely little in order to account for a great deal. The prevention of tabes would, according to this theory, rest with the chiropodists. According to Page, the disease may begin with the painful pressure of corns, and the protracted sensory disturbance may give rise to ultimate structural change, first in the peripheral nerves, and then travelling upwards to the cord. It appears to us that the discomfort which the majority of Europeans habitually experience from corns is bad enough, without crediting these little pests with such wickedness as eventually to conduce to tabes. That corns, especially in certain positions, may be often the beginning of considerable trophic changes in a limb, seems probable enough. But, in the case which forms the basis of Page's theory, an important link is wanting, inasmuch as it has not been shown that the patient was free from spinal disease at the time when the corn commenced to give trouble. The sequence of events is much more likely to be as follows: tabes, from whatever cause, is the primary change; this induces liability to peripheral nerve-disturbance, so that certain exciting causes, such as long-continued pressure, violence, or injury, are more likely to lead to mischief than in healthy persons, whose nerve-power is not defective. In this respect, the recent observations of Pitres and Vaillard are highly suggestive, as it has been shown that there is tendency to peripheral neuritis, not only in tabic subjects, but also after cerebral hæmorrhage, small-pox, alcoholism, and other distempers giving rise to a state of low vitality.

It cannot be a matter of surprise that that universal scapegoat, the sympathetic system of nerves, should have been accused of being at the

bottom of the complaint. Most of the older observations on disease of the sympathetic ganglia and fibres in tabes, must be received with a considerable degree of caution; but the recent researches of Raymond and Arthaud, which were made in Vulpian's laboratory, render it evident that, at least occasionally, atrophy of the ganglionic cells and fatty degeneration of the nuclei of Remak's fibres; together with corresponding changes in the blood-vessels and connective tissue, do occur. But to conclude from this that the sympathetic is primarily affected, and that all other alterations spring from that source, constitutes a *salto mortale* which we at least consider it somewhat too hazardous to undertake. We will not here allude to the fact that the more intimate structure of the sympathetic is by no means invariable in health, but that great variations occur habitually in the size and pigmentation of the ganglionic cells, and in the proportion of nerve-tubes furnished with a medullary sheath, and Remak's fibres. Suffice it to say that, even supposing that there were a constant change in the sympathetic, it would even then be impossible to explain why this should lead, as a rule, only to affection of the posterior, and not also of the antero-lateral columns, and other portions of the spinal cord.

When Duchenne was under the impression that he had discovered a new disease, "locomotor ataxy," he considered the cerebellum to be the most probable seat of the malady; and more recently Neftel, of New York, has asserted that tabes is really an affection of the brain. But what does pathology teach us? We know that the morbid change in tabes affects the cord from top to bottom, but that, in the vast majority of cases, it stops short in the medulla oblongata. Indeed, changes in the cortex and in other portions of the brain, have only been found where towards the end tabes had become complicated with epilepsy, general paralysis of the insane, and similar conditions. It is true that we meet occasionally, in the initial stage of tabes, with such symptoms as aphasia, apoplexy, hemiplegia, mental derangement, etc.; but all such symptoms are fleeting and evanescent, and can be differently explained. The theory of the cerebral origin of the disease, therefore, seems to rest only on certain somewhat imperfectly understood clinical observations, while the pathological evidence of innumerable well observed cases is directly antagonistic to it.

The process of exclusion, as well as of direct observation, teaches us that tabes is primarily a disease of the spinal cord, and more especially of its posterior columns.

What portion of the posterior columns is first affected? In cases of long standing, their entire transverse section, including both Burdach's and Goll's strands, is sclerosed, while in the earlier stages of

FIG. 3.—Schematic diagram of the different areas of degeneration in the posterior columns (after Strümpell). 1. Small portion of diseased tissue at either side of the posterior commissure. 2. Bulk of Goll's columns. 3. Lateral area of Burdach's columns. 4. Small round fold near apex of Goll's columns. 5. Postero-external fold. 6. Small strand at inner side of posterior roots.

the disease the process appears to be more strictly localised in certain areas. Pierret, who was the first to examine this point more minutely, concluded from his observations that the disease commenced with two symmetrical islands of degeneration in the posterior root-zones, or Burdach's columns, and that Goll's strands only became subsequently and secondarily affected.

This view was endorsed by Charcot and Vulpian, and was for some time unreservedly accepted by the profession; but more recent researches by German anatomists, more especially by Strümpell of Leipzig, have shown that such is not by any means invariably the case, but that, at least in a certain number of instances, the most internal portion of Goll's columns becomes sclerosed in the very commencement of the disease, while the postero-external field of Burdach's columns remains normal for a long time. Strümpell's researches appear to us to constitute a decided step in advance in our knowledge of the finer anatomy of the disease, and show us how much still remains to be done in this department. Want of time prevents us from enter-

ing fully into this matter, as we have yet to discuss the important question, in which tissue the malady actually begins. Is the fault primarily located in the blood-vessels, in the intertubular connective tissue, or in the nerve-tubes themselves?

Ordoñez, who has described a lesion of the arterioles of the posterior columns, has broached the view that certain changes in the coats of these vessels must interfere with the exchange of nutritive material between them and the tissues, and thus lead to atrophy and sclerosis of the nervous structures. This is, however, controverted by the circumstance that the vascular lesions are variable, and may be absent, while the changes in the nerve-tubes are constant. Moreover, the phenomena of Wallerian degeneration show that alterations in the blood-vessels are rather the consequence than the cause of the changes which take place in the nervous structures.

Interstitial degeneration of the neuroglia is believed by Adamkiewicz and some other observers, to be the essential lesion of tabes; but if we were to accept this theory, we should find it impossible to understand why a disease which primarily affected the neuroglia, should be so strictly localised in certain systems of the cord as tabes is known to be, while the anatomical distribution of the connective tissue is exactly the same in all the different strands of the cord.

It cannot then be doubted any longer that the primary change which occurs in tabes, takes place in the central nerve-tube, and more especially in that most important constituent of it, the axis-cylinder. Irritation of this structure gives rise to those singular attacks of lightning-pains, which form the most striking symptoms of the first stage of the disease. Repeated attacks of irritation lead to imperfect nutrition, and finally to atrophy. The dead axis-cylinder then acts as a foreign body, and causes further irritation in the neighbourhood. The nuclei of the nerve-fibres swell and multiply; the protoplasm surrounding the nuclei begins to proliferate; the myeline sheath becomes segmented, and ultimately reduced to small drops and granulations; and the intertubular connective tissue and the blood-vessels now begin to participate in the disease, which is thus shown to be one of parenchymatous degeneration or atrophy, rather than one of inflammation or interstitial degeneration.

An affection analogous to tabes, but affecting the lateral instead of the posterior columns, has been described as spastic spinal paralysis, or spasmodic tabes dorsalis, or primary lateral sclerosis, and has, during the last few years, excited a good deal of discussion. I am bound to say that it has as yet eluded the grasp of the pathologist.¹ Dr. Ross, of Manchester, in the second edition of his *Treatise on Diseases of the Nervous System*, credits his colleagues, Professors Morgan and Dreschfeld of that city, with the honour of having been the first to prove by dissection the connection of the symptoms of spastic paralysis with lateral sclerosis, in a primary and uncomplicated case of the disease. The account, however, which is given by those observers in the *Transactions of the International Medical Congress*, and more particularly in the *Journal of Anatomy*, shows quite clearly that their case was one of the amyotrophic variety of the disease, inasmuch as the giant-cells of the grey anterior cornua were likewise affected, there being simple atrophy, pigmentary atrophy, and occasionally complete absence of whole groups of these cells. All the other cases which have, until now, been recorded, are equally unsatisfactory, for one reason or another.

Leyden has long ago expressed his disbelief in Erb's spastic paralysis as a distinct disease, and has, in cases which showed the symptoms described by Erb during life, generally found myelitis of the dorsal portion of the cord after death.

Apart, however, from this, the phenomena of spastic paralysis may be owing, more particularly in children, to disease or deficiency in the motor area of the cortex (porencephaly), and it is, in such cases, sometimes not easy to decide whether the case is one of spinal or cerebral disease. Finally, there may be all the symptoms of spastic paralysis present in young girls, and lead to the diagnosis of lateral sclerosis, when sudden or gradual recovery, with or without treatment, shows the case to have been of a functional, or, as is then generally said, of a hysterical character.

There appears to be, therefore, no primary symmetrical sclerosis of the lateral columns of the spinal cord, analogous to the posterior sclerosis of tabes. Nevertheless, we shall be justified for the present in retaining the clinical expression, spastic paralysis, and in keeping in mind its frequent, if not invariable, connection with either structural disease or functional irritation of the lateral columns. An exclusive affection of these latter may possibly exist during some period of the malady, but will rarely be discovered *post mortem*, which corresponds

¹ Since this lecture was delivered, an uncomplicated case of this disease has been described by Nimkowski.

clinically with the fact that the symptoms of motor irritation and paralysis are, in the later stages of the disease, almost invariably complicated with sensory and trophic disturbances, such as anesthesia, paralysis of the bladder, bed-sores, muscular atrophy, etc.

REMARKS

ON A

CASE OF PHLEGMONOUS GASTRITIS.¹

By THOMAS WHIPHAM, M.B.,

Physician to and Lecturer on Pathology at St. George's Hospital.

MARTHA M., aged 49, was admitted into St. George's Hospital under my care on July 4th, 1883. It appeared from the history that she had been for several years subject to bilious attacks, in which severe vomiting with occasional diarrhoea were prominent symptoms. She had lost all her teeth, and was accustomed to drink large quantities of warm liquids. The diarrhoea had been very troublesome since the beginning of 1883; and on April 24th of that year she was an in-patient under Dr. Wadham, in a state of great prostration, after prolonged diarrhoea. There was on this occasion no vomiting. With rest and light diet (fish, etc.), the symptoms improved, and she was discharged relieved in a fortnight.

About a month after she had left the hospital, the diarrhoea returned; and on July 1st an old femoral hernia came down, causing her much abdominal pain, more especially in the epigastrium. When in the recumbent position, the hernia disappeared; but this was at first followed by profuse and continuous vomiting; at first of dark matter, then of thin bilious fluid. The bowels were freely open.

She was admitted into hospital at the request of Mr. Chippendale, of Upper Phillimore Gardens, who had been called in to attend her. At first, she was placed in a surgical ward, as it was supposed that the symptoms might depend upon the hernia. No rupture being discovered, she was at once transferred to me.

The patient was badly nourished and slightly jaundiced. She had lost all her teeth. She complained of abdominal pain, referred chiefly to the epigastrium. No tumour was detected. The abdominal walls were flaccid. The vomiting was almost continuous; the vomited matter was small in quantity, fluid, opaque, green, and strongly acid.

Small doses of acetate of morphia with sulphate of atropia were given to relieve the pain; while at the same time, with a view of maintaining the patient's strength, nutrient enemata containing brandy were administered. The injections were retained fairly well; but the vomiting did not entirely cease, nor did the character of the vomited matter alter.

On July 6th, the patient sank into a comatose state; the temperature gradually rose from normal to 104° Fahr., with the symptoms of peritonitis; and she died soon after midnight.

Necropsy, twelve hours after death.—*General Appearances.* Height, 5 feet; black hair; no teeth: jaw atrophied. *Lungs.* Posterior portions of both congested. *Heart.* Weight, 12 oz. The mitral valve was thickened and contracted; only one finger could be passed between its flaps. The remaining valves and orifices were natural. *Abdomen.* The peritoneal cavity contained a small quantity of purulent fluid. *Stomach.* The contents were thin greenish yellow fluid. The mucous surface at the cardiac third of the organ was very red, felt thick, and infiltrated with fluid. On transverse section, it was a quarter of an inch thick. The remaining two-thirds of the stomach, so far as the inner surface was concerned, presented no abnormal appearance; but, on transverse section, the pyloric end was found infiltrated with a purulent effusion between the muscular and mucous coats. On the posterior surface of the organ, below the lesser curvature, there was an area about one inch wide by three inches long, which was of a distinctly yellow colour; and, at this place, section showed an abundance of purulent fluid between the muscular and serous coats. The remainder of the alimentary canal was natural. *Liver.* Weighted

3 lbs. 12 oz.; its surface was smooth. The tissue was pale and fatty. *Spleen.* Healthy; weight, 7 oz. *Kidneys.* Weighted 8 oz. The capsules were adherent; the surfaces were granular; the cortical parts diminished, and mottled slightly with dark coloured spots. Microscopic examination confirmed the naked-eye appearances of the walls of the stomach. There was an extensive infiltration of the submucous tissue, chiefly by pus-cells. The pus was most abundant around the secreting glands, the cells of which were somewhat swollen, and their contents opaque and granular. At that part of the posterior surface of the organ where the yellow discoloration of the external surface was found, as described above, all the coats of the stomach were infiltrated with pus.

After the patient's death, the friends brought out a box labelled "podophyllin pills." They stated that she had taken one or two of them on the night before admission; but they were unable to say whether or not she was in the habit of taking them.

REMARKS.—Cases of acute inflammation of the walls of the stomach have received but little attention in the ordinary English text-books. In foreign works, however, more attention has been paid to this affection. Raynaud, in the *Bulletins de la Société Anatomique*, 1860; and Leube, in Ziemssen's *Cyclopadia*, vol. vii, devotes a considerable space to the disease—gastritis phlegmonosa. According to the latter, it occurs in two forms: 1. circumscribed suppuration; 2. diffuse purulent infiltration of the parietes, extending around and between the gastric glands. In the circumscribed form, the mucous as well as the serous layers are usually intact, or at most only erythematous. "The abscesses of the stomach are sometimes single, at other times multiple." "Cases of diffuse purulent infiltration of the parietes have been observed less frequently, and less exactly." In this latter, the condition of the mucous layer varies greatly in the different cases examined; it is sometimes unchanged, at other times infiltrated with pus or serum. Occasionally the muscular layer may be destroyed by the purulent infiltration.

The serous layer may be also unaffected, according to the seat and extent of the phlegmonous inflammation. In other cases, it may take on inflammation, accompanied by the accumulation of dirty grey thick masses of exudation upon the surface of the stomach. In these cases, the subserous layer may be infiltrated with serum or pus.

Of thirty-one cases collected by Leube, twenty-six occurred in males and five in females (Ziemssen's *Cyclopad.*, vol. vii, p. 157.) Bamberger (*Handbuch der Speciel. Pathol. und Therapie: Krankheiten des Chylpoëtisch. Syst.*) devotes but a single short paragraph to this affection of the stomach; his description of the morbid appearances agrees with that of Leube.

Etiology.—It appears, from the records of the cases which have been observed, that the cause is unknown. According to Leube: "In a number of cases, the origin of the affection has been attributed to excessive indulgence in alcohol; in a few, to some wound occurring in the region of the stomach; in others, again, to some error of diet." An obvious objection occurs at once to such explanations of the cause of phlegmonous gastritis—viz., that in countries such as England, to say nothing of other nations, where drinking is carried on to so great excess, and in which errors in diet are an equally prevalent fault, instances of this disease are of rare occurrence. I purposely omit those cases in which inflammation of the walls of the stomach has followed acute diseases, such as typhoid fever, puerperal fever, etc. It is within the knowledge of all medical practitioners that, in many persons, certain articles of diet exercise a deleterious effect—e.g., in some, lobster or other shell-fish invariably produce an attack of urticaria. In my own experience, a few strawberries always have this effect in a boy about ten years of age. Podophyllin is differently tolerated by different individuals. In some, its action is most beneficial; in others, it has the reverse effect. In a patient of my own, a quarter of a grain is sufficient to render her so irritable that, as she expressed it on one occasion on which I prescribed it, she was "unable to sit still all day;" while the idiosyncrasy which renders many persons intolerant even of a few grains of potassium iodide, is too well recognised to require comment.

It may be that, in certain persons, an attack of phlegmonous gastritis may result from the ingestion of articles of diet usually considered wholesome, or from the administration of drugs in ordinary use; hitherto, however, its occurrence under any circumstances is a rare event.

We have no evidence that the patient whose case forms the subject of the present communication had ever taken podophyllin before; and with such deficient information no conclusion can be drawn in this case, one way or the other.

Leube states that, "as regards age, those who died of gastritis phlegmonosa were, almost without exception, in middle life." The age (49) of the patient in the present instance is a case in point.

¹ Read before the Medical Society of London.