myself. The bulbous end has a mean diameter of No. 5; the stem is about 2½ or 3. The patient being under the influence of an anæsthetic, the little catheter which has been tied in is withdrawn, and the urethrotome is first introduced as far as to the deep-seated stricture, through which the terminal bulb is then insinuated. It must now be passed fully half or three-quarters of an inch further in, that is, beyond the stricture; and the blade being exposed in a direction towards the floor, is pressed firmly thereon and drawn forward, until all resistance, sometimes considerable, is completely overcome. A touch on the button near the handle sheaths the blade, and the outward movement proceeds until the site of the second stricture is reached, when another incision is made in the same manner as before. The urethrotome is then withdrawn, and the meatus freely divided by a scalpel or short urethrotome. I next take a blunt metal bougie or dilator, No. 15 or 16, English size, and ascertain if it will pass without obstruction into the bladder. It generally does so at once; if, however, its progress be arrested at any point, the situation of this is carefully noted, when I withdraw the dilator, reintroduce the original urethrotome, and divide the opposing tissue; but this, as I before said, is seldom necessary. I may add that I am very rarely satisfied with anything less than the free and easy passage after the operation of a metallic sound, No. 16 in size; in some cases, No. 17 or even 18 will pass. A gum-elastic catheter, No. 12 or 13, is then tied in, and should remain always 48 hours; with an extra 24 or 48 hours, if the incisions have been deeper than usual, or if hæmorrhage be free or continuous, the latter being a very exceptional occurrence.

It is sometimes objected that, if a small instrument can be easily passed through the stricture, and if this may be easily dilated so as to admit an urethrotome as large as No. 5, why should any cutting operation be performed? The reply need only be brief; the object of division is to remedy the tendency to recontract, which certain strictures exhibit, and which almost all do when they have continued several years. In other words, it is marked contractility in a stricture, and not mere narrowness, which renders operation by incision desirable.

An important inquiry remains: What are the results of internal urethrotomy in relation to the reappearance of stricture? Does the operation free the subject of it from that liability to return which constitutes the very serious character of the disease when treated only by dilatation?

The reply to this question cannot be a simple and categorical one; it must be made with limitations.

I may remark first, that it may be taken for granted that no mode of urethrotomy which fails to ensure complete division of the obstructing tissues has a chance of affording permanent immunity from the complaint. Whatever be the mode of operating, if incomplete division only be effected, the patient will find his urethra narrowing after a more or less prolonged interval of time. What happens then in those cases in which, as far as the operator can judge, he has effected a complete division? Well, I am free to confess that it is not possible to promise immunity from return. We may render the period of return as remote; we may produce a good calibre, easily maintained by occasional regular dilatation, a procedure which, before the operation, was not only ineffective, but painful and irritating. We may place a patient in a condition of health and comfort for several years, meantime saving his bladder, ureters, and kidney from the slowly but surely occurring grave changes which threaten his existence. Now and then, but rarely, I have met with a case in which the patient's troubles have never re-appeared. I am certain we cannot reckon on this result as a rule, and I mistrust claims to the contrary as due to error of judgment, or want of information, or as prematurely made, *i.e.*, without waiting sufficiently long to observe remote results. In enunciating this opinion, I have no hesitation whatever. But herein I have stated the worst in regard to the patient's prospects, and have shown you the reverse of the medal. Thus, supposing after a few years the patient finds himself unable, through re-appearance of contraction, to pass a bougie of adequate size, division can again be resorted to. It is not a dangerous proceeding, necessarily occasioning hesitation on the part of the patient, when his condition demands relief. Just as in calculus cases, a second, a third, or even a fourth stone can be, and often is, safely removed by lithotripsy when, after the lapse of time, fresh products are formed; thus also may a stricture be dealt with by this operation, a second or a third time if necessary. By the prompt resort to the remedy, should it be again required, the disease may be confined to the canal, and the real gravamen of the malady, the implication of vital organs, certain to result after long continued ineffective dilatation, is altogether avoided.

I have performed this operation now for between three and four hundred patients. Some of my earliest cases were, from want of suffi-

cient confidence and experience, less completely and freely cut than those on whom I have operated of late years, and a few of the early patients have been recut. I have not regretted such advice in a single instance, for the advantage to them has been undoubted.

The risk of the operation is very small. Estimating the number of patients on whom I have performed it as 340, which is within five, more or less, the deaths have been six, or not two per cent. These were due to pyæmia in three, to embolism in one, to extravasation and exhaustion in two; one of the latter was a case in hospital almost thirty years ago, among my earliest, and he was unfit for any operation.

The sum of my experience is the expression of a strong conviction that internal urethrotomy, completely performed, should be resorted to as the best and safest treatment of stricture, as soon as the easy use of the bougie fails to maintain the urethra patent, or to allay signs of irritation in the bladder arising from obstructed urethra. It is the best means not only for relieving urethral obstruction and its painful symptoms, but for ensuring the future sound condition of the organs which lie behind it.

A CLINICAL LECTURE ON TROPICAL LIVER-ABSCESS.

Delivered at Charing Cross Hospital, May, 1884.

BY SIR JOSEPH FAYRER, K.C.S.I., M.D., F.R.S.,

Honorary Physician to H.M. the Queen and to H.R.H. the Prince of Wales;
Physician to H.R.H. the Duke of Edinburgh; Physician to the
Secretary of State for India in Council; etc.

GENTLEMEN,—You will remember that, when recently directing your attention to the case of J. S., who was suffering from tropical diarrhoea, I pointed out to you certain complications which seemed to remove it from the category of ordinary examples of this disease, and to render the prognosis more doubtful than it might otherwise have been. The symptoms referred to were a continuous daily rise of evening temperature to 100° or 101°; a tendency to sweat; a dry, glazed, and fissured tongue; anxious expression of countenance; a sense of fulness, sometimes of pain or uneasiness, on pressure over the liver; some tension of the recti muscles; with defective respiration, signs of consolidation of the base of the right lung, and cough. At the same time, the diarrhoea from which he had been suffering had ceased, and the dejections had become formed though copious, once a day, as is the case when much milk is taken. On the 7th instant, I found that symptoms of hepatic mischief had become more pronounced, and there was an appearance of bulging of the right side, which was not, however, confirmed by measurement.

His condition had become one of grave anxiety. At my request, therefore, Mr. Barwell introduced an aspirating trocar and cannula between the eighth and ninth ribs, in a line between the axilla and nipple; and, as I anticipated, pus escaped through the tube. The opening was at once enlarged by a free incision, and about thirty ounces of thick dark coloured pus, mingled with blood, escaped. A large drainage-tube was placed in the wound, the strictest antiseptic precautions being observed throughout the proceeding. The patient experienced great relief, and there was a rapid fall of temperature, which has remained normal since the operation. The look of anxiety and suffering, and the pain in the hypochondrium, have passed away. The sweats have diminished. He eats and sleeps well.

The case is one of considerable interest, and so instructive in its character that I have thought it well to invite your attention to it, and at the same time make it the text for some remarks on the subject of tropical liver-abscess generally. I speak of this disease as "tropical," because it most frequently occurs in those who live in hot and tropical or subtropical countries, or who have resided there for some time previously, as was the case with our patient, J. S. Of course, the liver may suppurate from certain causes anywhere; but the form of liver-abscess I am now considering is essentially a disease peculiar, if not confined, to hot climates; or rather, I should say, the conditions which give rise to liver-abscess are prone to occur in those countries; the suppuration itself may appear after return to a temperate climate, as in S.'s case.

Hepatic suppuration is not limited to the tropics, it may occur anywhere; but, in this country, it is comparatively rare, and is generally due to septic absorption from some wound, operation, or ulcer of the stomach, gall-bladder, or its duct; or to suppuration in the pelvic cavity or about the hæmorrhoidal region. Such cases are pyæmic, and

very different from the large single abscesses found in tropical residents, but similar to certain forms which occur in connection with dysentery, and are due either to direct septic absorption through the veins, or to general septic poisoning. In the latter cases, the abscesses are apt to be numerous, and the liver may be studded with suppurating patches, where the pus has formed round portions of necrosed liver-tissue; these are most dangerous, perhaps generally fatal.

Let me briefly describe to you the different forms in which liver-abscess presents itself; but you must not expect that you will be able always to distinguish them easily; for, like other things classified, the forms may seem to run into each other, though typical examples are tolerably distinct.

First, then, the large single, or generally single, abscess is referable to heat, malaria, the effects of a tropical climate combined with altered conditions of living, food, drink; the excessive or injudicious use of alcoholic beverages; though you must not assume that, in all cases, intemperance is a cause. Tropical abscess may and does occur even in the most temperate, but it is more frequent in those who live freely. This abscess is often large and solitary (quite recently I removed eighty ounces and, again, fifty ounces, from the same liver within a week); there may be more than one abscess. The formation is generally preceded, especially in the chronic, by fever, pain, and all the signs of active inflammation. Sometimes it assumes a more chronic form, and is insidious in its course, presenting no well defined symptoms, or so slightly marked as to escape observation until the existence of an abscess is revealed by bulging of the thorax, pointing in the epigastrium or below the ribs, or by sudden evacuations of the contents through the stomach, lungs, or bowels, or into the peritoneum, or it may be not until examination after death from some other disease. I have frequently detected liver-abscess where the symptoms were very obscure, indeed absent; and where it was not suspected. As a general rule, the prognosis of liver-abscess of the simple form, if it be recognised and properly dealt with early, is not unfavourable, though one must always feel uncertainty; but if early evacuation of its contents be effected, recovery is hopeful. The evacuations may take place spontaneously through any of the channels before mentioned; but if you are able to detect pus, and can remove it by puncture, the chances of recovery are enhanced.

Secondly, liver-abscess, following or co-existing with dysentery and tropical diarrhoea, was regarded by Murchison and others as pyemic; and due to absorption of pus or septic matter from the ulcerated bowel, and therefore similar to the septic liver-abscess of temperate climates. That this is the case in some instances is indisputable; and when multiple liver-abscesses occur in a person who is or has recently been suffering from dysentery or chronic diarrhoea, complicated with ulcerated bowels, or fever with Peyerian ulceration, no better explanation of its origin could be given; such cases are very dangerous. Let me ask you to note the nature of the abscesses in cases of this truly pyemic character. They begin as patches of necrosed tissue, embolisms, or infarctions; which, if the patient live long enough, as generally happens, become abscesses from suppuration occurring around the dead tissue, as in the case of the core of a boil. This becomes the centre of a cavity, varying in size from a pea to an orange, containing debris of tissue, leucocytes, and finally pus; the whole being surrounded by an area of congested, and sometimes pus-infiltrated, liver-tissue. Such abscesses as these occur also in the other viscera, spleen, and kidneys; they are evidences of general septicæmia.

Liver-abscess, however, may be unassociated with dysentery or diarrhoea, and yet be multiple; on the other hand, it may accompany or follow dysentery or diarrhoea, and still be single. It is often difficult to come to any decided opinion as to the true character; each case requires careful study, and the etiological relation of dysentery with liver-abscess is always a matter of considerable interest. That liver-abscess and dysentery are dependent on each other, has been held by high authority. Annesley considered that dysentery was due to hepatitis. Budd, thirty-six or thirty-seven years ago, enunciated the opinion which has received more support here than in India—that liver-abscess is always due to purulent absorption from the bowel; that, in fact, tropical abscess is pathologically identical with liver-abscess in this country. Abercrombie thought they were co-existent but independent of each other. Murchison, Macpherson, and others concur in this. Sir R. Martin and others thought that liver-abscess was intimately connected with disease of the cæcum; but this was not confirmed by post mortem examination, for, of seventy-two cases where the cæcum was affected, in only twenty-two was there liver-abscess. Moore says that eight observers out of twelve accord a percentage of 18, or 25 cases of liver-abscess in 1,522 of dysentery; four other observers give a ratio of 39 per cent., or 62 out of 151 cases of dysen-

tery; 18 per cent. is more probably the correct ratio, the larger percentage being more probably due to all diseased conditions of the liver that have accompanied dysentery. Recent observations, I think, confirm this view of the relative frequency of hepatic abscess in connection with dysentery; and albeit in some cases of multiple abscess the condition is due to septic absorption from a dysenteric bowel, yet generally it is to be regarded rather as an expression of the general disease than as a direct consequence of dysentery. It would appear that, in different hot climates, it is more or less frequent according to circumstances, but that it is more prevalent in India than in any other country.

As to the extent to which hepatitis prevails in India, a late statistical report gives the following information. "In the European army of India, averaging 57,742 men, 28,780 admissions and 1,267 deaths were recorded under hepatitis during the ten-year period 1870 to 1879. This gives an admission-rate of 49.8, and a death-rate of 2.19 per 1,000. In the army of Bengal, the admission-rate was 47.1, and the death-rate 2.04 per 1,000. In the preceding ten-year period, the admission-rate was 59.2, and the death-rate 3.31 per 1,000. A considerable reduction has thus taken place in both the admission and the death-rate, and the diminution has been pretty evenly distributed over the different sections of the Presidency. There were 7,417 admissions and 349 deaths from hepatitis in the European army of Madras during the years 1870 to 1879, an admission-rate of 67.2, and a death-rate of 3.31 per 1,000; much higher rates than those for Bengal. During the immediately preceding ten-year period, the admission-rate from hepatitis was 70.2, so that very little diminution had taken place in the frequency of this disease. During the period 1870 to 1879, there were 4,234 admissions and 177 deaths from hepatitis in the European army of the Bombay Presidency, the admission-rate being 40.9, and the death-rate 1.71 per 1,000, slightly less than the corresponding rate for Bengal. For the years 1863-69, the average admission-rate from this disease in Bombay was 26.6 per 1,000. Hepatitis thus seems to have been more prevalent of late in the Bombay Presidency. The eighteenth annual report of the Sanitary Commissioner for 1881 tells us that the admissions for hepatitis per 1,000 men for that year were: Bengal 32, Madras 39, Bombay 29; the deaths per 1,000 in the same period were: Bengal 1.64, Madras 1.54, Bombay 1.52, from the same cause. It further remarks that, "of the chief diseases, hepatitis was the only one which, during 1881, yielded a higher death-rate than the preceding year, 1.60 per 1,000 against 1.34. The increase was, however, very trifling, and the admission-rates are nearly identical. In Madras, hepatic diseases in former years have prevailed to a greater extent than in the other Presidencies. The returns during the last two years have been very favourable compared with the preceding ten-year period." Also, "after anæmia and general debility, the chief causes of invaliding in the army of India were: hepatitis, 2.96 per 1,000; dysentery and diarrhoea, 2.83; phthisis, 2.21." As a great proportion of the deaths recorded under hepatitis were probably due to liver-abscess, it will be seen how large the experience of the disease is in India.

With reference to the relative proportion in which liver-abscess and dysentery were coincident or combined, Dr. Waring, in 1854, made the following notes:

"Of 300 cases of fatal liver-abscess in India, in only 82 cases, or in 27.3 per cent., was the hepatitis preceded by symptoms of dysentery. Of the 600 cases, the number of abscesses was not stated in 12. Of the remaining 288, 177 had one abscess, 39 had two abscesses, 11 had three, 17 had four, 10 had from five to ten or more, and 40 had more than ten."

Relative Frequency of Hepatitis in different Foreign Stations.¹

Madras.					
Year.	Strength.	Disease.	Admitted.	Died.	Invalided.
1870	9,936	hepatitis	685	41	131
1871	10,684	"	713	41	58
1872	10,892	"	586	5	70
—	—	abscess	43	35	—
China.					
1870	546	hepatitis	15	—	1
1871	516	"	8	0	0
1872	789	"	15	1	0
—	—	abscess	1	1	—
Cape, and St. Helena.					
1870	2,441	hepatitis	66	4	16
1871	2,473	"	24	0	6
1872	2,484	"	23	1	3
West Indies.					
1870	1,482	hepatitis	2	0	0
1871	966	"	12	0	0
1872	1,112	"	2	—	2
—	—	abscess	1	0	0

¹ Fayrer, Tropical Diseases.

Canada.					
1870	4,519	hepatitis	10	0	0
1871	2,388	"	12	1	1
1872	1,602	"	3	0	9
Malta.					
1870	4,799	hepatitis	26	5	6
1871	4,977	"	26	2	7
1872	4,855	"	27	0	4
		abscess	2	1	0

Morehead says that, in malarious countries, liver-abscess frequently occurs in association with remittent fever without any ulceration of the bowel. The combination of malarial poisoning with hepatic mischief is most frequent, and it is often difficult to decide how far certain rigors and chills may be due to malarial fever, how much to incipient suppuration in the liver, or to what extent both may be concerned. Of course, when there are physical signs of enlargement, pain, bulging or pointing, there is no difficulty in diagnosis; but when these symptoms are absent, or only slightly expressed, as often happens, then the difficulty in diagnosing the true state of things is great, and exploration is needed to resolve the doubt.

My own impression is that liver-abscess and dysentery are often co-existent, as is liver-abscess with malarial fever, and that one may follow closely or precede the other; yet the diseases are independent, though, doubtless, they aggravate each other. At the same time, I repeat that, though I regard liver-abscess associated with dysentery as generally merely an independent result of a common cause, yet in some cases, as already explained, I do believe the liver-abscess may be the result of direct absorption of purulent matter, in which case the abscess may be single, double, or multiple, this being a different state of things from the necrotic multiple abscesses which result also from general septicæmic infection. I would summarise the relation of liver-abscess to other disease in the following terms.

1. Pyæmic abscesses which originate in necrosis of parenchyma, embolic deposits, or infarcts; form cavities varying in size from a mere speck to a pea, a walnut, or an orange; they contain debris of tissue, sanies, puriform matter, leucocytes, and, finally, pus. These are seen in various stages of development, and occur in other viscera, as well as in the liver; they are truly septicæmic.

2. A form of liver-abscess coexistent with, and due to, dysentery, is the result of direct absorption of pus or septic matter from the bowel into the liver, through the veins. Such may be solitary, double, or triple; this is also a very dangerous form of the disease, though not necessarily fatal, as it is to be feared must be the case in the former variety.

3. Dysentery, malarial fever, and hepatitis, may coexist with or supervene on each other, as results of climatic influences. In such cases, it seems natural to ascribe the liver-abscess to the dysentery; but it is probable that they are coincidences rather than consequences of each other, and that the cause which affects the glandular structures of the large intestine, as dysentery, may determine, coincidentally, mischief in the liver. Such cases are obviously different from those before mentioned.

4. The ordinary tropical abscess is independent of dysentery.

Each of these forms of liver-abscess may occur in this country in persons who have resided in India or other tropical climates. I should also allude to the co-existence of liver-abscess with diarrhoea. In such cases the diarrhoea may depend directly on the liver-mischief; or it may be kept up by pressure and irritation of the bowel. It may precede the formation of the abscess, which sometimes takes place so insidiously as to escape notice until revealed by some of the symptoms previously described.

A few words now on the symptomatology and treatment of liver-abscess.

The early symptoms of the acute form, either when it occurs in India or here, are enlargement of the liver, pain, and high temperature. The fever is preceded by rigors, or chills, and followed by sweats. There is cough, pain in the right shoulder, inability to lie on the left side, owing to the weight of the swollen and dependent liver, nausea or vomiting, from pressure on the stomach or irritation of the vagus; a yellow tinge of the skin and conjunctivæ (sometimes, but not often, jaundice), and high coloured urine, containing lithates.

These symptoms, together with bulging or pointing, leave little doubt of the presence of abscess, and the diagnosis is not difficult; but there are cases in which the symptoms are not so well marked, and yet an abscess may have formed, and escape detection until sudden bulging and fluctuation become apparent, or until the abscess bursts into the bowels, stomach, or lung, or it may be into the pericardium or peritoneum; in the latter cases with fatal consequences, in the former with great danger from shock or suffocation, but, happily, not often immediately fatal; for such cases not unfrequently recover.

The amount of pain in liver-abscess appears to depend a good deal on the part of the organ implicated. If the pus be deeply seated, at a distance from the peritoneal covering, the pain is comparatively slight, or even absent. If it approach the surface, the pain is severe. There is, in any case, a sense of distress and uneasiness, felt most distinctly when the patient turns on the left side. Tension of the recti abdominis muscles, with a sense of nausea on pressure, is also a characteristic symptom.

It often happens that the patient is suffering at the same time from malarial poisoning, and is liable to recurrences of febrile paroxysms, which make it difficult to decide whether they indicate fever or suppuration; and in some cases, when there has been no previous malarial fever, the rigors or chills of incipient suppuration are so slight as to have escaped notice, and the patient denies having had them at all.

Again, if the abscess be small and central, the pressure is so slight on neighbouring parts that the stomach is neither directly nor sympathetically affected, nor is pain felt in the right shoulder or in the hypochondrium in decubitus on the left side; these signs being either absent or so faintly expressed as to give little aid in diagnosis. In such cases, the patient may be thought to be suffering from malarial cachexia and climatic deterioration of health, till an abscess reveals itself as before described, or it may be that it is not discovered till after death, when an encysted abscess is detected, a contracted cavity containing a certain amount of caseous matter, which reveals the former presence of an abscess, the contents of which have undergone absorption. Such cases are rare, but doubtless they do exist. In these, after years of quiescence, inflammation may light up renewed suppuration, and prove fatal.

The value of the thermometer is great in such cases; it is, perhaps, the most reliable indication we have of the change going on in the liver; the absence of other symptoms making diagnosis so obscure. All one can do in these circumstances is to wait, and watch the temperature and pulse, with the state of the skin and excretions. Occasional sweatings, which are far from being well marked, will tend to establish the probability of suppuration, and exploration may confirm it. I feel certain that such cases occur, and that I have seen them recover.

The treatment in the outset of the acute form is that of congestion; free purgation by mercurials and salines, chloride of ammonium, and ipecacuanha, hot fomentations; and, where the pain is severe and the temperature high, a few leeches on the side or to the ants may be useful. Rest and restricted diet, when pus has formed, favour its evacuation; as soon as possible, make a free opening with antiseptic precautions, and use a drainage-tube. When any doubt exists as to the presence of pus, explore with an aspirating cannula-needle; there is little danger in this proceeding if properly done. Slight inflammation may follow, but it will soon subside, and, even if no pus be detected, it still in some cases seems to do good.

If pus be detected, it should, if possible, be removed, a free opening being made either below the ribs or between them, according to circumstances, and free drainage being ensured through a good sized tube. The strictest antiseptic precautions are to be observed, both during the operation and in subsequent dressings.

I can report most favourably on the results of several cases which have come under my observation since I wrote on the subject some years ago. Where the matter lies deep, and much liver-tissue intervenes, it will be well to introduce a large trocar and cannula, having first made an incision, an inch or an inch and a half in length, through the skin, and leave the cannula in the wound to act as a drainage-tube. This in a short time can be replaced, if necessary, by a flexible tube. After a few days, in favourable cases, the cavity will begin to contract, and the tube will be thrust out, when you may, with advantage, gradually shorten it. The discharge will soon cease to be purulent, assume a serous character, gradually cease altogether, and the wound cicatrises. No special treatment beyond attending to the diet and the state of the bowels is necessary; some simple tonic, with quinine, especially if there be any febrile tendency, may be useful. Quiet and care in avoiding chill are essential until convalescence is well advanced.

The previous history of the case will serve to some extent as a guide to the nature of the abscess; if it be single, and dealt with early, and if no complications arise, the prognosis is often favourable; it may be that another abscess shall form, and still recovery occur. But in cases where multiple abscesses are of the pyæmic or septicæmic character before alluded to, the prospect of success after puncture is very small; indeed, in such cases, the operation would seldom be performed unless one of the abscesses, having attained considerable development, pointed; when relief, though it may be only of a temporary character, would be given by emptying it of its contents.

The question whether liver-abscess be single or multiple is always difficult, and, I fear, one about which it is often impossible at first to form an exact diagnosis, and until puncture has been resorted to. Generally, the evacuation of pus is followed by an immediate fall of temperature, which remains at or near the normal standard, provided the discharge be free and the pus protected from decomposition. A rise in the temperature, or its remaining high with recurrence of pain and sweating, give cause for apprehension either that a fresh abscess is forming, or that decomposition of the discharge is taking place. In either case, you must keep a strict watch on the symptoms, and interfere as early as possible to give exit to newly formed pus, or to remove decomposing discharge from the cavity of the abscess. If the symptoms of suppuration continue, although no physical sign may betray the presence of pus, explore the liver freely with the aspirating needle. I have frequently given exit to fresh collections of pus which have formed under or near the site of the first abscess. As the wound heals, the diet may be increased, but you must protect your patient from all excesses or errors in food.

There are one or two points which you will do well to bear in mind, such as, when operating, the amount of liver-substance you may have to cut through before you reach the pus; this may cause considerable hæmorrhage, but it will generally soon cease. As you wish to avoid loss of blood as much as possible, it is well in these cases to use a cannula sufficiently large to allow the free exit of pus, and leave it in the wound for a few days, substituting for it, after that period, a larger flexible tube, and making a larger opening if necessary.

The pus is often thick, and contains flakes of lymph which may block the tube, and render it necessary to clear it out. Remember that the drainage-tube should not be allowed to press against the wall of the contracting cavity, and that it will require shortening as the wound gradually closes; also the imperative necessity for having free discharge and of keeping it sweet, to which end you must be most scrupulous in attending minutely to the strictest antiseptic precautions. When you are in doubt as to the probable formation of more pus (another abscess), use the exploring aspirating needle; there is no danger in this, if done with due care and precaution, either below or between the ribs.

Should you find another abscess, if possible give exit to its contents by making a free opening. It is very desirable not to delay in these matters. My belief is, that earlier interference would in many cases have saved life and diminished suffering.

In any case a liver-abscess is a most serious affair, and the result is uncertain; but, by observing and acting on the principles and practice inculcated, you may hope to meet with a considerable amount of success.

Never allow a liver-abscess to point and cause gangrene of the abdominal wall from pressure, and consequent inevitable death by exhaustion, in such cases where a simple incision would, in all probability, have relieved suffering, and might have saved life.

NOTE.—The patient who was the subject of this lecture was doing well in all respects on June 2nd, 1884.

BEQUESTS AND DONATIONS.—The Royal Free Hospital has received £1,000, and the National Hospital for Consumption at Ventno £600, under the will of Mrs. Esther Bunning.—Mr. George William Nugent, of Cambridge House, Maida Vale, has bequeathed £200 each to the Seamen's Hospital, Greenwich, the London Hospital, St. Mark's Hospital for Fistula, etc., the Royal Free Hospital, the Metropolitan Free Hospital, the Western General Dispensary, St. Mary's Hospital, the Hospital for Sick Children, the Female Lock Hospital and Asylum, the Middlesex Hospital, University College Hospital, and the Convalescent Hospital, Stone, Staffordshire.—Miss Mary Davidson Blair, of Forbes Road, Greenhill, Edinburgh, has bequeathed £100 to the Society for the Relief of the Destitute Sick, Edinburgh, £50 to the Smyrna Medical Mission, and £50 to the Longmore Hospital for Incurables.—The Lincoln County Hospital has received £200 under the will of Mr. T. Garfit, of Boston, and £50 under that of Miss Anne Torr, of Grimsby.—Mr. William John Campbell Allen, of Fannoran, Carrickfergus, has bequeathed £100 to the Royal Hospital, Belfast.—The Grocers' Company have given £100 to the Royal Medical Benevolent College.—The Hon. Mrs. Robert Bruce has given £50 to the Charing Cross Hospital, and £50, additional, to the Middlesex Hospital.—The Royal Hospital for Diseases of the Chest has received £43 10s. from Captain Wiltshire, being the proceeds of a concert at Myddelton Hall, Islington, on April 30th.

CLINICAL REMARKS

CASES ILLUSTRATING THE ESSENTIAL IDENTITY OF PROGRESSIVE MUSCULAR ATROPHY AND PROGRESSIVE BULBAR PARALYSIS.¹

By J. MAGEE FINNY, M.D. Univ. Dubl., F.K.Q.C.P.I.,
King's Professor of Practice of Medicine in the School of Physic; Clinical Physician to Sir Patrick Dun's Hospital.

CASES of primary labio-glosso-laryngeal paralysis are of sufficient rarity to secure for the recital of their clinical features some interest; for, although bulbar symptoms are not very uncommonly present at the termination of various systemic cerebral and spinal diseases, examples of primary bulbar paralysis are not frequently met with in the experience of physicians in general practice; and it would seem, indeed, to be among the rarer features of nerve-paresis witnessed by those whose opportunities of studying neurotic lesions are extensive, if we may judge from the fact that Erb, when writing the article bearing directly on this subject (Ziemssen's *Cyclopædia of Medicine*), refers to but nine cases of which he had accurate notes, and Hammond states that he had met with but six.

With the exception of the two individuals, the record of whose state I am about to submit to the Academy, and of a third case which I saw when a senior student, but which, I confess, I failed at the time to recognise, I have not met with any other instances in my own practice, nor seen any in the wards of the hospital under any of my colleagues.

Apart from the intrinsic interest attaching to the pathological lesion of the medulla to which is due the disease termed labio-glosso-laryngeal paralysis, or, as Kussmaul, with special reference thereto, called it, "progressive bulbo-nuclear paralysis," I look upon these cases, the subject of this communication, as illustrating, in a marked way, the intimate relations which exist between it and progressive muscular atrophy.

Each of these diseases is to be met with separately. Although chronic progressive bulbar paralysis does occur alone, as a typical disease, unaccompanied by any general amyotrophic changes of the trunk and extremities—as, of Erb's nine cases, six were limited to paralysis of the lips, tongue, and larynx—yet it is proved to be one of the most frequent complications of progressive muscular atrophy. A striking example of this has been recently published by Dr. W. Fox (BRITISH MEDICAL JOURNAL, 1881, vol. i, p. 82). It is fairly formulated by Eulenburg (Ziemssen's *Cyclopædia of Medicine*, vol. xiv, p. 132). "One cause of death in progressive muscular atrophy is so frequent, and is clinically so nearly allied to progressive muscular atrophy, that it must be considered not as an accidental complication, but as a combined disease arising from the same or similar pathogenic causes. This is progressive bulbar paralysis. It may form a cause of death, either by producing a gradual palsy of the muscles of swallowing and of the larynx, or, when it occurs in an apoplectic form, with immediate palsy of the respiratory centres."

In like manner, in cases of primary bulbar paralysis, one of the commonest complications is that of progressive muscular atrophy. Hence the question presses as to the relative position these two lesions bear to one another. Is the fact of their being both present in an individual to be attributed to chance coincidence? Is one sequential on the other? Or are the two groups of symptoms but evidences of the same disease affecting different but allied parts of the cerebro-spinal centres—branches of the same tree which, though springing from different parts of the parent stem, produce similar fruits?

When glosso-laryngeal disease was for the first time systematically demonstrated by Duchenne in 1861, although it had been recognised twenty years previously by Trouseau, the close relation it bore to progressive muscular atrophy was thoroughly admitted by this accurate observer.

Duchenne considered, however, the two lesions to be quite distinct, the one from the other, and laid down the characteristic feature of progressive bulbar paralysis to be "paralysis without atrophy," in contrast to that of progressive muscular atrophy, which was "atrophy without paralysis." For several years subsequently this view of Duchenne's was generally adopted, and even in the present time it appears to have its adherents, among whom Hammond and Friedreich may be numbered.

¹ Read before the Academy of Medicine in Ireland.