August 3rd.—The patient was dressed, and sitting up. The head could be placed quite straight without pain.

August 4th (third day after operation).—The boy recommenced the medical gymnastic treatment. The two sterno-mastoids were of exactly the same length. The patient could hold the head perfectly straight on paying attention.

August 7th.—The left muscle contracted a little while the boy was lying on his back, on attempting to raise the head from off the couch (flexion of the neck).

August 9th.—The divided muscle had healed sufficiently to contract rigorously, although only the ninth day after the operation.

August 30th.—The patient ceased treatment. The spine was quite straight. The two sterno-mastoids remained of equal length. A year afterwards, there had not been any relapse, and the head remained quite straight.

From the above notes, it will be observed that I employ no apparatus whatever, either before or after the operation, and that I do not insist upon any rubbing of the affected muscle. A couch with movable back and horizontal seat (of which I gave an illustration in the *BRITISH MEDICAL JOURNAL* of March 5th, 1881) is useful, on which the patient is trained to hold his head in the best possible position. Walking about is allowed as usual during the whole treatment, except during the two days immediately after the tenotomy.

The results in these three cases will compare favourably, I believe, with any published, especially in regard to the rapidity with which a complete cure was obtained, and with no tendency to relapse a year or more afterwards.

The treatment I here venture to bring forward is based on rational surgical principles, and I trust that other surgeons may be induced to try it. It consists essentially of (1) a week's preliminary daily training of the patient's neck-muscles; (2) tenotomy; and (3) two or three weeks' daily further perseverance with prescribed medical gymnastic exercises, and a re-education of the patient's muscular sense as to the proper position of the head, without the employment of any mechanical contrivances throughout the treatment.

PATHOLOGY, SYMPTOMS, AND TREATMENT OF CELLULITIS OF THE NECK.¹

By G. HARRISON YOUNGE, L.K.Q.C.P.I., etc.,
Surgeon Army Medical Department.

STRANGE to say, cellulitis of the neck, although a most important and dangerous affection, is not even mentioned in the ordinary text-books on surgery. Yet a glance at the anatomy of the neck is sufficient to show how important is this disease. When inflammation of the cellular tissue of the neck takes place, serum and pus are effused under the cervical fascia, giving rise to severe and dangerous pressure, not only on the larynx and trachea, pharynx and œsophagus, but also on the large blood-vessels passing to and from the brain, and on the sympathetic and pneumogastric nerves. As Mr. H. G. Croly of Dublin says, when writing of this affection, "Diffuse inflammation of the areolar tissue of the neck is one of the most serious and fatal forms of disease which the erysipelatos type assumes."

Causes.—The causes of cellulitis of the neck may be divided into predisposing and exciting, the former being the same as in all other erysipelatos inflammations. In fact, this disease may be predisposed to by anything that tends to lower the general health, such as want of attention to hygienic conditions, intemperance, want of proper food, and also the opposite extreme, namely, overfeeding combined with deficient exercise.

Diseased states of the blood, especially those occurring in Bright's disease and in the gouty diathesis, predispose to erysipelas, and therefore to this disease. In many cases, however, the disease comes on suddenly in strong healthy persons, without the previous existence of any of these conditions. The exciting cause is always exposure to wet and cold. The disease, therefore, almost always occurs in winter, and is most prevalent in low damp situations. Cellulitis of the neck, unlike other erysipelatos diseases, is rarely, if ever, traumatic, and is, probably, never due to infection.

Symptoms.—The symptoms of the disease are always well marked and characteristic. For a few days before the local symptoms become established, the patient complains of general malaise, there are frequent chills, with a feeling of great languor and inability for all exercise. A dull aching pain in the back is almost invariably complained of. At the same time there is severe headache; the tongue becomes thickly furred; there is nausea, and occasionally vomiting, and the bowels are obstinately constipated. The temperature is always above normal, but

rarely exceeds 102° Fahr. After the general symptoms have lasted from twenty-four to forty-eight hours, the local symptoms set in. The patient now complains of severe pain in the neck; there is some pain and difficulty in swallowing, especially solids, and there is a clear, ringing, metallic cough, very like that heard in the first stage of croup. Then the neck becomes swollen, the swelling extending from one angle of the jaw to the other, and then downwards along the sterno-cleido-mastoid muscles to the clavicle. In a small number of cases the swelling is unilateral, and then extends from the angle of the jaw downwards to the inner end of the clavicle. The right side is, I believe, invariably affected when the disease is unilateral. The swelling is at first tense and elastic, but, after about twenty-four hours, it becomes soft, has a brawny feel, and pits on pressure. The neck has now a most characteristic appearance. When looked at from the front, it is seen to be considerably swollen from the jaw down to the clavicle, while the centre of the swelling presents a horizontal depression corresponding to the attachment of the cervical fascia to the os hyoides. This gives to the swelling an appearance exactly as if a muff had been twice folded round the neck. When this brawny feeling of the tissues is established, the symptoms of the disease become greatly intensified, as now effusion of serum and pus has taken place beneath the deep cervical fascia, and is exercising injurious, and if not relieved fatal, pressure on the deep structures of the neck.

There is now great difficulty of breathing, in bad cases amounting to orthopnoea, while the voice is a mere whisper, or there may be complete aphonia. The patient is entirely unable to swallow solids, and occasionally even fluids, these regurgitating through the nose when attempts are made to swallow them. The pressure on the veins of the neck retards the return of the venous circulation. The face, therefore, becomes congested and livid, while cold clammy drops of sweat collect on the forehead. At the same time congestion of the brain takes place, giving rise to great anxiety, restlessness, and occasionally to delirium.

In cases where the effusion extends underneath the sterno-cleido-mastoid muscles, serious pressure is exercised on the sympathetic and pneumogastric nerves, which greatly increases the dyspnoea, and gives rise, at first, to violent and irregular action of the heart, and afterwards to great weakness of this organ.

If, when effusion has taken place, the tension be not relieved by free and deep incisions into the cervical fascia, œdema of the glottis may set in and prove rapidly fatal; or the dyspnoea may gradually increase, and the patient die comatose from carbonic acid poisoning; or, what still more frequently happens, sudden spasm of the glottis may occur, the result being suffocation. Even if these dangers be avoided, deep and destructive sloughing of the tissues of the neck will take place, if the matter be not allowed free vent.

There is never any difficulty in diagnosing this affection. The characteristic appearance of the neck, the brawny feel of the tissues, and the subsequent pitting on pressure, distinguish it from every other disease found in this region. In this disease, although there is abundant effusion of serum and pus, fluctuation is never obtained. This is due, I believe, to the fact that the effusions are always situated under the deep cervical fascia, which binds them down so firmly that we cannot detect fluctuation.

Pathology.—This disease consists essentially of an erysipelatos inflammation of the areolar tissue, which fills up the spaces between, and connects together the deep cervical muscles. This tissue rapidly breaks down, giving rise to sloughs, while at the same time serum and pus are effused into the intermuscular spaces. As nearly all this mischief takes place beneath the deep cervical fascia, which is dense and unyielding, we can at once see what an amount of pressure is exercised on the important anatomical structures of the neck, and how necessary it is to give relief to this pressure by free incisions.

Treatment.—Of all the diseases to which man is subject, none require more prompt treatment than cellulitis of the neck; for if relief be not afforded by surgical means, the disease is almost necessarily fatal, or if recovery do chance to take place without operation, it will only be after sloughing has set in, and extensive disorganisation of the muscles of the neck has taken place, which will give rise to stiffness and great deformity. From the first, our treatment should be such as to support the strength of the patient, and all remedies which tend to lower the powers of the system should be carefully avoided. It is to be remembered that the patient is suffering from a disease of a highly asthenic type, and will also have to pass through a period of prolonged suppuration before recovery takes place. Lowering treatment also tends to increase the chances of pyæmia occurring. The patient should have a good purgative at the commencement of the disease, as the bowels are always constipated. He should then be placed on a mixture containing chlorate of potash combined with tincture of the perchloride of

¹ Read before the Bengal Branch of the British Medical Association.

iron, or nitric acid with decoction of cinchona. It seems to me that a combination of chlorate of potash with tincture of iron has a peculiar and most beneficial effect in all erysipelatous diseases, and I now invariably prescribe it. At the same time, plenty of milk, beef-tea, chicken-broth, and raw eggs and brandy should be given. Overstimulation, however, is to be carefully avoided, and the quantity of brandy given should be regulated by the amount of asthenia present. If the patient cannot swallow these articles they should be given in the form of enemata. If given thus, the benefit to the patient will be greatly increased by the addition to each enema of a little pepsine and hydrochloric acid about half an hour before it is administered. Local applications, as large linseed-poultices, hot fomentations, and poppy-stupes, should be applied, as they relieve pain and tension. However, all these measures are of small importance in comparison to the surgical treatment, as they have little or no curative power over the disease, and if they alone be used the patient will probably die. As Mr. H. G. Croly says, "The symptoms are urgent and alarming, and if relief be not afforded by prompt and bold surgical aid, the patient may be suffocated."

As soon as the brawny feeling of the tissues, with pitting on pressure, sets in, free and deep incisions should be made into the cervical fascia. When these incisions have been made, the relief which they give is immediate, even though no serum or pus escape; as the cervical fascia, being divided, stretches and tears, and the tension is thus greatly lessened. I have seen patients who had almost complete aphonia and dysphagia able to speak and swallow in five minutes after incisions had been made, and this even though no fluid escaped.

To illustrate the suddenness of the relief afforded I may mention a case which I treated at Fort Pitt, Chatham. The patient was admitted into hospital in January 1882, complaining of general febrile

symptoms and slight sore throat. On the evening of his admission he was seized with a severe rigor and pain in the right side of his neck. The symptoms and appearances above described soon manifested themselves, only in this case the disease was entirely confined to the right side of the neck. About forty-eight hours after the swelling of the neck began I was called to see him, and found that violent dyspnoea had suddenly set in. On examination, it was found that the swelling had greatly increased, and had pushed the larynx considerably to the left side. As the patient was in imminent danger of suffocation, an incision was at once made through the skin one inch to the outer side and one inch above the inner articular end of the clavicle, where

the swelling seemed to bulge most. The cervical fascia was then divided on a director, and a large quantity of sanious pus at once escaped. The man expressed immediate relief, the voice became clear, and he drank some beef-tea in a few minutes afterwards.

In making the incisions, it is only necessary to remember the anatomy of the neck to do so with perfect safety. On the accompanying sketch, which is taken from one by Mr. Croly, in his pamphlet on *Cellulitis of the Neck*, the lines of safety and danger are well shown. The lines of danger are shown by the black lines, while those of safety are shown by the dotted lines. The line extending from the angle of the jaw to the centre of the clavicle represents the course of the external jugular vein; while the curved line, along the inner border of the sterno-mastoid muscle, shows the course of the carotid vessels. The lines of safety will be seen to extend along the mesial line from the chin to the interclavicular notch, and along the posterior border of the sterno-mastoid muscle—taking care, however, to avoid the spot where the muscle is crossed by the external jugular vein. In making the incisions, it is best to adopt the method recommended by Hilton—viz., to make an incision through the skin with a scalpel; and then, having passed an ordinary pair of dissecting-forceps through the cervical fascia, forcibly separate their blades, and thus tear through this structure. In the supra-hyoid region, however, along the mesial line, this is not necessary, as here the scalpel can be passed to the full length of its blade without danger.

Before concluding, I would refer to a variety of this disease, first described by the late Dr. Osbrey of Dublin, and which is nearly always fatal—namely, that occurring in children recovering from malignant scarlatina. In these cases, the disease sets in very insidiously, and is invariably followed by sloughing. Dr. Osbrey believed it was not good practice to make incisions into the cervical fascia in these cases, as the patients are not able to bear them, and they, in his opinion, only hasten a fatal termination. These cases are to be treated with chlorate of potash and tincture of iron, together with all the nourishment the patient can be got to take.

NOTES OF A CASE OF GONORRHOEA, WITH ACUTE RHEUMATISM AND ENDOCARDITIS OCCURRING IN ITS COURSE.

By T. C. RAILTON, M.D. LOND., B.Sc., F.R.C.S.,

Senior Assistant-Physician to the Manchester Clinical Hospital for Women and Children.

THE following case, which I shall endeavour to report as briefly as possible, is of considerable interest, and deserves to be recorded on account of the rarity with which gonorrhoeal rheumatism is complicated with endocarditis. Rosenstein, in Ziemssen's *Cyclopaedia*, vol. vi, almost denies the occurrence of endocardial complication in this form of rheumatism; while Senator, in vol. xvi of the same work, asserts that, if more than two or three joints be affected, if there be abundant sweating and affection of the heart, we have not to deal with gonorrhoeal rheumatism at all, but with a case of acute rheumatism, accidentally associated with a gonorrhoea. It is difficult to see why this so-called gonorrhoeal rheumatism, very similar in its character to the forms following scarlet fever or childbirth, in which endocarditis is by no means of uncommon occurrence, should differ from them in any special immunity from valvular complications, or in any of the other clinical features. In all three, there appears to be reason for assigning their origin to something akin to pyæmia.

J. R. C., aged 21, bookbinder, was first seen on September 17th, on account of a sore-throat. His previous and family-history were as follows. He had never suffered from rheumatism in any form, nor from scarlet fever. He had always enjoyed good health. His parents were both dead from phthisis, neither of them having had rheumatism to his knowledge. About the beginning of August, he contracted gonorrhoea for the first time in his life; for this he had an injection prescribed, which caused the discharge to diminish. On the 25th, he discontinued using it, thinking he was cured; but he still noticed the end of the urethra to be occasionally glued up, and a drop of pus to escape when the orifice was pressed open.

Without having been particularly exposed to cold or wet previously, on September 15th he began to feel ill. He had repeated shiverings, aching in his limbs, and sore-throat, and felt feverish.

On the 16th, his throat became worse; swallowing was painful and difficult, and his head ached excruciatingly.

On the 17th, when seen for the first time, his tonsils were swollen, and intensely red; his headache was so severe, that he said it almost blinded him. His temperature was only moderately high (102° Fahr.).